

Gossypium hirsutum

Source: AGRICOLA database (1970-1996)

References (Biological Abstracts 1988-2000):

Agrawal, A. A. (2000). Host-range evolution: Adaptation and trade-offs in fitness of mites on alternative hosts. *Ecology* Washington D C. [print] February 81(2): 500-508. {a} Department of Botany, University of Toronto, 25 Wilcocks Avenue, Toronto, ON, M5S 3B2, Canada

Agrawal, A. A., R. Karban, et al. (2000). How leaf domatia and induced plant resistance affect herbivores, natural enemies and plant performance. *Oikos*. April 89(1): 70-80. {a} Dept of Botany, Univ. of Toronto, 25 Willcocks Street, Toronto, ON, M5S 3B2, Canada

Ahuja, S. L. and S. K. Banerjee (2000). Stability for bollworm resistance, jassid grade, seed cotton yield and its components of cytotypes in cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research*. [print] June 34(2): 71-77. {a} Central Institute for Cotton Research Regional Station, Sirsa, 125055, India

Anadranistakis, M., A. Liakatas, et al. (2000). Crop water requirements model tested for crops grown in Greece. *Agricultural Water Management*. [print] August 45(3): 297-316. {a} Agricultural University of Athens, 75 Iera Odos, GR-118 55, Athens, Greece

Andersland, J. M. and B. A. Triplett (2000). Selective extraction of cotton fiber cytoplasm to identify cytoskeletal-associated proteins. *Plant Physiology and Biochemistry* Paris. March 38(3): 193-199. {a} ARS, Southern Regional Research Center, USDA, 1100 Robert E. Lee Blvd., New Orleans, LA, 70124-4305, USA

Ashraf, M. and S. Ahmad (2000). Influence of sodium chloride on ion accumulation, yield components and fibre characteristics in salt-tolerant and salt-sensitive lines of cotton (*Gossypium hirsutum* L.). *Field Crops Research*. May 66(2): 115-127. {a} ABC Road, 51-C Sheikh Colony, Faisalabad, Pakistan

Baird, R., W. Batson, et al. (2000). First report of *Rhizoctonia solani* AG-7 on cotton in Mississippi. *Plant Disease*. [print] October 84(10). {a} Entomology and Plant Pathology Department, Mississippi State University, Mississippi State, MS, 39762, USA

Banerjee, A. K., D. C. Agrawal, et al. (2000). Recovery of in vitro cotton shoots through micrografting. *Current Science* Bangalore. [print] 78(5): 623-626. {a} Plant Tissue Culture Division, National Chemical Laboratory, Pune, 411 008, India

Banumathy, S. and S. Patil (2000). Genetic analysis of fuzzlessness and other attributes in *Gossypium hirsutum* L. *Crop Research* Hisar. [print] July 20(1): 102-104. {a} Centre for Plant Breeding and Genetics, Tamil Nadu Agricultural University, Coimbatore, 641 003, India

Bararpour, M. T. and A. Abdollahi (2000). Velvetleaf (*Abutilon theophrasti*) interference and control. *Zeitschrift fuer Pflanzenkrankheiten und Pflanzenschutz*(17): 589-594. {a} Department of Agronomy, University of Mazandaran, Sari, Iran

Barbour, M. M. and G. D. Farquhar (2000). Relative humidity- and ABA-induced variation in carbon and oxygen isotope ratios of cotton leaves. *Plant Cell and Environment*. [print] May 23(5): 473-485.

Bauer, P. J., J. R. Frederick, et al. (2000). Canopy photosynthesis and fiber properties of normal- and late-planted cotton. *Agronomy Journal*. [print] May June 92(3): 518-523. {a} Coastal Plains Soil, Water, and Plant Res. Cent., USDA-ARS, 2611 W. Lucas St., Florence, SC, 29501-1242, USA

Baumhardt, R. L. and R. J. Lascano (2000). Water budget and yield of dryland cotton intercropped with terminated winter wheat. *Agronomy Journal*. Nov. Dec. 91(6): 922-927. {a} Conservation and Production Res. Lab., USDA-ARS, Bushland, TX, 79012-0010, USA

Bednarz, C. W., D. C. Bridges, et al. (2000). Analysis of cotton yield stability across population densities. *Agronomy Journal*. [print] January February 92(1): 128-135. {a} Coastal Plain Experiment Station, University of Georgia, Tifton, GA, 31793, USA

Bednarz, C. W., G. H. Harris, et al. (2000). Agronomic and economic analyses of cotton starter fertilizers. *Agronomy Journal*. [print] July August 92(4): 766-771. {a} Coastal Plain Experiment Station, Univ. of Georgia, Tifton, GA, 31793, USA

Beyrouthy, C. A., J. K. Keino, et al. (2000). Phytotoxic concentrations of subsoil aluminum as influenced by soils and landscape position. *Soil Science*. Feb. 165(2): 135-143. {a} Department of Crop, Soil, and Environmental Sciences, University of Arkansas, 115 Plant Science Bldg., Fayetteville, AR, 72701, USA

Booker, F. L. (2000). Influence of carbon dioxide enrichment, ozone and nitrogen fertilization on cotton (*Gossypium hirsutum* L.) leaf and root composition. *Plant Cell and Environment*. [print] June 23(6): 573-583. {a} Agricultural Research Service, Air Quality-Plant Growth and Development Research Unit, and Department of Crop Science, United States Department of Agriculture, North Carolina State University, 3908 Inwood Road, Raleigh, NC, 27603, USA

Booker, F. L., S. R. Shafer, et al. (2000). Carbon dioxide enrichment and nitrogen fertilization effects on cotton (*Gossypium hirsutum* L.) plant residue chemistry and decomposition. *Plant and Soil*. [print] 220(1-2): 89-98. {a} Air Quality - Plant Growth and Development Research Unit, and Department of Crop Science, U.S. Department of Agriculture, Agricultural Research Service, North Carolina State University, 3908 Inwood Road, Raleigh, NC, 27603, USA

Borole, V. K., D. B. Dhumale, et al. (2000). Embryo culture studies in interspecific crosses between arboreum and hirsutum cotton. *Indian Journal of Genetics and Plant Breeding*. Feb. 60(1): 105-110. {a} Department of Agricultural Botany, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola, 444 104, India

Briddon, R. W., S. Mansoor, et al. (2000). Clones of cotton leaf curl geminivirus induce symptoms atypical of cotton leaf curl disease. *Virus Genes*. Feb. 20(1): 19-26. {a} Department of Virus Research, John Innes Centre, Norwich Research Park, Colney Lane, Norwich, NR4 7UH, UK

Cantrell, R. G. and D. D. Davis (2000). Registration of NM24016, an interspecific-derived cotton genetic stock. *Crop Science*. [print] July August 40(4). {a} Dep. Agronomy and Horticulture, New Mexico State Univ., Las Cruces, NM, 88003-8003, USA

Cantrell, R. G., C. L. Roberts, et al. (2000). Registration of 'Acala 1517-99' cotton. *Crop Science*. [print] July August 40(4): 1200-1201. {a} Dep. Agronomy and Horticulture, New Mexico State Univ., Las Cruces, NM, 88003-8003, USA

Chen, C. x., Y. j. Yu, et al. (2000). Genetic analysis of salt-tolerance variant in cotton. *Xibei Zhiwu Xuebao*. [print] March 20(2): 234-237. {a} Laboratory of Plant Genetic Engineering, Shandong Agricultural University, Taian, Shandong, 271018, China

Chimbekujwo, I. B. (2000). Frequency and pathogenicity of fusarium wilts (*Fusarium solani* and *Fusarium equiseti*) of cotton (*Gossypium hirsutum*) in Adamawa in Nigeria. *Revista de Biologia Tropical*. [print] March 48(1): 1-5. {a} Department of Biological Sciences, Federal University of Technology, Yola, Adamawa State, Nigeria

Chipika, J. T. and G. Kowero (2000). Deforestation of woodlands in communal areas of Zimbabwe: Is it due to agricultural policies? *Agriculture Ecosystems and Environment*. [print] July 79(2-3): 175-185. {a} Center for International Forestry Research (CIFOR), Regional Office, c/o Institute of Environmental Studies, University of Zimbabwe, Harare, Zimbabwe

Chu, C. C., T. Freeman, et al. (2000). Bemisia argentifolii adult, nymph and egg densities and egg distribution on selected upland cottons. *Journal of Entomological Science*. Jan. 35(1): 39-47. {a} Western Cotton Research Laboratory, USDA, ARS, Phoenix, AZ, 85040, USA

- Colson, H. E. S. and B. J. Deverall (2000). Effect of 2,6-dichloroisonicotinic acid, its formulation materials and benzothiadiazole on systemic resistance to alternaria leaf spot in cotton. *Plant Pathology Oxford*. April 49(2): 171-178. {a} Department of Crop Sciences, University of Sydney, Sydney, NSW, 2006, Australia
- Crow, W. T., D. W. Dickson, et al. (2000). Yield reduction and root damage to cotton induced by *Belonolaimus longicaudatus*. *Journal of Nematology*. [print] June 32(2): 205-209. {a} Texas A and M University Agricultural Research and Extension Center-Dallas, 17360 Coit Road, Dallas, TX, 75252, USA
- Crow, W. T., D. P. Weingartner, et al. (2000). Population dynamics of *Belonolaimus longicaudatus* in a cotton production system. *Journal of Nematology*. [print] June 32(2): 210-214. {a} Texas A and M University Agricultural Research and Extension Center-Dallas, 17360 Coit Road, Dallas, TX, 75252, USA
- Cui, Y., A. A. Bell, et al. (2000). Expression of potential defense response genes in cotton. *Physiological and Molecular Plant Pathology*. Jan. 56(1): 25-31. {a} Dept. of Plant Pathology and Microbiology, Texas A and M University, College Station, TX, 77843-2132, USA
- Culpepper, A. S. and A. C. York (2000). Weed management in ultra narrow row cotton (*Gossypium hirsutum*). *Weed Technology*. Jan. March 14(1): 19-29. {a} Department of Agronomy, University of Georgia, Tifton, GA, 31793, USA
- De, B. P. J., P. J. Hart, et al. (2000). The biology of two *Eretmocerus* spp. (*Haldeman*) and three *Encarsia* spp. Forster and their potential as biological control agents of *Bemisia tabaci* biotype B in Australia. *Entomologia Experimentalis et Applicata*. Jan. 94(1): 93-102. {a} CSIRO Entomology, Indooroopilly, QLD, 4068, Australia
- Dixon, D. C., W. R. Meredith, Jr., et al. (2000). An assessment of alpha-tubulin isotype modification in developing cotton fiber. *International Journal of Plant Sciences*. Jan. 161(1): 63-67. {a} Southern Regional Research Center, USDA-ARS, 1100 Robert E. Lee Boulevard, New Orleans, LA, 70179-0687, USA
- Dubey, B. K., J. Gupta, et al. (2000). Sulphur response in cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences*. [print] April 70(4): 253-254. {a} Lucknow University, Lucknow, Uttar Pradesh, 226 002, India
- Feng, R. and R. M. Brown, Jr. (2000). A novel cotton ovule culture: Induction, growth, and characterization of submerged cotton fibers (*Gossypium hirsutum* L.). *In Vitro Cellular and Developmental Biology Plant*. [print] July 36(4): 293-299. {a} Molecular Genetics and Microbiology, University of Texas at Austin, Austin, TX, 78712, USA
- Grossmann, K. (2000). Involvement of ethylene in the induction of cotton defoliation by the isoindoldione cinidon-ethyl. *Pesticide Biochemistry and Physiology*. [print] September 68(1): 42-48. {a} BASF Agricultural Center Limburgerhof, D-67114, Limburgerhof, Germany
- Gupta, G. P., G. K. Mahapatro, et al. (2000). Bio-potency of insecticidal crystal proteins of *Bacillus thuringiensis* against cotton (*Gossypium hirsutum*) bollworms. *Indian Journal of Agricultural Sciences*. March 70(3): 194-196. {a} Indian Agricultural Research Institute, New Delhi, 110 012, India
- Gupta, S. K., P. K. Singh, et al. (2000). Effect of light intensity on in vitro multiple shoot induction and regeneration of cotton (*Gossypium hirsutum* L. cv Khandawa-2). *Indian Journal of Experimental Biology*. [print] April 38(4): 399-401. {a} National Botanical Research Institute, Rana Pratap Marg, Lucknow, 226 001, India
- Hanson, R. E., F. M. N. Islam, et al. (2000). Ty1-copia-retrotransposon behavior in a polyploid cotton. *Chromosome Research* 8(1): 73-76. {a} Department of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843-2474, USA
- He, J. X., R. Q. Jiang, et al. (2000). Genetic analysis for introgressive genes of petal color from *Gossypium bickii* to *G. hirsutum* L. *Acta Genetica Sinica*. [print] 27(4): 344-350. {a} Institute of Genetics, Chinese Academy of Sciences, Beijing, 100101, China

Howell, C. R., L. E. Hanson, et al. (2000). Induction of terpenoid synthesis in cotton roots and control of *Rhizoctonia solani* by seed treatment with *Trichoderma virens*. *Phytopathology* . March 90(3): 248-252. {a} Southern Crops Research Laboratory, Agriculture Research Service, United States Department of Agriculture, 2765 F and B Road, College Station, TX, 77845, USA

Hulugalle, N. R. (2000). Carbon sequestration in irrigated Vertisols under cotton-based farming systems. *Communications in Soil Science and Plant Analysis*. March 31(5-6): 645-654. {a} Cooperative Research Centre for Sustainable Cotton Production, NSW Agriculture, Australian Cotton Research Institute, Narrabri, NSW, 2390, Australia

Jiang, C., R. J. Wright, et al. (2000). QTL analysis of leaf morphology in tetraploid *Gossypium* (cotton). *Theoretical and Applied Genetics*. Feb. 100(3-4): 409-418. {a} Department of Crop and Soil Science, University of Georgia, Athens, GA, 30602, USA

Jiang, C. x., P. W. Chee, et al. (2000). Multilocus interactions restrict gene introgression in interspecific populations of polyploid *Gossypium* (cotton). *Evolution* . [print] June 54(3): 798-814. {a} Garst Seeds, Slater, IA, 50244, USA

Johnson, A. W., C. C. Dowler, et al. (2000). Population dynamics of *Meloidogyne incognita*, *M. arenaria*, and other nematodes and crop yields in rotations of cotton, peanut, and wheat under minimum tillage. *Journal of Nematology*. March 32(1): 52-61. {a} Coastal Plain Station, USDA ARS, Tifton, GA, 31793, USA

Jost, P. H. and J. T. Cothren (2000). Growth and yield comparisons of cotton planted in conventional and ultra-narrow row spacings. *Crop Science*. [print] May June 40(2): 430-435. {a} Department of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843-2474, USA

Katkar, R. N. (2000). Effect of intercropping on yield of cotton. *Crop Research Hisar*. [print] May 19(3): 409-413. {a} Cotton Research Scheme, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola, 444 104, India

Klueva, N. Y., R. C. Joshi, et al. (2000). Genetic variability and molecular responses of root penetration in cotton. *Plant Science Shannon*. [print] June 12th 155(1): 41-47. {a} Department of Plant and Soil Science, Texas Tech University, Lubbock, TX, 79409-2122, USA

Koenning, S. R., F. F. A. Alim, et al. (2000). Stem canker on cotton caused by *Phoma exigua* in North Carolina and Virginia. *Plant Disease*. [print] November 84(11). {a} Department of Plant Pathology, North Carolina State University, Raleigh, NC, 27695-7616, USA

Korolev, N., J. Katan, et al. (2000). Vegetative compatibility groups of *Verticillium dahliae* in Israel: Their distribution and association with pathogenicity. *Phytopathology* . May 90(5): 529-536. {a} Department of Plant Pathology, ARO, Volcani Center, Bet Dagan, 50250, Israel

Kumaresan, D., J. Ganesan, et al. (2000). Genetic analysis of quantitative characters in cotton (*Gossypium hirsutum* L.). *Crop Research Hisar*. [print] May 19(3): 481-484. {a} Department of Agricultural Botany, Annamalai University, Chidambaram, Tamil Nadu, India

Larik, A. S., S. I. Malik, et al. (2000). Assessment of heritability and genetic advance for yield and yield components in *Gossypium hirsutum* L. *Scientific Khyber*. Jan. 13(1): 39-44. {a} Agriculture Development Institute, Khanpur Usta Muhammad, Pakistan

Lawrence, C. and A. S. Holaday (2000). Effects of mild night chilling on respiration of expanding cotton leaves. *Plant Science Shannon*. [print] August 157(2): 233-244. {a} Department of Biological Sciences, Texas Tech University, Lubbock, TX, 79409-3131, USA

Lin, T. B., S. Wolf, et al. (2000). Silverleaf whitefly stress impairs sugar export from cotton source leaves. *Physiologia Plantarum*. [print] July 109(3): 291-297. {a} Faculty of Agricultural, Food and Environmental Quality Sciences, The Hebrew University of Jerusalem, Rehovot, 76100, Israel

- Liu, X. z., S. z. Kang, et al. (2000). Effects of soil moisture and short-term shade on photosynthetic and stomatal response in cotton leaves. *Xibei Zhiwu Xuebao*. [print] July 20(4): 561-567. {a} Institute of Soil and Water Conservation, Ministry of Water Resources, Chinese Academy of Sciences, Yangling, Shaanxi, 712100, China
- Maitra, S., A. K. Mondal, et al. (2000). Effect of spacing and fertility levels on upland cotton (*Gossypium hirsutum*) in Sundarbans. *Indian Journal of Agricultural Sciences*. [print] May 70(5): 323-324. {a} R K Ashram Krishi Vigyan Kendra, Nimpith Ashram, WB, 743 338, India
- Mangrich, M. E. and M. E. Saltveit (2000). Heat shocks reduce chilling sensitivity of cotton, kenaf, okra, and rice seedling radicles. *Journal of the American Society for Horticultural Science*. [print] May 125(3): 377-382. {a} Mann Laboratory, Department of Vegetable Crops, University of California, One Shields Ave., Davis, CA, 95616-8631, USA
- Mehta, Y. R. and R. L. Brogin (2000). Phytotoxicity of a culture filtrate produced by *Stemphylium solani* of cotton. *Plant Disease*. [print] August 84(8): 838-842. {a} Instituto Agronomico do Parana-IAPAR, CEP-8600-970, Londrina, PR., Brazil
- Murugan, K., N. S. Kumar, et al. (2000). Influence of *Helicoverpa armigera* (Hubner) diet on its parasitoid *Campoletis chloridae* Uchida. *Insect Science and its Application*. [print] January March 20(1): 23-31. {a} Division of Entomology, Department of Zoology, Bharathiar University, Coimbatore, 641 046, India
- Orford, S. J. and J. N. Timmis (2000). Expression of a lipid transfer protein gene family during cotton fibre development. *Biochimica et Biophysica Acta*. Jan. 17(2): 275-284. {a} Department of Genetics, University of Adelaide, Adelaide, SA, 5005, Australia
- Parker, C. D., Jr., V. J. Mascarenhas, et al. (2000). Survival rates of tobacco budworm (Lepidoptera: Noctuidae) larvae exposed to transgenic cottons expressing insecticidal protein of *Bacillus thuringiensis* Berliner. *Journal of Entomological Science*. April 35(2): 105-117. {a} Department of Entomology and Plant Pathology, Mississippi State University, Mississippi State, MS, 39762, USA
- Patel, J. C., U. G. Patel, et al. (2000). Performance of intra-hirsutum and Asiatic cotton hybrids based on male sterility system. *Indian Journal of Genetics and Plant Breeding*. Feb. 60(1): 111-115. {a} Main Cotton Research Station, Gujarat Agricultural University, Surat, 395 007, India
- Perez, A. E., P. M. D. Garcia, et al. (2000). Differentiation of cotton-defoliating and nondefoliating pathotypes of *Verticillium dahliae* by RAPD and specific PCR analyses. *European Journal of Plant Pathology*. [print] July 106(6): 507-517. {a} Departamento de Proteccion de Cultivos, IAS, CSIC, 14080, Cordoba, Spain
- Pettigrew, W. T. (2000). Potassium deficiency increases specific leaf weights and leaf glucose levels in field-grown cotton. *Agronomy Journal*. Nov. Dec. 91(6): 962-968. {a} Crop Genetics and Production Res. Unit, USDA-ARS, Stoneville, MS, 38776, USA
- Phogat, D. S., D. P. Singh, et al. (2000). Correlation studies on bollworm incidence, gossypol seed cotton yield and fibre characters in American cotton (*Gossypium hirsutum* L.). *Annals of Biology Hissar*. June 16(1): 89-93. {a} Department of Plant Breeding, CCS Haryana Agricultural University, Hisar, HR, 125 004, India
- Punitha, D. and T. S. Raveendran (2000). Metroglyph analysis for intraspecific coloured linted cotton genotypes (*Gossypium hirsutum* L.). *Crop Research Hisar*. [print] September 20(2): 317-320. {a} Department of Cotton, Tamil Nadu Agricultural University, Coimbatore, TN, India
- Rajasekaran, K., R. L. Hudspeth, et al. (2000). High-frequency stable transformation of cotton (*Gossypium hirsutum* L.) by particle bombardment of embryogenic cell suspension cultures. *Plant Cell Reports*. [print] May 19(6): 539-545. {a} Southern Regional Research Center, USDA, ARS, 1100 Robert E. Lee Blvd., New Orleans, LA, 70124, USA

- Rathinavel, K., C. Dharmalingam, et al. (2000). Effect of seed pelleting on the productivity of rice fallow cotton (*Gossypium hirsutum* L.) cv. MCU 7. *Advances in Plant Sciences*. [print] June 13(1): 213-217. {a} Seed Technology, Regional Station, Central Institute for Cotton Research, Coimbatore, 641003, India
- Ren, G. j., Y. z. Chen, et al. (2000). Studies on flower bud differentiation and changes of endogenous hormones of *Gossypium hirsutum*. *Xibei Zhiwu Xuebao*. [print] Sept 20(5): 847-851. {a} Department of Biology, Shandong University, Ji'nan, 250010, China
- Rhoton, F. E. (2000). Influence of time on soil response to no-till practices. *Soil Science Society of America Journal*. [print] March April 64(2): 700-709. {a} National Sedimentation Laboratory, USDA-ARS, Oxford, MS, 38655, USA
- Rosolem, C. A. and A. Costa (2000). Cotton growth and boron distribution in the plant as affected by a temporary deficiency of boron. *Journal of Plant Nutrition*. [print] June 23(6): 815-825. {a} Department of Crop Science, Sao Paulo State University, 18603-970, Botucatu, SP, Brazil
- Ruan, Y. L., D. J. Llewellyn, et al. (2000). Pathway and control of sucrose import into initiating cotton fibre cells. *Australian Journal of Plant Physiology*. [print] 27(8-9): 795-800. {a} CSIRO Plant Industry, Canberra, ACT, 2601, Australia
- Sacchi, G. A., A. Abruzzese, et al. (2000). Efflux and active re-absorption of glucose in roots of cotton plants grown under saline conditions. *Plant and Soil*. [print] 220(1-2): 1-11. {a} Dipartimento di Fisiologia delle Piante Coltivate e Chimica Agraria, Universita degli Studi di Milano, I-20133, Milano, Italy
- Samaras, V. and C. Kallianou (2000). Effect of sewage sludge application on cotton yield and contamination of soils and plant leaves. *Communications in Soil Science and Plant Analysis*. Feb. 31(3-4): 331-343. {a} Laboratory of Soils and Agricultural Chemistry, Agricultural University of Athens, Iera Odos 75, 11855, Athens, Greece
- Sanamyan, M. F., J. E. Petlyakova, et al. (2000). The development and a cytogenetic study of monosomics of *Gossypium hirsutum* L. *Biologia Plantarum Prague*. [print] 43(2): 193-197. {a} Department of Biology, Tashkent State University, 700095, Tashkent, Uzbekistan
- Schwab, G. J., G. L. Mullins, et al. (2000). Growth and nutrient uptake by cotton roots under field conditions. *Communications in Soil Science and Plant Analysis*. Jan. 31(1-2): 149-164. {a} Department of Crop and Soil Environmental Sciences, Virginia Tech, 424 Smyth Hall (0403), Blacksburg, VA, 24061, USA
- Singh, T. and J. S. Brar (2000). Scheduling of first and last irrigation to cotton (*Gossypium hirsutum* L.). *Annals of Biology Hissar*. June 16(1): 1-2. {a} Regional Research Station, PAU, Bathinda, PU, 151 001, India
- Smith, D. T., R. V. Baker, et al. (2000). Palmer amaranth (*Amaranthus palmeri*) impacts on yield, harvesting, and ginning in dryland cotton (*Gossypium hirsutum*). *Weed Technology*. Jan. March 14(1): 122-126. {a} Department of Soil and Crop Sciences, Texas Agricultural Experiment Station, Texas A and M University, College Station, TX, 77843-2474, USA
- Soomro, A. R. (2000). Assessment of *Gossypium hirsutum* F1 hybrids against commercial varieties NIAB-78 and CRIS-9 in Sakrand conditions. *Pakistan Journal of Scientific and Industrial Research*. [print] June 43(3): 184-187. {a} Central Cotton Research Institute Sakrand, District Nawabshah, Sindhi, Pakistan
- Stapel, J. O., A. M. Cortesero, et al. (2000). Disruptive sublethal effects of insecticides on biological control: Altered foraging ability and life span of a parasitoid after feeding on extrafloral nectar of cotton treated with systemic insecticides. *Biological Control*. March 17(3): 243-249. {a} Laboratoire d'Ecobiologie des Insectes Parasitoides, Universite de Rennes 1, Avenue du General Leclerc, 35042, Rennes Cedex, France
- Stewart, A. M., K. L. Edmisten, et al. (2000). Boll openers in cotton: Effectiveness and environmental influences. *Field Crops Research*. [print] July 67(1): 83-90. {a} Department of Crop Science, North Carolina State University, NCSU, Raleigh, NC, 27695-7620, USA

- Sun, X. f., Q. s. Zheng, et al. (2000). Salinity injury to germination and growth of cotton (*Gossypium hirsutum* L.) at emergence and seedling stages. *Journal of Plant Resources and Environment*. [print] July 9(3): 22-25. {a} Key Laboratory of Crop Growth Regulation, Ministry of Agriculture of China, Nanjing Agricultural University, Nanjing, 210095, China
- Sundararaju, D. and P. C. S. Babu (2000). Improved mass culture technique for the neem mosquito bug, *Helopeltis antonii* Signoret (Heteroptera : Miridae). *Journal of Entomological Research New Delhi*. [print] March 24(1): 73-82. {a} National Research Centre for Cashew, Shanthigodu, Puttur, Dakshina Kannada, KRN, 574 202, India
- Tan, X. P., W. Q. Liang, et al. (2000). Expression pattern of (+)-delta-cadinene synthase genes and biosynthesis of sesquiterpene aldehydes in plants of *Gossypium arboreum* L. *Planta Berlin*. March 210(4): 644-651. {a} National Laboratory of Plant Molecular Genetics, Institute of Plant Physiology of Shanghai Institutes for Biological Science, The Chinese Academy of Sciences, 300 Fenglin Road, Shanghai, 200032, China
- Thakur, A., P. S. Thakur, et al. (2000). Effect of bioregulators, bioextracts and potassium on growth and related attributes in bell pepper (*Capsicum annum*) varieties under water stress. *Indian Journal of Agricultural Sciences*. [print] April 70(4): 255-258. {a} Dr Y S Parmar University of Horticulture and Forestry, Nauni-Solan, Himachal Pradesh, 173 230, USA
- Timmannavar, S. R. and B. C. Patil (2000). Performance of cotton genotypes of different species under rainfed conditions. *Crop Research Hisar*. [print] September 20(2): 312-316. {a} Department of Crop Physiology, University of Agricultural Sciences, Dharwad, KRN, 5, India
- Triantafilis, J., G. M. Laslett, et al. (2000). Calibrating an electromagnetic induction instrument to measure salinity in soil under irrigated cotton. *Soil Science Society of America Journal*. [print] May June 64(3): 1009-1017. {a} Australian Cotton Cooperative Research Centre, Dep. of Agricultural Chemistry and Soil Science, The University of Sydney, Sydney, NSW, 2006, Australia
- Walker, N. R., T. L. Kirkpatrick, et al. (2000). Influence of *Meloidogyne incognita* and *Thielaviopsis basicola* populations on early-season disease development and cotton growth. *Plant Disease*. April 84(4): 449-453. {a} Department of Entomology and Plant Pathology, Oklahoma State University, Stillwater, OK, 74078, USA
- Webster, E. P., D. R. Shaw, et al. (2000). Influence of cultivation timing on pyrithiobac performance in cotton (*Gossypium hirsutum*). *Weed Technology*. Jan. March 14(1): 116-121. {a} Mississippi State University, Mississippi State, MS, 39762, USA
- Wheeler, M. H. (2000). Cryostorage of cottonseed with liquid nitrogen. *Seed Science and Technology*. [print] 28(2): 357-366. {a} Southern Plains Agricultural Research Center, Cotton Pathology Research Unit, USDA, ARS, 2765 F and B Road, College Station, TX, 77845, USA
- Williams, L., III and N. P. Tugwell (2000). Histological description of tarnished plant bug (Heteroptera: Miridae) feeding on small cotton floral buds. *Journal of Entomological Science*. April 35(2): 187-195. {a} USDA-ARS Southern Insect Management Research Unit, Stoneville, MS, 38776-0346, USA
- Wise, R. R., C. G. F. Sassenrath, et al. (2000). A comparison of leaf anatomy in field-grown *Gossypium hirsutum* and *G. barbadense*. *Annals of Botany London*. [print] October 86(4): 731-738. {a} Department of Biology, University of Wisconsin-Oshkosh, Oshkosh, WI, 54901, USA
- Xie, Y. Q., Z. Zhu, et al. (2000). Isolation of a strong promoter and its functional study. *Acta Botanica Sinica*. [print] Jan 42(1): 50-54. {a} Institute of Genetics, Chinese Academy of Sciences, Beijing, 100101, China
- Yu, X. H., Y. Q. Zhu, et al. (2000). Alterations of root and fiber in transgenic cotton plants with chimeric Ph/P-ipt gene expression. *Acta Botanica Sinica*. [print] Jan 42(1): 59-63. {a} National Laboratory of Plant Molecular Genetics, Shanghai Research Center of Life Sciences, Institute of Plant Physiology of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai, 200032, China

Yu, X. H., Y. Q. Zhu, et al. (2000). Isolation of GAE6-3A 5'-upstream fragment from *Gossypium arboreum* and its expression in tobacco. *Acta Phytophysiological Sinica*. [print] 26(2): 143-147. {a} National Laboratory of Plant Molecular Genetics, Shanghai Institute of Plant Physiology, Chinese Academy of Sciences, Shanghai, 200032, China

Yuan, Y. L., Y. H. Chen, et al. (2000). Effects of the dominant glandless gene Gl2e on agronomic and fibre characters of Upland cotton. *Plant Breeding*. Feb. 119(1): 59-64. {a} Key Laboratory of Crop Germplasm and Breeding, Ministry of Agriculture, and Department of Agronomy, Nanjing Agricultural University, Nanjing, Jiangsu, 210095, China

Zhang, B. H., F. Liu, et al. (2000). Plant regeneration via somatic embryogenesis in cotton. *Plant Cell Tissue and Organ Culture*. [print] 60(2): 89-94. {a} 1632 West 6th Street, APT F, Austin, TX, 78703, USA

Zhao, D. and D. M. Oosterhuis (2000). Dynamics of non-structural carbohydrates in developing leaves, bracts and floral buds of cotton. *Environmental and Experimental Botany*. [print] June 43(3): 185-195. {a} Department of Crop, Soil, and Environmental Sciences, Altheimer Laboratory, University of Arkansas, 276 Altheimer Drive, Fayetteville, AR, 72704, USA

Zhao, D. and D. M. Oosterhuis (2000). Cotton responses to shade at different growth stages: Growth, lint yield and fibre quality. *Experimental Agriculture*. Jan. 36(1): 27-39. {a} Department of Crop, Soil and Environmental Sciences, Altheimer Laboratory, University of Arkansas, 276 Altheimer Drive, Fayetteville, AR, 72704, USA

Abraham, K. J., M. L. Pierce, et al. (1999). The phytoalexins desoxyhemigossypol and hemigossypol are elicited by *Xanthomonas* in *Gossypium* cotyledons. *Phytochemistry Oxford*. Nov. 52(5): 829-836. {a} Department of Biochemistry and Molecular Biology, Oklahoma State University, Stillwater, OK, 74078-3035, USA

Adcock, C. W., A. O. Abaye, et al. (1999). Subsurface liming effects on cotton development and root growth: A greenhouse study. *Communications in Soil Science and Plant Analysis* 30(3-4): 317-327. {a} Department of Crop and Soil Environmental Sciences, Virginia Polytechnic Institute and State University, 330 Smyth Hall, Blacksburg, VA, 24061-0404, USA

Akade, J. H., A. H. Sonone, et al. (1999). Cyto-morphological investigations in some sterile plants of upland cotton (*Gossypium hirsutum* L.). *Journal of Maharashtra Agricultural Universities*. [print] September 24(3): 331-333. {a} Department of Botany, Mahatma Phule Krishi Vidyapeeth, Rahuri, Distt. Ahmednagar, 413 722, India

Akhtar, L. H., S. Ahmad, et al. (1999). Effects of macronutrients and CaCl₂ on callus growth of *Gossypium hirsutum* L. cv. Acala SJ2. *Scientific Khyber*. July 12(2): 5-12. {a} Regional Agricultural Research Institute, Bahawalpur, Pakistan

Akhtar, L. H., J. Gorham, et al. (1999). Heat stress tolerance studies in callus cultures of *Gossypium hirsutum* L. cv. Acala SJ2. *Pakistan Journal of Botany*. [print] December 31(2): 271-281. {a} Regional Agricultural Research Institute, Bahawalpur, Pakistan

Anderson, P. and H. Alborn (1999). Effects on oviposition behaviour and larval development of *Spodoptera littoralis* by herbivore-induced changes in cotton plants. *Entomologia Experimentalis et Applicata*. July 92(1): 45-51. {a} Department of Plant Protection Sciences, Swedish University of Agricultural Sciences, S-230 53, Alnarp, Sweden

Ashraf, M. and S. Ahmad (1999). Exploitation of intra-specific genetic variation for improvement of salt (NaCl) tolerance in upland cotton (*Gossypium hirsutum* L.). *Hereditas Lund*. Dec. 131(3): 253-256. {a} Department of Botany, University of Agriculture, Faisalabad, Pakistan

Ashraf, M. and Z. U. Zafar (1999). Some physiological characteristics in resistant and susceptible cotton cultivars infected with cotton leaf curl virus. *Biologia Plantarum Prague* 42(4): 615-620. {a} 51-C Sheikh Colony, ABC Road, Faisalabad, Pakistan

- Ashraf, M., Z. U. Zafar, et al. (1999). Some morpho-anatomical characteristics of cotton (*Gossypium hirsutum* L.) in relation to resistance to cotton leaf curl virus (CLCuV). *Journal of Applied Botany*. Sept. 73(3-4): 76-82. {a} Institute of Pure and Applied Biology, B.Z. University, Multan, Pakistan
- Askew, S. D. and J. W. Wilcut (1999). Cost and weed management with herbicide programs in glyphosate-resistant cotton (*Gossypium hirsutum*). *Weed Technology*. April June 13(2): 308-313. {a} Crop Science Department, North Carolina State University, Raleigh, NC, 27695-7620, USA
- Azhar, F. M. and S. U. K. Ajmal (1999). Diallel analysis of oil content in seed of *Gossypium hirsutum* L. *Journal of Genetics and Breeding*. March 53(1): 19-23. {a} Department of Plant Breeding and Genetics, University of Agriculture, Faisalabad, Pakistan
- Bauer, P. J. and D. W. Reeves (1999). A comparison of winter cereal species and planting dates as residue cover for cotton grown with conservation tillage. *Crop Science*. Nov. Dec. 39(6): 1824-1830. {a} USDA-ARS Coastal Plains Soil, Water, and Plant Research Center, 2611 West Lucas Street, Florence, SC, 29501, USA
- Bednarz, C. W., N. W. Hopper, et al. (1999). Effects of foliar fertilization of Texas Southern High Plains cotton: Leaf phosphorus, potassium, zinc, iron, manganese, boron, calcium, and yield distribution. *Journal of Plant Nutrition*. June 22(6): 863-875. {a} University of Georgia Coastal Plain Experiment Station, Tifton, GA, 31793, USA
- Bharpoda, T. M., H. P. Patel, et al. (1999). Evaluation of economic threshold levels for *Helicoverpa armigera* on 'H 6' cotton (*Gossypium hirsutum*) in central Gujarat region. *Indian Journal of Agricultural Sciences*. April 69(4): 304-305. {a} Gujarat Agricultural University, Campus Anand, Anand, 388110, India
- Bhat, R. G. and K. V. Subbarao (1999). Host range specificity in *Verticillium dahliae*. *Phytopathology*. Dec. 89(12): 1218-1225. {a} Department of Plant Pathology, University of California, Davis, c/o U.S. Agricultural Research Station, Salinas, CA, USA
- Bi, I. V., J. P. Baudoin, et al. (1999). Development of high-gossypol cotton plants with low-gossypol seeds using trispecies bridge crosses and in vitro culture of seed embryos. *Euphytica* 106(3): 243-251. {a} Faculte Universitaire des Sciences Agronomiques, Unite de Phytotechnie des Regions Intertropicales, 2 Passage des Deportes, B-5030, Gembloux, Belgium
- Bi, I. V., A. Maquet, et al. (1999). Breeding for "low-gossypol seed and high-gossypol plants" in upland cotton. Analysis of tri-species hybrids and backcross progenies using AFLPs and mapped RFLPs. *Theoretical and Applied Genetics*. Nov. 99(7-8): 1233-1244. {a} Faculte Universitaire des Sciences Agronomiques, Unite de Phytotechnie des Regions Intertropicales, 2 Passage des deportes, B-5030, Gembloux, Belgium
- Bianchini, G. M., R. D. Stipanovic, et al. (1999). Induction of delta-cadinene synthase and sesquiterpenoid phytoalexins in cotton by *Verticillium dahliae*. *Journal of Agricultural and Food Chemistry*. Oct. 47(10): 4403-4406. {a} Southern Crops Research Laboratory, Agricultural Research Service, U.S. Department of Agriculture, College Station, TX, 77845, USA
- Bondada, B. R., D. M. Oosterhuis, et al. (1999). Cotton growth and yield as influenced by different timing of late-season foliar nitrogen fertilization. *Nutrient Cycling in Agroecosystems*. May 54(1): 1-8. {a} Department of Plant and Soil Science, University of Tennessee, Knoxville, TN, 37901, USA
- Brubaker, C. L., A. H. D. Brown, et al. (1999). Production of fertile hybrid germplasm with diploid Australian *Gossypium* species for cotton improvement. *Euphytica* 108(3): 199-213. {a} Centre for Plant Biodiversity Research, CSIRO Plant Industry, Canberra, ACT, 2601, Australia
- Burkle, R., Y. R. Mehta, et al. (1999). Reaction of cotton cultivars to *Colletotrichum gossypii* var. *cephalosporioides*. *Summa Phytopathologica*. July Sept 25(3): 278-281. {a} IAPAR, CEP 86.001-970, Londrina, PR, Brazil

Caesar, A. J., G. Campobasso, et al. (1999). Effects of European and U.S. strains of *Fusarium* spp. pathogenic to leafy spurge on North American grasses and cultivated species. *Biological Control*. June 15(2): 130-136. {a} Pest Management and Agricultural Systems Research Unit, Northern Plains Agricultural Research Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Sidney, MT, 59270, USA

Campbell, C. S. and K. J. McInnes (1999). Response of in situ leaf psychrometer to cuticle removal by abrasion. *Agronomy Journal*. Sept. Oct. 91(5): 859-862. {a} Dep. of Soil and Crop Science, Texas A and M University, College Station, TX, USA

Celini, L. and J. Vaillant (1999). Spatio-temporal distribution of winged *Aphis gossypii* (Hemiptera: Aphididae) presences in cotton Malvaceae field. *Canadian Entomologist*. Nov. Dec. 131(6): 813-824. {a} Ministère de la Coopération, 10, rue Alphonse karr, 75019, Paris, France

Chachar, Q. I., D. L. Laidman, et al. (1999). Radiolabelled (Me-3H) thymidine incorporation into DNA in roots of cotton seedling. *Pakistan Journal of Botany*. June 31(1): 79-84. {a} Department of Physiology and Biochemistry, Faculty of Crop Production, Sindh Agriculture University, Tandojam, Pakistan

Chapman, K. D., B. Venables, et al. (1999). N-Acylethanolamines in seeds. Quantification of molecular species and their degradation upon imbibition. *Plant Physiology Rockville*. Aug. 120(4): 1157-1164. {a} Department of Biological Sciences, Division of Biochemistry and Molecular Biology, University of North Texas, Denton, TX, 76203-5220, USA

Chu, C. c., A. C. Cohen, et al. (1999). *Bemisia tabaci* (Hemiptera:Aleyrodidae) biotype B colonisation and leaf morphology relationships in upland cotton cultivars. *Australian Journal of Entomology*. May 38(2): 127-131. {a} Western Cotton Research Laboratory, USDA-ARS, 4135 E. Broadway Rd, Phoenix, AZ, 85040, USA

Cothren, J. T., P. H. Jost, et al. (1999). Cotton desiccation and defoliation by paraquat influenced by time of day. *Crop Science*. May June 39(3): 859-862. {a} Dep. of Soil and Crop Sciences, Texas A and M Univ., College Station, TX, 77843-2474, USA

Crafts, B. S. J. and C. c. Chu (1999). Insect clip cages rapidly alter photosynthetic traits of leaves. *Crop Science*. Nov. Dec. 39(6): 1896-1899. {a} Western Cotton Research Laboratory, USDA-ARS, 4135 E. Broadway Rd., Phoenix, AZ, 85040-8803, USA

Cronn, R. C., R. L. Small, et al. (1999). Duplicated genes evolve independently after polyploid formation in cotton. *Proceedings of the National Academy of Sciences of the United States of America*. Dec. 96(25): 14406-14411. {a} Department of Botany, Iowa State University, Bessey Hall 353, Ames, IA, USA

Culpepper, A. S. and A. C. York (1999). Weed management and net returns with transgenic, herbicide-resistant, and nontransgenic cotton (*Gossypium hirsutum*). *Weed Technology*. April June 13(2): 411-420. {a} Crop Science Department, North Carolina State University, Raleigh, NC, 27695-7620, USA

Dani, R. G. (1999). Genetic improvement of seed oil content, following indirect selection for earliness and fibre yield in cotton (*Gossypium hirsutum* L.). *Advances in Plant Sciences*. Dec. 12(2): 479-492. {a} Division of Crop Improvement, Central Institute for Cotton Research, Shankarnagar P.O., Nagpur, 440010, India

Deguine, J. P., J. Martin, et al. (1999). Extreme polyphagy of *Aphis gossypii* Glover (Hemiptera: Aphididae) during the dry season in northern Cameroon. *Insect Science and its Application*. Jan. March 19(1): 23-36. {a} Unite de Formation et de Recherche d'Ecologie Animale et de Zoologie Agricole, ENSA-INRA Ecole Nationale Supérieure Agronomique, 2 Place Viala, 34060, Montpellier Cedex 1, France

Deshmukh, V. V., V. K. Mohod, et al. (1999). Variability, heritability and genetic advance in upland cotton (*G. hirsutum* L.). *PKV Research Journal*. Jan. 23(1): 21-23.

Elena, K. (1999). Genetic relationships among *Verticillium dahliae* isolates from cotton in Greece based on vegetative compatibility. *European Journal of Plant Pathology*. Sept. 105(6): 609-616. {a} Benaki Phytopathological Institute, 8 S. Delta str., GR 145 61 Kifissia, Athens, Greece

Ellington, J. and M. Southward (1999). Quadrat sample precision and cost with a high-vacuum insect sampling machine in cotton ecosystems. *Environmental Entomology*. Aug. 28(4): 722-728. {a} Entomology, Plant Pathology and Weed Science Department, Agricultural Experiment Station, New Mexico State University, Las Cruces, NM, 88003, USA

Endondo, C. and M. Samatana (1999). Influence of cowpea sowing date on cotton yield in a cotton/cowpea intercropping sequence. *Cahiers Agricultures*. May June 8(3): 215-217. {a} Institut de recherche agricole pour le developpement (IRAD), Maroua, Cameroon

Farooq, S., N. Iqbal, et al. (1999). Isozyme markers in cotton breeding 1. Standardization of different isozyme systems for identification of different cultivars of cotton (*Gossypium hirsutum*). *Pakistan Journal of Botany*. June 31(1): 5-20. {a} Nuclear Institute for Agriculture and Biology (NIAB), Jhang Road, Faisalabad, Pakistan

Farooq, S. and H. Sayyed (1999). Isozyme markers in cotton breeding: III. Variation in the intensity of isozymes of enzyme peroxidase exhibited by different loci of different cotton varieties and there relationship with cotton leaf virus disease. *Pakistan Journal of Botany*. [print] December 31(2): 361-370. {a} Nuclear Institute for Agriculture and Biology (NIAB), Jhang Road, Faisalabad, Pakistan

Farooq, S. and H. Sayyed (1999). Isozyme markers in cotton breeding: II: Inter and intravarietal variation in the activity of isozymes of the enzyme peroxidase as affected by area of cotton cultivation. *Pakistan Journal of Botany*. [print] December 31(2): 347-359. {a} Nuclear Institute for Agriculture and Biology (NIAB), Jhang Road, Faisalabad, Pakistan

Ferreira, A. and L. F. Mesquita (1999). Evaluation of cotton genotypes resistance to *Alabama argillacea* (Huebner, 1818) (Lepidoptera: Noctuidae): II. Antibiosis. *Bragantia* . [print] 58(2): 287-292. {a} Departamento de Fitossanidade, FCAV, UNESP, Rodovia Carlos Tonanni, km 5, 14870-000, Jaboticabal, SP, Brazil

Flint, H. M. and N. J. Parks (1999). Seasonal infestation by pink bollworm, *Pectinophora gossypiella* (Saunders), of transgenic and non-transgenic cultivars of cotton, *Gossypium Hirsutum* L., in central Arizona. *Southwestern Entomologist*. March 24(1): 13-20. {a} ARS, PWA, Western Cotton Research Laboratory, USDA, 4135 E. Broadway Road, Phoenix, AZ, 85040-8803, USA

Francovig, P. C., Y. R. Mehta, et al. (1999). Sources of resistance to *Stemphylium solani* in cotton cultivars. *Summa Phytopathologica*. July Sept 25(3): 217-222. {a} Area de Protecao de Plantas, Instituto Agronomico do Parana, IAPAR, Londrina, PR, Brazil

Galanopoulou, S. S. and D. Roupakias (1999). Performance of cotton F1 hybrids and its relation to the mean yield of advanced bulk generations. *European Journal of Agronomy*. June 11(1): 53-62. {a} Laboratory of Agronomy, School of Agriculture, University of Thessaly, 38334, Volos, Greece

Godoy, A. S. and G. A. Palomo (1999). Genetic analysis of earliness in upland cotton (*Gossypium hirsutum* L.). II. Yield and lint percentage. *Euphytica* 105(2): 161-166. {a} Instituto Nacional de Investigaciones Forestales y Agropecuarias, Torreon, COAH, Mexico

Godoy, A. S. and G. A. Palomo (1999). Genetic analysis of earliness in upland cotton (*Gossypium hirsutum* L.). I. Morphological and phenological variables. *Euphytica* 105(2): 155-160. {a} Instituto Nacional de Investigaciones Forestales y Agropecuarias, Torreon, COAH, Mexico

Grantz, D. A. and D. L. Vaughn (1999). Vertical profiles of boundary layer conductance and wind speed in a cotton canopy measured with heated brass surrogate leaves. *Agricultural and Forest Meteorology*. Nov. 97(3): 187-197. {a} Kearney Agricultural Center, 9240 S. Riverbend Avenue, Parlier, CA, USA

Hallmann, J., K. R. Rodriguez, et al. (1999). Chitin-mediated changes in bacterial communities of the soil, rhizosphere and within roots of cotton in relation to nematode control. *Soil Biology and Biochemistry* 31(4): 551-560. {a} Alabama Agricultural Experiment Station, Department of Plant Pathology, Biological Control Institute, Auburn University, Auburn, AL, 36849-5409, USA

Headrick, D. H., T. S. Bellows, et al. (1999). Development and reproduction of a population of *Eretmocerus eremicus* (Hymenoptera: Aphelinidae) on *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Environmental Entomology*. April 28(2): 300-306. {a} Department of Entomology, University of California, Riverside, CA, 92521, USA

Heagle, A. S., J. E. Miller, et al. (1999). Ozone stress, carbon dioxide enrichment, and nitrogen fertility interactions in cotton. *Crop Science*. May June 39(3): 731-741. {a} Plant Growth and Development Research Unit, USDA-ARS Air Quality, 3908 Inwood Road, Raleigh, NC, 27603, USA

Hebbar, K. P., B. A. Bailey, et al. (1999). An improved granular formulation for a mycoherbicidal strain of *Fusarium oxysporum*. *Weed Science*. July Aug 47(4): 473-478. {a} Biocontrol of Plant Diseases Laboratory, BARC-W, USDA-ARS, Building 011A, Beltsville, MD, 20705, USA

Hill, M. K., K. J. Lyon, et al. (1999). Identification of disease response genes expressed in *Gossypium hirsutum* upon infection with the wilt pathogen *Verticillium dahliae*. *Plant Molecular Biology*. May 40(2): 289-296. {a} Cooperative Research Centre for Sustainable Cotton Production, School of Biological Sciences A12, University of Sydney, Sydney, NSW, 2006, Australia

Hoang, C. V., H. G. Wessler, et al. (1999). Identification and expression of cotton (*Gossypium hirsutum* L.) plastidial carbonic anhydrase. *Plant and Cell Physiology*. Dec. 40(12): 1262-1270. {a} Department of Biological Sciences, University of North Texas, Denton, TX, 76203-5220, USA

Holman, E. M. and D. M. Oosterhuis (1999). Cotton photosynthesis and carbon partitioning in response to floral bud loss due to insect damage. *Crop Science*. Sept. Oct. 39(5): 1347-1351. {a} Northeast Res. Stn., Louisiana State Univ. Agric. Center, Saint Joseph, LA, 71366, USA

Howard, D. D., M. E. Essington, et al. (1999). Vertical phosphorus and potassium stratification in no-till cotton soils. *Agronomy Journal*. March April 91(2): 266-269. {a} Plant and Soil Sci. Dep., Univ. of Tenn., Knoxville, TN, 37901-1071, USA

Hron, R. J., Sr., H. L. Kim, et al. (1999). Determination of (+)-, (-)-, and total gossypol in cottonseed by high-performance liquid chromatography. *Journal of the American Oil Chemists' Society*. Nov. 76(11): 1351-1355. {a} Southern Regional Research Center, New Orleans, LA, USA

Hsu, C. Y., R. G. Creech, et al. (1999). Analysis of promoter activity of cotton lipid transfer protein gene LTP6 in transgenic tobacco plants. *Plant Science Shannon*. May 143(1): 63-70. {a} Department of Biochemistry and Molecular Biology, Mississippi State University, Mississippi State, MS, 39762, USA

Hulugalle, N. R., P. C. Entwistle, et al. (1999). Sowing wheat or field pea as rotation crops after irrigated cotton in a grey Vertosol. *Australian Journal of Soil Research* 37(5): 867-889. {a} Co-operative Research Centre for Sustainable Cotton Production, Narrabri, NSW, 2390, Australia

Hunt, P. G., T. A. Matheny, et al. (1999). Sample size influences estimation of cotton dry matter, nitrogen accumulation, and lint yield. *Communications in Soil Science and Plant Analysis*. Nov. 30(19-20): 2801-2810. {a} Coastal Plains Soil, Water, and Plant Research Center, USDA-ARS, 2611 W. Lucas St., Florence, SC, 29501-1242, USA

Isaacs, R., M. Cahill, et al. (1999). Host plant evaluation behaviour of *Bemisia tabaci* and its modification by external or internal uptake of imidacloprid. *Physiological Entomology*. June 24(2): 101-108. {a} Department of Entomology, 203 Pesticide Research Center, Michigan State University, East Lansing, MI, 48824, USA

Ji, Y., D. M. De, et al. (1999). New ribosomal RNA gene locations in *Gossypium hirsutum* mapped by meiotic FISH. *Chromosoma Berlin* 108(3): 200-207. {a} Department of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843, USA

Ji, Y., W. A. Raska, et al. (1999). Identification and distinction among segmental duplication-deficiencies by fluorescence in situ hybridization (FISH)-adorned multivalent analysis. *Genome*. Aug. 42(4): 763-771. {a} Department of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843, USA

Johnson, A. W., N. A. Minton, et al. (1999). Bahiagrass, corn, cotton rotations, and pesticides for managing nematodes, diseases, and insects on peanut. *Journal of Nematology*. June 31(2): 191-200. {a} USDA ARS, Coastal Plain Station, Tifton, GA, 31793, USA

Johnson, A. W., N. A. Minton, et al. (1999). Managing nematodes, fungal diseases, and thrips on peanut with pesticides and crop rotations of bahiagrass, corn, and cotton. *Peanut Science*. Jan. June 29(1): 32-39. {a} Coastal Plain Exp. Stn., USDA-ARS, Tifton, GA, 31793-0748, USA

Jonsson, M. and P. Anderson (1999). Electrophysiological response to herbivore-induced host plant volatiles in the moth *Spodoptera littoralis*. *Physiological Entomology*. Dec. 24(4): 377-385. {a} Department of Plant Protection Sciences, Swedish University of Agricultural Sciences, S-230 53, Alnarp, Sweden

Kairon, M. S., M. R. K. Rao, et al. (1999). Cotton (*Gossypium* species) research: A pursuit towards self-sufficiency and export. *Indian Journal of Agricultural Sciences*. Aug. 68(8 spec. issue): 460-467. {a} Central Institute for Cotton Research, Nagpur, MAH, 440 010, India

Kakani, A., S. Saha, et al. (1999). Genetic mechanism and chromosomal location of pollen-specific gene(s) in *Gossypium*. *Crop Science*. May June 39(3): 668-673. {a} Crop Sci. Res. Lab., USDA-ARS, Mississippi State, MS, 39762, USA

Karban, R. and J. S. Thaler (1999). Plant phase change and resistance to herbivory. *Ecology* Washington D C 80(2): 510-517. {a} Department of Entomology, University of California, Davis, CA, 95616, USA

Karimov, K. K. and S. V. Dontsova (1999). Soluble proteins in cotton seeds as related to their viability. *Fiziologiya Rastanii Moscow*. May June 46(3): 484-491.

Kasperbauer, M. J. (1999). Cotton seedling root growth responses to light reflected to the shoots from straw-covered versus bare soil. *Crop Science*. Jan. Feb. 39(1): 164-167. {a} USDA-ARS, Coastal Plains Soil Water Plant Res. Cent., 2611 W. Lucas St., Florence, SC 29501-1242, USA

Kersting, U., S. Satar, et al (1999). Effect of temperature on development rate and fecundity of apterous *Aphis gossypii* Glover (Hom., Aphididae) reared on *Gossypium hirsutum* L. *Journal of Applied Entomology* 123(1): 23-27. {a} Department of Plant Protection, University of Cukurova, Adana, Turkey

Khan, M. u. D., M. Z. Iqbal, et al. (1999). Phenotypic stability in *Gossypium hirsutum* L. for yield, ginning out turn and staple length in Multan Cotton Zone. *Pakistan Journal of Scientific and Industrial Research*. [print] Dec 42(6): 355-359. {a} Cotton Research Station, Multan, Pakistan

Kohel, R. J. and A. A. Bell (1999). Genetic analysis of two terpenoid variants in cotton (*Gossypium hirsutum* L.). *Journal of Heredity* 90(1): 249-251. {a} USDA-ARS, Crop Germplasm Res. Unit, 2765 F and B Rd., College Station, TX 77845, USA

Kumar, P. R. and T. S. Raveendran (1999). Genetic analysis of yield and quality traits in upland cotton (*Gossypium hirsutum*) through generation mean analysis. *Indian Journal of Agricultural Sciences*. July 69(7): 538-540. {a} Centre for Plant Breeding and Genetics, Tamil Nadu Agricultural University, Coimbatore, 641 003, India

Kumar, V., M. D. Gohil, et al. (1999). Methanol enhanced productivity in cotton. *Indian Journal of Plant Physiology*. April June 4(2): 105-107. {a} Main Cotton Research Station, Gujarat Agricultural University, Surat, 395 007, India

Kumaresan, D., P. Senthilkumar, et al. (1999). Combining ability studies for quantitative traits in cotton (*Gossypium hirsutum* L.). *Crop Research Hisar*. Nov. 18(3): 430-432. {a} Department of Plant Breeding and Genetics, Agricultural College and Research Institute, Madurai, Tamil Nadu, 625 104, India

- Kuznetsov, V. V., V. Y. Rakitin, et al. (1999). Effects of preliminary heat-shock treatment on accumulation of osmolytes and drought resistance in cotton plants during water deficiency. *Physiologia Plantarum*. Dec. 107(4): 399-406. {a} Laboratory of Molecular Mechanisms of Adaptation, Institute of Plant Physiology, Russian Academy of Sciences, 35 Botanicheskaya St., Moscow, 127276, Russia
- Lara, F. M., A. Ferreira, et al. (1999). Resistance of cotton genotypes to *Alabama argillacea* (Huebner) (Lepidoptera: Noctuidae): I - Nonpreference. *Anais da Sociedade Entomologica do Brasil*. Dec. 28(4): 739-744. {a} Depto de Fitossanidade, FCAV/UNESP, Rod. Carlos Tonanni, Km 5, 14870-000, Jaboticabal, SP, Brazil
- Larney, F. J., J. F. Leys, et al. (1999). Dust and endosulfan deposition in a cotton-growing area of Northern New South Wales, Australia. *Journal of Environmental Quality* 28(2): 692-701. {a} Research Centre, Agriculture and Agri-Food Canada, Lethbridge, AB, T1J 4B1, Canada
- Law, R. D. and B. S. J. Crafts (1999). Inhibition and acclimation of photosynthesis to heat stress is closely correlated with activation of ribulose-1,5-bisphosphate carboxylase/oxygenase. *Plant Physiology* Rockville. May 120(1): 173-181. {a} United States Department of Agriculture-Agricultural Research Service, Western Cotton Research Laboratory, 4135 East Broadway Road, Phoenix, AZ, 85040-8803, USA
- Leidi, E. O., M. Lopez, et al. (1999). Variation in carbon isotope discrimination and other traits related to drought tolerance in upland cotton cultivars under dryland conditions. *Field Crops Research* 61(2): 109-123. {a} Dpto. de Biologia Vegetal, IRNAS-CSIC, Avda Reina Mercedes 10, 41012, Seville, Spain
- Lin, T. B., A. Schwartz, et al. (1999). Non-stomatal factors limit cotton photosynthesis under silverleaf whitefly stress. *Physiologia Plantarum*. Nov. 107(3): 303-311. {a} Faculty of Agricultural, Food and Environmental Quality Sciences, The Hebrew University of Jerusalem, Rehovot, 76100, Israel
- Lin, T. B., A. Schwartz, et al. (1999). Photosynthesis and productivity of cotton under silverleaf whitefly stress. *Crop Science*. Jan. Feb. 39(1): 174-184. {a} Dep. Field Crops Vegetables Genetics, Fac. Agric. Food Environ. Quality Sci., Hebrew Univ. Jerusalem, P.O. Box 12, Rehovot 76100, Israel
- Liu, Q., S. P. Singh, et al. (1999). Molecular cloning and expression of a cDNA encoding a microsomal omega-6 fatty acid desaturase from cotton (*Gossypium hirsutum*). *Australian Journal of Plant Physiology* 26(2): 101-106. {a} Division of Plant Industry, Commonwealth Scientific and Industrial Research Organisation, Dickson, ACT, 2601, Australia
- Loguercio, L. L., H. C. Scott, et al. (1999). Hmg-coA reductase gene family in cotton (*Gossypium hirsutum* L.): Unique structural features and differential expression of hmg2 potentially associated with synthesis of specific isoprenoids in developing embryos. *Plant and Cell Physiology*. July 40(7): 750-761. {a} EMBRAPA/CNPMS (Brazilian Agriculture Research Corporation/National Maize and Sorghum Research Center), Rod. MG 424, Km 65, Sete Lagoas, MG, 35701-970, Brazil
- Loguercio, L. L., J. Q. Zhang, et al. (1999). Differential regulation of six novel MYB-domain genes defines two distinct expression patterns in allotetraploid cotton (*Gossypium hirsutum* L.). *Molecular and General Genetics*. June 261(4-5): 660-671. {a} Department of Agronomy and Range Science, University of California, One Shields Avenue, Davis, CA, 95616-8515, USA
- Longer, D. E. and D. M. Oosterhuis (1999). Cotton regrowth and recovery from early season leaf loss. *Environmental and Experimental Botany* 41(1): 67-73. {a} Department of Agronomy, University of Arkansas, Fayetteville, AR, 72701, USA
- Luo, J. and J. H. Gould (1999). In vitro shoot-tip grafting improves recovery of cotton plants from culture. *Plant Cell Tissue and Organ Culture* 57(3): 211-213. {a} Forest Science Department, Texas Agricultural Experiment Station, Texas A and M University, College Station, TX, 77843-2135, USA
- Maliwal, G. L., H. C. Parmar, et al. (1999). Production potential and economics of different cotton (*Gossypium hirsutum*) based cropping systems. *Indian Journal of Agronomy*. March 44(1): 200-202. {a} Department of Agricultural Chemistry and Soil Science, B.A. College of Agriculture, Anand, GUJ, India

- Mane, V. S., S. S. Mehetre, et al. (1999). Response of cotton to plant geometry and fertilizer levels. *Journal of Maharashtra Agricultural Universities*. [print] May 24(2): 174-176. {a} Mahatma Phule Krishi Vidyapeeth, Rahuri, 413 722, India
- Marcone, C., F. Hergenahhn, et al. (1999). Dodder transmission of pear decline, European stone fruit yellows, rubus stunt, picris echioides yellows and cotton phyllody phytoplasmas to periwinkle. *Journal of Phytopathology Berlin*. March 147(3): 187-192. {a} Biologische Bundesanstalt fuer Land- und Forstwirtschaft, Institut fuer Pflanzenschutz im Obstbau, D-69221, Dossenheim, Germany
- Masangkay, R. F., M. O. Mabbayad, et al. (1999). Host range of *Alternaria alternata* f.sp. *sphenocleae* causing leaf blight of *Sphenoclea zeylanica*. *Canadian Journal of Botany*. Jan. 77(1): 103-112. {a} Agronomy, Plant Physiology, and Agroecology Division, International Rice Research Institute, 1271, Makati City, Philippines
- Matocha, C. J. and L. R. Hossner (1999). Mobility of the herbicide pyriithiobac through intact soil columns. *Journal of Agricultural and Food Chemistry*. April 47(4): 1755-1759. {a} Department of Plant and Soil Sciences, University of Delaware, Newark, DE, 19717, USA
- McBride, R. G., R. L. Mikkelsen, et al. (1999). A comparison of three methods for determining root-knot nematode infection of cotton roots. *Nematropica* . [print] Dec 29(2): 147-151. {a} Department of Soil Science, N.C. State University, Raleigh, NC, 27695, USA
- McConnell, J. S., E. D. Vories, et al. (1999). Effect of irrigation termination on the yield, earliness, and fiber qualities of cotton. *Journal of Production Agriculture*. April June 12(2): 263-268. {a} Univ. of Arkansas Southeast Res. and Ext. Cent., Monticello, AR, 71656, USA
- Meena, R. A., M. N. Mishra, et al. (1999). Effect of storage containers on seed longevity in cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences*. Oct. 69(10): 740-742. {a} Regional Station Central Institute for Cotton Research, Sirsa, HR, 125 055, India
- Meena, R. A., K. Rathinavel, et al. (1999). Storage potential of tetraploid and diplid cottons under ambient conditions. *Seed Research New Delhi*. [print] June 27(1): 125-127. {a} Regional Station, Central Institute for Cotton Research, Sirsa, 125 055, India
- Mellon, J. E. and P. J. Cotty (1999). Raffinose content may influence cottonseed susceptibility to aflatoxin contamination. *Journal of the American Oil Chemists' Society*. July 76(7): 883-886. {a} ARS, Southern Regional Research Center, USDA, New Orleans, LA, 70179, USA
- Mensah, R. K. (1999). Habitat diversity: Implications for the conservation and use of predatory insects of *Helicoverpa* spp. in cotton systems in Australia. *International Journal of Pest Management*. April June 45(2): 91-100. {a} NSW Agriculture, Australian Cotton Research Institute, PMB Myall Vale Narrabri, NSW, 2390, Australia
- Moire, L., A. Schmutz, et al. (1999). Glycerol is a suberin monomer. New experimental evidence for an old hypothesis. *Plant Physiology Rockville* 119(3): 1137-1146. {a} Institut fur Botanische Biologie, Universitat Freiburg, A. Gockelstrasse 3, CH-1700, Freiburg, Switzerland
- Monks, C. D., M. G. Patterson, et al. (1999). The effects of row width and leaf shape on cotton growth and yield. *Alabama Agricultural Experiment Station Auburn University Bulletin*. [print] October 638: 1-7. {a} Department of Agronomy and Soils, Auburn University, Auburn, AL, USA
- Monks, C. D., M. G. Patterson, et al. (1999). Effect of pyriithiobac, MSMA, and DSMA on cotton (*Gossypium hirsutum* L.) growth and weed control. *Weed Technology*. Jan. March 13(1): 6-11. {a} Department of Agronomy and Soils and Alabama Agricultural Experiment Station, Auburn University, Auburn, AL, 36849, USA
- Morrison, J. E., Jr., D. F. Williams, et al. (1999). Effect of crop seed water content on the rate of seed damage by red imported fire ants (Hymenoptera: Formicidae). *Journal of Economic Entomology* 92(1): 215-

219. {a} Nat. Resource Conserv. Res. Unit, USDA-ARS-GSWRL, 808 E. Blackland Rd., Temple, TX 76502, USA
- Mullins, G. L., G. J. Schwab, et al. (1999). Cotton response to surface applications of potassium fertilizer: A 10-year summary. *Journal of Production Agriculture*. July Sept 12(3): 434-440. {a} Virginia Tech, 424 Smyth Hall (0403), Blacksburg, VA, 24061, USA
- Murray, F., D. Llewellyn, et al. (1999). Expression of the *Talaromyces flavus* glucose oxidase gene in cotton and tobacco reduces fungal infection, but is also phytotoxic. *Molecular Breeding* 5(3): 219-232. {a} CSIRO Plant Industry, Canberra City, ACT, 2601, Australia
- Nadjimov, U. K., I. M. Scott, et al. (1999). Conditioning of fasciation by gibberellin and genotype in cotton (*Gossypium hirsutum* L.). *Journal of Plant Growth Regulation*. Winter 18(1): 45-48. {a} Institute of Biological Sciences, University of Wales, Aberystwyth, SY23 3DA, UK
- Nadler, A., A. Gamliel, et al. (1999). Practical aspects of salinity effect on TDR-measured water content: A field study. *Soil Science Society of America Journal*. Sept. Oct. 63(5): 1070-1076. {a} Dep of Physical Chemistry of the Soil, Volcani Center, Soil and Water Inst., ARO, Bet Dagan, 50-250, Israel
- Nehl, D. B., P. A. McGee, et al. (1999). Patterns of arbuscular mycorrhiza down the profile of a heavy textured soil do not reflect associated colonization potential. *New Phytologist*. June 142(3): 495-503. {a} Cooperative Research Centre for Sustainable Cotton Production, NSW Agriculture, Myall Vale Mail Run, Narrabri, NSW, 2390, Australia
- Ogallo, J. L., P. B. Goodell, et al. (1999). Management of root-knot nematodes with resistant cotton cv. NemX. *Crop Science* 39(2): 418-421. {a} Dep. of Nematology, Univ. of California, Riverside, CA, 92521, USA
- Opondo, R. M., G. A. Ombakho, et al. (1999). Registration of P799, R64, and S212 cotton germplasm. *Crop Science*. Sept. Oct. 39(5). {a} Natl. Fibre Res. Ctr., Kibos, Kenya Agric. Res. Inst., Kisumu, Kenya
- Opondo, R. M., G. A. Ombakho, et al. (1999). Registration of E790, F962, and H314 cotton germplasm. *Crop Science*. Sept. Oct. 39(5): 1538-1539. {a} Natl. Fibre Res. Ctr., Kibos, Kenya Agric. Res. Inst., Kisumu, Kenya
- Opondo, R. M., G. A. Ombakho, et al. (1999). Registration of A540, B803, and D883 cotton germplasm. *Crop Science*. Sept. Oct. 39(5). {a} Natl. Fibre Res. Ctr., Kibos, Kenya Agric. Res. Inst., Kisumu, Kenya
- Orford, S. J., T. J. Carney, et al. (1999). Characterisation of a cotton gene expressed late in fibre cell elongation. *Theoretical and Applied Genetics*. April 98(5): 757-764. {a} Department of Genetics, University of Adelaide, Adelaide, 5005, Australia
- Ouzounidou, G. and H. A. Constantinidou (1999). Changes in growth and physiology of tobacco and cotton under Ag exposure and recovery: Are they of direct or indirect nature? *Archives of Environmental Contamination and Toxicology*. Nov. 37(4): 480-487. {a} Department Botany, Aristotle University of Thessaloniki, 540 06, Thessaloniki, Greece
- Pace, P. F., H. T. Cralle, et al. (1999). Photosynthate and dry matter partitioning in short- and long-season cotton cultivars. *Crop Science*. July Aug 39(4): 1065-1069. {a} Dep. of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843-2474, USA
- Parker, C. D., Jr. and R. G. Luttrell (1999). Interplant movement of *Heliothis virescens* (Lepidoptera: Noctuidae) larvae in pure and mixed plantings of cotton with and without expression of the Cry1Ac delta-endotoxin protein of *Bacillus thuringiensis* Berliner. *Journal of Economic Entomology*. Aug. 92(4): 837-845. {a} Department of Entomology and Plant Pathology, Mississippi Agricultural and Forestry Experiment Station, Mississippi State University, Mississippi State, MS, 39762, USA

- Patel, U. G., J. C. Patel, et al. (1999). Phenotypic stability in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences*. Feb. 69(2): 116-117. {a} Main Cotton Research Station, Gujarat Agricultural University, Surat, 395 007, India
- Pathan, A. R. K., S. R. Bhunia, et al. (1999). Irrigation requirement of American cotton (*Gossypium hirsutum*) under shallow water table condition. *Indian Journal of Agricultural Sciences*. Oct. 69(10): 738-739. {a} Agricultural Research Station, Rajasthan Agricultural University, Sriganganagar, RAJ, 335 001, India
- Patil, S. and T. S. Raveendran (1999). Genetic variability in F2 populations in cotton (*Gossypium hirsutum*). *PKV Research Journal*. Jan. 23(1): 9-13. {a} College of Agriculture, Nagpur, India
- Patil, S. A., P. M. Salimath, et al. (1999). Nature of gene action for leaf area in cotton. *Crop Research Hisar*. Sept. 18(2): 230-233. {a} Department of Genetics and Plant Breeding, University of Agricultural Sciences, Dharwad, 580 005, India
- Patricio, F. R. A., G. R. A. Klein, et al. (1999). Cottonseed treatment with fungicides. *Summa Phytopathologica*. July Sept 25(3): 250-256. {a} Instituto Biologico, CEP 13.001.970, Campinas, SP, Brazil
- Paulsgrove, M. D. and J. W. Wilcut (1999). Weed management in bromoxynil-resistant *Gossypium hirsutum*. *Weed Science*. Sept. Oct. 47(5): 596-601. {a} Department of Crop Science, North Carolina State University, Raleigh, NC, USA
- Pavasia, M. J., P. T. Shukla, et al. (1999). Combining ability analysis over environments for fibre characters in upland cotton. *Indian Journal of Genetics and Plant Breeding*. Feb. 59(1): 77-81. {a} Main Cotton Research Station, Gujarat Agricultural University, Surat, 395 007, India
- Payne, T., J. Clement, et al. (1999). Heterologous myb genes distinct from GL1 enhance trichome production when overexpressed in *Nicotiana tabacum*. *Development Cambridge* 126(4): 671-682. {a} Department of Botany, Institute for Cellular and Molecular Biology, University of Texas at Austin, Austin, TX, 78713, USA
- Pillai, U. P. and D. McGarry (1999). Structure repair of a compacted Vertisol with wet-dry cycles and crops. *Soil Science Society of America Journal* 63(1): 201-210. {a} Dep. of Agriculture, The Univ. of Queensland, St. Lucia, Queensland, 4072, Australia
- Pirtle, R. M., D. W. Yoder, et al. (1999). Characterization of a palmitoyl-acyl carrier protein thioesterase (FatB1) in cotton. *Plant and Cell Physiology* 40(2): 155-163. {a} Department of Biological Sciences, University of North Texas, Denton, TX, 76203-5220, USA
- Pizzinatto, M. A., L. F. Razera, et al. (1999). Cotton seed quality of regional trial of Sao Paulo state varieties. *Summa Phytopathologica*. April June 25(2): 139-144. {a} Instituto Agronomico de Campinas, CEP 13.001-970, Campinas, SP, Brazil
- Plant, R. E. and M. Keeley (1999). Relationships among final plant map indices in Acala cotton. *Journal of Production Agriculture*. Jan. March 12(1): 61-68. {a} Dep. of Agronomy and Range Sci., Univ. of California, Davis, CA, 95616-8515, USA
- Potikha, T. S., C. C. Collins, et al. (1999). The involvement of hydrogen peroxide in the differentiation of secondary walls in cotton fibers. *Plant Physiology Rockville* 119(3): 849-858. {a} Department of Plant Sciences, Hebrew University of Jerusalem, Givat-Ram, Jerusalem, 91904, Israel
- Prasifka, J. R., P. C. Krauter, et al. (1999). Predator conservation in cotton: Using grain sorghum as a source for insect predators. *Biological Control*. Oct. 16(2): 223-229. {a} Biological Control Laboratory, Department of Entomology, Texas A and M University, College Station, TX, 77843-2475, USA

Punitha, D. and T. S. Raveendran (1999). Heterosis and combining ability studies for quantitative characters in coloured linted cotton genotypes (*Gossypium hirsutum* X *Gossypium barbadense*). *Crop Research Hisar*. Nov. 18(3): 423-429. {a} Senior Research Fellow, Hybrid Rice Evaluation Centre, Gudalur, 643 212, India

Punitha, D. and T. S. Raveendran (1999). Heterosis and combining ability studies in intraspecific coloured linted upland cotton (*Gossypium hirsutum* L.). *Crop Research Hisar*. March 17(2): 245-249. {a} Hybrid Rice Evaluation Centre, TNAU, Gudalur, 643-212, India

Punitha, D., T. S. Raveendran, et al. (1999). Heterosis and combining ability studies for quantitative characters in coloured linted cotton genotypes (*Gossypium hirsutum* X *Gossypium barbadense*). *PKV Research Journal*. Jan. 23(1): 17-20. {a} Hybrid Rice Evaluation centre, Gudalur, India

Punitha, D., T. S. Raveendran, et al. (1999). Combining ability studies for yield and quality traits in interspecific coloured linted cotton (*Gossypium hirsutum* X *Gossypium barbadense*). *PKV Research Journal*. Jan. 23(1): 14-16. {a} Hybrid Rice Evaluation Centre, Gudalur, India

Rabadia, V. S., V. S. Thaker, et al. (1999). Relationship between water content and growth of seed and fibre of three cotton genotypes. *Journal of Agronomy and Crop Science*. Nov. 183(4): 255-261. {a} Department of Biosciences, Saurashtra University, Rajkot, GUJ, 360 005, India

Rajasegar, G., R. S. R. Sree, et al. (1999). Isolation and culture of protoplasts from callus and cell suspension cultures of cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research*. Sept. 33(3): 147-152. {a} Department of Botany, National University of Singapore, Singapore, 0511, Singapore

Rathinavel, K. and C. Dharmalingam (1999). Effect of seed pelleting on elite seedling production in cotton cv. MCU 7 (*Gossypium hirsutum* L.). *Crop Research Hisar*. July 18(1): 137-141. {a} Department of Seed Technology, Tamil Nadu Agricultural University, Coimbatore, 641 003, India

Rathinavel, K., C. Dharmalingam, et al. (1999). Effect of seed pelleting on the productivity of rice fallow cotton (*Gossypium hirsutum* L.) cv. MCU 7. *PKV Research Journal*. Jan. 23(1): 5-9. {a} Seed Technology, Central Institute for Cotton Research, Regional Station, Coimbatore, India

Reddy, K. R., G. H. Davidonis, et al. (1999). Temperature regime and carbon dioxide enrichment alter cotton boll development and fiber properties. *Agronomy Journal*. Sept. Oct. 91(5): 851-858. {a} Dep. of Plant and Soil Sciences, Mississippi State Univ., Mississippi State, MS, USA

Robinson, A. F., C. G. Cook, et al. (1999). Resistance to *Rotylenchulus reniformis* and *Meloidogyne incognita* race 3 in the major cotton cultivars planted since 1950. *Crop Science*. May June 39(3): 850-858. {a} Southern Crops Research Laboratory, USDA-ARS, 2765 F and B Rd., College Station, TX, 77845, USA

Rosolem, C. A., J. P. T. Witacker, et al. (1999). The significance of root growth on cotton nutrition in an acidic low-P soil. *Plant and Soil* 212(2): 185-190. {a} Department of Crop Science, Sao Paulo State University, 18603-970, Botucatu, Sao Paulo, Brazil

Rowland, M. W., D. S. Murray, et al. (1999). Full-season Palmer amaranth (*Amaranthus palmeri*) interference with cotton (*Gossypium hirsutum*). *Weed Science*. May June 47(3): 305-309. {a} Department of Plant and Soil Sciences, Oklahoma State University, Stillwater, OK, 74078, USA

Saini, R. K. and R. Singh (1999). Host plant preference for oviposition by the spiny bollworm *Earias insulana* Boisid. (Lep., Noctuidae). *Journal of Applied Entomology*. May 123(4): 241-245. {a} Department of Entomology, CCS Haryana Agricultural University, Hisar, 125 004, India

Saminathan, V. R., R. K. M. Baskaran, et al. (1999). Biology and predatory potential of green lacewing (*Chrysoperla carnea*) (Neuroptera: Chrysopidae) on different insect hosts. *Indian Journal of Agricultural Sciences*. July 69(7): 502-505. {a} Agricultural College and Research Institute, Tamil Nadu Agricultural University, Madurai, Tamil Nadu, 625 104, India

- Sheshagiri, R. and B. M. Khadi (1999). Somatic instability for chlorophyll pigmentation in cotton (*Gossypium* spp.). *Current Science Bangalore*. Aug. 77(3): 443-446. {a} Agricultural Research Station, Dharwad Farm, Dharwad, 580 007, India
- Shin, H. and R. M. Brown, Jr. (1999). GTPase activity and biochemical characterization of a recombinant cotton fiber annexin. *Plant Physiology Rockville* 119(3): 925-934. {a} Department of Botany, University of Texas, Austin, TX, 78713-7640, USA
- Shrivastava, M. K., K. C. Jain, et al. (1999). Studies on effect of phosphorus application in cotton (*G. hirsutum* L.). *Advances in Plant Sciences*. June 12(1): 279-281. {a} Agril. Research Station, JNKVV, Badnawar (Dhar), MP, 454 660, India
- Singh, T. and Z. S. Brar (1999). Effect of growth-regulators and defoliant on yield and maturity of upland cotton (*Gossypium hirsutum*) under irrigated conditions. *Indian Journal of Agronomy*. March 44(1): 179-184. {a} Department of Agronomy, PAU, Ludhiana, 141 004, India
- Small, R. L., J. A. Ryburn, et al. (1999). Low levels of nucleotide diversity at homoeologous *Adh* loci in allotetraploid cotton (*Gossypium* L.). *Molecular Biology and Evolution*. April 16(4): 491-501. {a} Department of Botany, Iowa State University, Ames, IA, 50011, USA
- Solaiappan, U., N. Subbaraman, et al. (1999). Cotton (*Gossypium hirsutum*)-based intercropping system under different sowing environments. *Indian Journal of Agronomy*. March 44(1): 59-63. {a} Agricultural Research Station, Tamil Nadu Agricultural University, Kovilpatti, 628 501, India
- Starr, J. L. and C. W. Smith (1999). Registration of three root-knot nematode resistant germplasm lines of upland cotton. *Crop Science*. Jan. Feb. 39(1): 303. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843, USA
- Swaminathan, V. R., R. K. Muralibaskaran, et al. (1999). Influence of intercropping on the conservation of *Chrysoperla carnea* (Stephens) in cotton. *Journal of Biological Control*. [print] 13(1-2): 111-114. {a} Department of Agricultural Entomology, Agricultural College and Research Institute (TNAU), Madurai, Tamil Nadu, 625 104, India
- Tengecho, B. and B. M. Khaemba (1999). Effects of diet change on the reproductive parameters of the cotton stainers *Dysdercus cardinalis* and *D. fasciatus* (Hemiptera: Pyrrhocoridae). *Insect Science and its Application*. July Sept 18(3): 211-215. {a} Zoology Department, Kenyatta University, Nairobi, Kenya
- Teuber, S. S., K. C. Jarvis, et al. (1999). Identification and cloning of a complementary DNA encoding a vicilin-like proprotein, *Jug r 2*, from English walnut kernel (*Juglans regia*), a major food allergen. *Journal of Allergy and Clinical Immunology*. Dec. 104(6): 1311-1320. {a} Division of Rheumatology, Allergy and Clinical Immunology, School of Medicine, University of California, Davis, One Shields Ave, TB 192, Davis, CA, 95616, USA
- Thaker, V. S., V. S. Rabadia, et al. (1999). Physiological and biochemical changes associated with cotton fibre development. VII. Carbohydrate metabolism. *Acta Physiologiae Plantarum* 21(1): 57-61. {a} Department of Biosciences, Saurashtra University, Rajkot, 360005, India
- Torbert, H. A., S. A. Prior, et al. (1999). Land management effects on nitrogen and carbon cycling in an Ultisol. *Communications in Soil Science and Plant Analysis*. May 30(9-10): 1345-1359. {a} Soil and Water Research Laboratory, USDA-ARS Grassland, 808 East Blackland Road, Temple, TX, 76502, USA
- Trindade, M. L. B. and L. G. Chiavegato (1999). Effects of decomposition products of deltamethrin on the populational development of *Tetranychus urticae* Koch (Acari: Tetranychidae). *Anais da Sociedade Entomologica do Brasil*. Sept. 28(3): 511-517. {a} Curso de Pos-graduacao em Agronomia, Area de Concentracao Agricultura-Doutoramento, Departamento de Agricultura e Melhoramento Vegetal, Faculdade de Ciencias Agronomicas, UNESP, Bolsista da FAPESP, Botucatu, SP, Brazil
- Triplett, B. A. and D. S. Johnson (1999). Adding gelling agents to cotton ovule culture media leads to subtle changes in fiber development. *In Vitro Cellular and Developmental Biology Plant*. May June 35(3): 265-

270. {a} Cotton Fiber Bioscience Research, USDA-ARS, Southern Regional Research Center, 1100 Robert E. Lee Boulevard, New Orleans, LA, 70124, USA

Van, E. G. A., D. T. Bowman, et al. (1999). Genetic similarity indices for ancestral cotton cultivars and their impact on genetic diversity estimates of modern cultivars. *Crop Science* 39(2): 323-328. {a} Dep. of Crop Science, North Carolina State University, Raleigh, NC, 27695-8604, USA

Van, t. H. J. (1999). Increased nuclear DNA content in developing cotton fiber cells. *American Journal of Botany*. June 86(6): 776-779. {a} Biology Department, Brookhaven National Laboratory, Upton, NY, 11973, USA

Varco, J. J., S. R. Spurlock, et al. (1999). Profitability and nitrogen rate optimization associated with winter cover management in no-tillage cotton. *Journal of Production Agriculture*. Jan. March 12(1): 91-95.

Venugopalan, M. V. and R. Pundarikakshudu (1999). Long-term effect of nutrient management and cropping system on cotton yield and soil fertility in rainfed vertisols. *Nutrient Cycling in Agroecosystems*. Oct. 55(2): 159-164. {a} Central Institute for Cotton Research, Shankar Nagar (PO), Nagpur, 440 010, India

Vreeland, J. M., Jr. (1999). The revival of colored cotton. *Scientific American*. April 280(4): 112-118.

Walker, N. R., T. L. Kirkpatrick, et al. (1999). Effect of temperature on and histopathology of the interaction between *Meloidogyne incognita* and *Thielaviopsis basicola* on cotton. *Phytopathology* . Aug. 89(8): 613-617. {a} Department of Plant Pathology, Southwest Research and Extension Center, University of Arkansas, Hope, AR, 71801, USA

Wallace, T. P. (1999). Small plot evaluation of an electro-optical cotton yield monitor. *Computers and Electronics in Agriculture*. [print] June 23(1): 1-8. {a} Mississippi Agricultural and Forestry Experiment Station, Mississippi State, MS, 39762, USA

Wang, B., M. L. Dale, et al. (1999). Studies on a pathogenicity assay for screening cotton germplasms for resistance to *Fusarium oxysporum* f. sp. *vasinfectum* in the glasshouse. *Australian Journal of Experimental Agriculture* 39(8): 967-974. {a} Cooperative Research Centre for Tropical Plant Pathology, Department of Botany, The University of Queensland, Brisbane, Qld, 4072, Australia

Wang, B., M. L. Dale, et al. (1999). Effects of plant residue, soil characteristics, cotton cultivars and other crops on fusarium wilt of cotton in Australia. *Australian Journal of Experimental Agriculture* 39(2): 203-209. {a} Cooperative Research Centre for Tropical Plant Pathology, Department of Botany, The University of Queensland, Brisbane, Qld, 4072, Australia

Wang, W., W. X. Chen, et al. (1999). Studies on highly efficient planting of transgenic cotton. *Acta Botanica Sinica*. [print] Oct 41(10): 1072-1075. {a} Institute of Genetics, The Chinese Academy of Sciences, Beijing, 100101, China

Wang, W., Z. Zhu, et al. (1999). Obtaining a transgenic upland cotton harboring two insecticidal genes. *Acta Botanica Sinica*. April 41(4): 384-388. {a} Institute of Genetics, Chinese Academy of Sciences, Beijing, 100101, China

Warwick, S. I., H. J. Beckie, et al. (1999). Transgenic crops: New weed problems for Canada? *Phytoprotection* . [print] Aug 80(2): 71-84. {a} Eastern Cereal and Oilseed Research Centre, Central Experimental Farm, Agriculture and Agri-Food Canada, Ottawa, ON, K1A 0C6, Canada

Weis, K. G., K. R. Jacobsen, et al. (1999). Cytochemistry of developing cotton fibers: A hypothesized relationship between motes and non-dyeing fibers. *Field Crops Research*. June 62(2-3): 107-117. {a} Department of Agronomy and Range Science, University of California, Davis, CA, 95616, USA

Wheeler, T. A., J. R. Gannaway, et al. (1999). Identification of resistance to *Thielaviopsis basicola* in diploid cotton. *Plant Disease*. Sept. 83(9): 831-833. {a} Texas Agricultural Experiment Station, Rt. 3, Lubbock, TX, 79401, USA

Wheeler, T. A., H. W. Kaufman, et al. (1999). Comparison of variable and single-rate applications of aldicarb on cotton yield in fields infested with *Meloidogyne incognita*. *Journal of Nematology*. Dec. 31(4 Suppl): 700-708. {a} Texas Agricultural Experiment Station, Rt. 3, Lubbock, TX, 79401, USA

Whittaker, D. J. and B. A. Triplett (1999). Gene-specific changes in alpha-tubulin transcript accumulation in developing cotton fibers. *Plant Physiology Rockville*. Sept. 121(1): 181-188. {a} Cotton Fiber Bioscience, Southern Regional Research Center, United States Department of Agriculture-Agricultural Research Service, 1100 Robert E. Lee Boulevard, New Orleans, LA, 70124, USA

Wood, M. L., D. S. Murray, et al. (1999). Full-season interference of *Ipomoea hederacea* with *Gossypium hirsutum*. *Weed Science*. Nov. Dec. 47(6): 693-696. {a} Department of Plant and Soil Sciences, Oklahoma State University, Stillwater, OK, 74078, USA

Wright, P. R. (1999). Premature senescence of cotton (*Gossypium hirsutum* L.): Predominantly a potassium disorder caused by an imbalance of source and sink. *Plant and Soil* 211(2): 231-239. {a} CRC for Sustainable Cotton Production, Australian Cotton Research Institute, Myall Vale Mail Run, Narrabri, NSW, 2390, Australia

Wright, R. J., P. M. Thaxton, et al. (1999). Molecular mapping of genes affecting pubescence of cotton. *Journal of Heredity* 90(1): 215-219. {a} Dep. Crop and Soil Sci., Univ. Ga., Athens, GA 30602, USA

Xia, J. Y., d. W. W. van, et al (1999). Influence of temperature on bionomics of cotton aphid, *Aphis gossypii*, on cotton. *Entomologia Experimentalis et Applicata*. Jan. 90(1): 25-35. {a} Department of Theoretical Production Ecology, Wageningen Agricultural University, 6700 AK, Wageningen, Netherlands

Xia, J. Y., D. W. W. Van, et al. (1999). Temperature and prey density on bionomics of *Coccinella septempunctata* (Coleoptera: Coccinellidae) feeding on *Aphis gossypii* (Homoptera: Aphididae) on cotton. *Environmental Entomology*. April 28(2): 307-314. {a} Department of Theoretical Production Ecology, Wageningen Agricultural University, 6700 AK, Wageningen, Netherlands

Yee, J. and W. Ferguson (1999). Cotton pest management strategies and related pesticide use and yield. *Journal of Production Agriculture*. Oct. Dec. 12(4): 618-623. {a} USDA Economic Research Service, 1800 M Street NW, Washington, DC, 20036-5831, USA

Yoder, D. W., M. Nampaisansuk, et al. (1999). Molecular cloning and nucleotide sequence of a gene encoding a cotton palmitoyl-acyl carrier protein thioesterase. *Biochimica et Biophysica Acta*. Sept. 3(3): 403-413. {a} Department of Biological Sciences, University of North Texas, Denton, TX, 76203-5220, USA

Zapata, C., S. H. Park, et al. (1999). Transformation of a Texas cotton cultivar by using *Agrobacterium* and the shoot apex. *Theoretical and Applied Genetics* 98(2): 252-256. {a} Department of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843, USA

Zapata, C., M. Srivatanakul, et al. (1999). Improvements in shoot apex regeneration of two fiber crops: Cotton and kenaf. *Plant Cell Tissue and Organ Culture* 56(3): 185-191. {a} Department of Soil and Crop Sciences, Texas A and M University, College Station, TX, 77843-2474, USA

Zeng, F., S. B. Ramaswamy, et al. (1999). Comparison of monoclonal antibody and laboratory rearing techniques to identify *Heliothentinae* (Lepidoptera: Noctuidae) eggs from Mississippi cotton fields. *Environmental Entomology*. April 28(2): 275-281. {a} Department of Entomology, Kansas State University, 123 Waters Hall, Manhattan, KS, 66506, USA

Zeringue, H. J., Jr., B. Y. Shih, et al. (1999). Identification of the bright-greenish-yellow-fluorescence (BGY-F) compound on cotton lint associated with aflatoxin contamination in cottonseed. *Phytochemistry* Oxford. Dec. 52(8): 1391-1397. {a} Southern Regional Research Center, U.S. Department of Agriculture, Agricultural Research Service, New Orleans, LA, 70179-0687, USA

Zhao, D. and D. M. Oosterhuis (1999). Photosynthetic capacity and carbon contribution of leaves and bracts to developing floral buds in cotton. *Photosynthetica Prague* 36(1-2): 279-290. {a} Altheimer Laboratory, Department of Agronomy, University of Arkansas, 276 Altheimer Drive, Fayetteville, AR, 72704, USA

- Zhao, D. and D. M. Oosterhuis (1999). Dynamics of mineral nutrient element concentrations in developing cotton leaves, bracts, and floral buds in relation to position in the canopy. *Journal of Plant Nutrition*. July 22(7): 1107-1122. {a} Department of Agronomy, 276 Altheimer Laboratory, University of Arkansas, Fayetteville, AR, 72704, USA
- Zodape, S. T. (1999). Influence of pulsed concentrated solar radiation on phytase activity in germinating seeds of cotton. *Indian Journal of Plant Physiology*. April June 4(2): 144-146. {a} Central Salt and Marine Chemicals Research Institute, Bhavnagar, 364002, India
- Abdullaev, A., Z. N. Abdurakhmanova, et al. (1998). Effect of exogenous phosphoenolpyruvate on photosynthetic metabolism of carbon in cotton leaves. *Fiziologiya Rastenii Moscow* 45(1): 28-32. {a} Inst. Plant Physiol. Genet., Acad. Sci. Tajikistan, ul. S. Aini 299/2, Dushanbe 734063, Tajikistan
- Abrao, M. M. and P. Mazzafera (1998). Physiological alterations in cotton due to infection by *Meloidogyne incognita* race 3: Influence of nitrogen fertilization. *Nematologia Brasileira*. Dec. 22(2): 66-79. {a} Aluna de Pos-Graduacao do Departamento de Fisiologia Vegetal IB, Unicamp, 13083-970, Campinas, SP, Brazil
- Adamczyk, J. J., Jr., J. W. Holloway, et al. (1998). Larval survival and development of the fall armyworm (Lepidoptera: Noctuidae) on normal and transgenic cotton expressing the *Bacillus thuringiensis* CryIA (c) variant delta-endotoxin. *Journal of Economic Entomology* 91(2): 539-545. {a} Dep. Entomol., Louisiana State Univ. Agric. Cent., 402 Life Sci. Build., Baton Rouge, LA 70803, USA
- Agrawal, D. C., A. K. Banerjee, et al. (1998). Effect of cefotaxime on the growth of excised embryoaxes of six cultivars of cotton (*Gossypium hirsutum* L.). *Journal of Plant Physiology* 152(4-5): 580-582. Plant Tissue Culture Div., Natl. Chem. Lab., Pune-411008 India
- Ahuja, S. L., S. K. Banerjee, et al. (1998). Genotype X environment interaction of morphotypes in cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research*. June 32(2): 93-100. {a} Central Institute for Cotton Research Regional Station, Sirsa, HR, 125 055, India
- Ahuja, S. L., D. Singh, et al. (1998). Genetic components of variation for physiological attributes in early types of *Gossypium hirsutum* cotton. *Agricultural Science Digest* 18(1): 12-14. {a} Regional Station, CICR, Sirsa, Haryana, 125 055, India
- Akey, D. H. and T. J. Henneberry (1998). Beet armyworm (Lepidoptera: Noctuidae) behavior and effects on insecticide efficacy in open and closed cotton flowers and boll abscission or damage from flower infestation. *Journal of Economic Entomology* 91(1): 263-266. Western Cotton Res. Lab., USDA-ARS, Pacific West Area, 4135 E. Broadway, Phoenix, AZ 85040-8803, USA
- Andersland, J. M., D. C. Dixon, et al. (1998). Isolation and characterization of cytoskeletons from cotton fiber cytoplasts. *In Vitro Cellular and Developmental Biology Plant* 34(3): 173-180. {a} USDA-ARS, Cotton Fiber Bioscience, Southern Regional Res. Centre, P.O. Box 19687, New Orleans, LA 70179, USA
- Anonymous (1998). New adventitious species in Mordovia. *Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody Otdel Biologicheskii*. Nov. Dec. 103(6): 59-60. Cambridge, UK
- Austin, N., M. U. D. Khan, et al. (1998). Genetic studies of cotton (*Gossypium hirsutum* L) I. Combining ability and heterosis studies in yield and yield components. *Pakistan Journal of Scientific and Industrial Research* 41(1): 54-56. {a} Cotton Research Station, Old Shujabad Road, Multan, Pakistan
- Ayliffe, M. A., N. S. Scott, et al. (1998). Analysis of plastid DNA-like sequences within the nuclear genomes of higher plants. *Molecular Biology and Evolution* 15(6): 738-745. {a} Dep. Genetics, Univ. Adelaide, Adelaide, South Australia 5005, Australia
- Babu, R. (1998). Cotton (*Gossypium hirsutum*)-based cropping systems in Ghataprabha command area of Karnataka. *Indian Journal of Agronomy* 43(2): 261-263. {a} Agric. Res. Stn., Bidar 585 401, India

- Baisakh, B., V. K. Khanna, et al. (1998). Influence of time of application of growth regulators on boll retention and set of upland cotton (*Gossypium hirsutum*) and tree cotton (*Gossypium arboreum*) crosses. *Indian Journal of Agricultural Sciences* 68(5): 264-267. {a} NARP, Bhawanipatna, Orissa 766 001, India
- Baisakh, B., V. K. Khanna, et al. (1998). Standardization of the best timing of growth regulator application in *Gossypium hirsutum* and *G. arboreum* crosses to increase boll formation and seed set. *Acta Agronomica Hungarica* 46(1): 9-14. Dep. Genet. Plant Breed., NARP, Bhawanipatana, Kalahandi, Orissa 766001, India
- Balasubramanian, A., N. R. Mahadeven, et al. (1998). Influence of intercropping on infestation of early season sucking pests of cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 68(6): 315-316. {a} Agric. Coll. Res. Inst., Tamil Nadu Agric. Univ., Madurai, Tamil Nadu 625 104, India
- Bauer, P. J., O. L. May, et al. (1998). Planting date and potassium fertility effects on cotton yield and fiber properties. *Journal of Production Agriculture* 11(4): 415-420. {a} USDA-ARS Coastal Plains Soil Water and Plant Res. Cent., 2611 W. Lucas St., Florence, SC 29501-1242, USA
- Bednarz, C. W. and G. J. Gascho (1998). Effects of 'Carpramid' on nutrient uptake and lint yield of cotton. *Journal of Plant Nutrition* 21(11): 2417-2427. {a} Dep. Crop Soil Sci., Univ. Ga. Coastal Plain Exp. Stn., P.O. Box 748, Tifton, GA 31793-0748, USA
- Bednarz, C. W., N. W. Hopper, et al. (1998). Effects of foliar fertilization of Texas southern high plains cotton: Leaf nitrogen and growth parameters. *Journal of Production Agriculture* 11(1): 80-84. {a} Univ. Ga., Coastal Plain Exp. Stn., P.O. Box 748, Tifton, GA 31793, USA
- Bednarz, C. W. and D. M. Oosterhuis (1998). Development of a protocol to study the effects of potassium deficiency in cotton under controlled environmental conditions. *Journal of Plant Nutrition* 21(2): 329-339. {a} Crop Soil Sci. Dep., Univ. Ga., Coastal Plain Exp. Stn., P.O. Box 748, Tifton, GA 31793-0748, USA
- Bednarz, C. W., D. M. Oosterhuis, et al. (1998). Leaf photosynthesis and carbon isotope discrimination of cotton in response to potassium deficiency. *Environmental and Experimental Botany* 39(2): 131-139. {a} Univ. Ga., Coastal Plain Experiment Station, Crop Soil Science Dep., P.O. 748, Tifton, GA 31793, USA
- Bondada, B. R. and D. M. Oosterhuis (1998). Relationships between nitrogen content and net gas exchange components of a cotton leaf during ontogeny. *Photosynthetica Prague* 35(4): 631-635. {a} Univ. Tennessee, Dep. Plant Soil Science, P.O. Box 1071, Knoxville, TN 37901, USA
- Bondada, B. R. and D. M. Oosterhuis (1998). Decline in photosynthesis as related at alterations in chloroplast ultrastructure of a cotton leaf during ontogeny. *Photosynthetica Prague* 35(3): 467-471. {a} Univ. Tennessee, Dep. Plant Soil Sci., P.O. Box 1071, Knoxville, TN 37901, USA
- Bourland, F. M., R. E. McGowen, Jr., et al. (1998). Registration of Arkot A129 and Arkot A132 germplasm lines of cotton with high fiber strength. *Crop Science* 38(2): 567-568. {a} Northeast Res. Ext. Cent., P.O. Box 96, Keiser, AR 72351, USA
- Bowman, D. T. (1998). Using crop performance data to select hybrids and varieties. *Journal of Production Agriculture* 11(2): 256-259. {a} Dep. Crop Soil Sci., N.C. State Univ., Raleigh, NC 27695-8604, USA
- Brar, K. S., B. S. Sandhu, et al. (1998). Tissue culture response of cultivated and wild cotton species. *Crop Improvement* 25(1): 59-65. PAU Regional Res. Stn., Bathinda 151 001, India
- Brisson, L. F., I. Zelitch, et al. (1998). Manipulation of catalase levels produces altered photosynthesis in transgenic tobacco plants. *Plant Physiology Rockville* 116(1): 259-269. {a} Dep. Biochemistry Genetics, Connecticut Agricultural Experiment Station, P.O. Box 1106, New Haven, CT 06504, USA
- Charles, G. W., R. D. Murison, et al. (1998). Competition of noogoora burr (*Xanthium occidentale*) and fierce thornapple (*Datura ferox*) with cotton (*Gossypium hirsutum*). *Weed Science* 46(4): 442-446. {a} NSW Agric., Aust. Cotton Res. Inst., Narrabri, NSW 2390, Australia

Chemengich, B. T. and B. M. Khaemba (1998). Comparative studies on biological performance of the cotton stainers *Dysdercus cardinalis* Gerst and *Dysdercus fasciatus* Signoret bred on different hosts. *Insect Science and its Application*. Jan. March 18(1): 25-29. {a} Department of Zoology, Kenyatta University, Nairobi, Kenya

Chitarra, L. G., A. B. Chitarra, et al. (1998). Neutral sugars, uronic acids and electric conductivity of cottonseeds (*Gossypium hirsutum* L.) exudate in function of the chemical delinting. *Brazilian Archives of Biology and Technology*. March 41(1): 129-143. {a} Departamento de Fitossanidade, Universidade Federal de Lavras, CEP 37200-000, Lavras, MG, USA

Chu, C. C., E. T. Natwick, et al. (1998). Effect's of aldicarb on cotton insects and plant growth and yield. *Journal of the Agricultural Association of China*. Dec.(184): 98-108. {a} Western Cotton Research Laboratory, USDA-ARS, 4135 E. Broodway Rd., Phoenix, AZ, 85040, USA

Conteh, A., G. J. Blair, et al. (1998). Soil organic carbon fractions in a Vertisol under irrigated cotton production as affected by burning and incorporating cotton stubble. *Australian Journal of Soil Research* 36(4): 655-656. {a} CRC Sustainable Cotton Production, Div. Agron. Soil Sci., Sch. Rural Sci. Nat. Resource, Univ. New Engl., Armidale, NSW 2351, Australia

Corkern, C. B., D. B. Reynolds, et al. (1998). Bromoxynil antagonizes Johnsongrass (*Sorghum halepense*) control with graminicides. *Weed Technology* 12(2): 205-208. {a} Dep. Plant Pathol. Crop Physiol., Baton Rouge, LA 70803, USA

Cortesero, A. M., M. C. M. De, et al. (1998). Correction of PREVIEWS 99653084. Comparisons and contrasts in host-foraging strategies of two larval parasitoids with different degrees of host specificity. Correction of author name from C. M. D. Moraes. *Journal of Chemical Ecology* 24(11): 1589-1606. {a} Insect Biol. Population Manage. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Tifton, GA 31793, USA

Culpepper, A. S., A. C. York, et al. (1998). Interaction of bromoxynil and postemergence graminicides on large crabgrass (*Digitaria sanguinalis*). *Weed Technology* 12(3): 554-559. Crop Sci. Dep., N.C. State Univ., Box 7620, Raleigh, NC 27695-7620, USA

Daayf, F., M. Nicole, et al. (1998). Hyaline mutants from *Verticillium dahliae*, an example of selection and characterization of strains for host-parasite interaction studies. *Plant Pathology Oxford* 47(4): 523-529. {a} Agric. Agri-Food Canada, 440 University Ave., P.O. Box 1210, Charlottetown, PE C1A 7M8, Canada

de, M. E. N., T. R. M. A. Gondim, et al. (1998). Cotton fiber number and convolution angle measurement and its relation to yarn and fiber technological properties. *Bragantia* 57(2): 279-283. {a} Centro de Algodao e Fibras Diversas, Instituto Agronomico (IAC), 13001-970, Campinas, SP, Brazil

Dolgikh, Y. I., E. Y. Shaikina, et al. (1998). Synthetic trisaccharide fragment of xyloglucan as regulator of plant morphogenesis. *Doklady Akademii Nauk* 360(3): 417-419. {a} K. A. Timiryazev Inst. Plant Physiol., Russ. Acad. Sci., Moscow, Russia

El, Z. K. M. and P. M. Thaxton (1998). Registration of five glabrous multi-adversity resistant (MAR-6) germplasm lines of upland cotton. *Crop Science* 38(6): 1730. {a} Dep. Soil Crop Sci., Tex. Agric. Exp. Stn., Tex. A and M Univ. Syst., College Station, TX 77843-2474, USA

El, Z. K. M. and P. M. Thaxton (1998). Registration of seven pubescent multi-adversity resistant (MAR-6) germplasm lines of upland cotton. *Crop Science* 38(6): 1729. {a} Dep. Soil Crop Sci., Tex. Agric. Exp. Stn., Tex. A and M Univ. System, College Station, TX 77843-2474, USA

Elena, K. and E. J. Paplomatas (1998). *Phytophthora boehmeriae* Boll Rot: A new threat to cotton cultivation in the Mediterranean region. *Phytoparasitica* 26(1): 20-26. Benaki Phytophthological Inst., 14561 Kifissia, Athens, Greece

Feller, U., B. S. J. Crafts, et al. (1998). Moderately high temperatures inhibit ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) activase-mediated activation of Rubisco. *Plant Physiology Rockville*

116(2): 539-546. {a} U.S. Dep. Agric.-Agric. Res. Serv. Western Cotton Res. Lab., 4135 E. Broadway Rd., Phoenix, AZ 85040-8830, USA

Ganieva, R. A., S. R. Allahverdiyev, et al. (1998). Effect of salt stress and synthetic hormone polystimuline K on the photosynthetic activity of cotton (*Gossypium hirsutum*). Turkish Journal of Botany 22(4): 217-221. {a} Inst. Bot., Acad. Sci., Baku, Azerbaijan

Gao, Y. F., Z. Zhu, et al. (1998). Isolation of soybean Kunitz trypsin inhibitor gene and its application in plant insect-resistant genetic engineering. Acta Botanica Sinica 40(5): 405-411. {a} Inst. Genet., Chinese Acad. Sci., Beijing 100101, China

Giri, A. N., R. H. Bhosle, et al. (1998). Weed management in cotton (*Gossypium hirsutum*)-sunflower (*Helianthus annuus*) cropping sequence. Indian Journal of Agronomy 43(2): 264-268. {a} Cotton Res. Scheme, MAU, Parbhani, Maharashtra 431 402, India

Giri, A. N., R. H. Bhosle, et al. (1998). Weed management in cotton (*Gossypium hirsutum*)-groundnut (*Arachis hypogaea*) sequence. Indian Journal of Agronomy 43(1): 50-56. All-India Co-ordinated Res. Programme Weed Control, Dep. Agron., Marathwada Agric. Univ., Parbhani Maharashtra 431 402, India

Gokani, S. J., R. Kumar, et al. (1998). Potential role of abscisic acid in cotton fiber and ovule development. Journal of Plant Growth Regulation 17(1): 1-5. {a} Dep. Biosci., Saurashtra Univ., Rajkot 360 005, India

Goolsby, J. A., M. A. Ciomperlik, et al. (1998). Laboratory and field evaluation of exotic parasitoids of *Bemisia tabaci* (Gennadius) (Biotype "B") (Homoptera: Aleyrodidae) in the Lower Rio Grande Valley of Texas. Biological Control 12(2): 127-135. {a} USDA-APHIS-PPQ Mission Plant Protection Cent., P.O. Box 2140, Mission, TX 78573, USA

Gordh, G. (1998). A new species of *Sierola* parasitic on moth larvae in Western Australia (Hymenoptera: Bethyridae). Proceedings of the Hawaiian Entomological Society 33: 83-88. {a} Dep. of Entomology, Univ. of Queensland, Brisbane, QLD 4072, Australia

Groves, R. L., G. G. Kennedy, et al. (1998). Inoculation of tomato spotted wilt virus into cotton. Plant Disease 82(8): 959. {a} Dep. Entomol., North Carolina State Univ., Raleigh, NC 27695, USA

Guertal, E. A., A. O. Abaye, et al. (1998). Boron uptake and concentration in cotton and soybean as affected by boron source. Communications in Soil Science and Plant Analysis 29(19-20): 3007-3014. {a} Dep. Agron. Soils, Auburn Univ., Auburn, AL 36849, USA

Gupta, G. P., G. K. Mahapatro, et al. (1998). Partial larval rearing of *Earias vittella* (Fabricius) on artificial diet. Insect Science and its Application. Oct. Dec. 18(4): 371-375. {a} Division of Entomology, Indian Agricultural Research Institute, New Delhi, 110012, India

Gwathmey, C. O. and D. D. Howard (1998). Potassium effects on canopy light interception and earliness of no-tillage cotton. Agronomy Journal 90(2): 144-149. {a} Dep. Plant Soil Sci., Univ. Tennessee, 605 Airways Blvd., Jackson, TN 38301, USA

Hafeez, U. R. W. S. K. (1998). Expressivity of H2 gene of hairiness and L0 gene of leaf shape of cotton under different genetic backgrounds. Pakistan Journal of Botany 30(1): 95-100. {a} Cotton Res. Inst., Faisalabad, Pakistan

Hallmann, J., H. A. Quadt, et al. (1998). Interactions between *Meloidogyne incognita* and endophytic bacteria in cotton and cucumber. Soil Biology and Biochemistry 30(7): 925-937. {a} Dep. Plant Pathol., Biol. Control Inst., Alabama Agricultural Exp. Station, Auburn, AL 36849-5409, USA

Hanson, R. E., X. P. Zhao, et al. (1998). Evolution of interspersed repetitive elements in *Gossypium* (Malvaceae). American Journal of Botany 85(10): 1364-1368. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA

Hattori, J., K. A. Boutilier, et al. (1998). A conserved BURP domain defines a novel group of plant proteins with unusual primary structures. *Molecular and General Genetics* 259(4): 424-428. {a} East. Cereal and Oilseed Res. Cent., Agric. Agri-Food Can., Room 2091 K.W. Neatby Build., Ottawa, ON K1A 0C6, Canada

Heitholt, J. J. and W. R. Meredith, Jr. (1998). Yield, flowering, and leaf area index of okra-leaf and normal-leaf cotton isolines. *Crop Science* 38(3): 643-648. {a} USDA-ARS, P.O. Box 345, Stoneville, MS 38776-0345, USA

Heitholt, J. J., W. R. Meredith, Jr., et al. (1998). Leaf area index response of four obsolete and four modern cotton cultivars to two nitrogen levels. *Journal of Plant Nutrition* 21(11): 2319-2328. USDA-ARS, Cotton Physiol. Genet., P.O. Box 345, Stoneville, MS 38776, USA

Hemphill, J. K., C. G. A. Maier, et al. (1998). Rapid in vitro plant regeneration of cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 17(4): 273-278. {a} Cottonseed Dev. Group, Dep. Biol. Sci., Univ. North Tex., Denton, TX 76203-5220, USA

Henneberry, T. J., L. F. Jech, et al. (1998). *Bemisia argentifolii* (Homoptera: Aleyrodidae) population relationships to cotton and lint stickiness in long and short staple cottons. *Journal of Economic Entomology* 91(5): 1196-1207. {a} West. Cotton Res. Lab., USDA-ARS, Phoenix, AZ, USA

Hoelmer, K. A., W. J. Roltsch, et al. (1998). Selectivity of whitefly traps in cotton for *Eretmocerus eremicus* (Hymenoptera: Aphelinidae), a native parasitoid of *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Environmental Entomology* 27(4): 1039-1044. {a} Phoenix Plant Prot. Cent., Irrigated Desert Res. Stn., U.S. Dep. Agric., APHIS, PPQ, 4151 Highway 86, Brawley, CA 92227, USA

Hood, M. A., D. K. V. Van, et al. (1998). Factors affecting attachment of *Enterobacter cloacae* to germinating cotton seed. *Microbial Ecology* 36(1): 101-110. {a} Univ. West Florida, Dep. Biol., Pensacola, FL 32514, USA

Howard, D. D., C. O. Gwathmey, et al. (1998). Potassium fertilization of cotton produced on a low K soil with contrasting tillage systems. *Journal of Production Agriculture* 11(1): 74-79. {a} Plant Soil Sci. Dep., Univ. Tenn., 605 Airways Blvd., Jackson, TN 38301, USA

Howard, D. D., C. O. Gwathmey, et al. (1998). Foliar feeding of cotton: Evaluating potassium sources, potassium solution buffering, and boron. *Agronomy Journal* 90(6): 740-746. {a} Plant Soil Sci. Dep., West Tenn. Exp. Stn., 605 Airways Blvd., Jackson, TN 38301, USA

Hulugalle, N. R., P. C. Entwistle, et al. (1998). Effect of long-fallow on soil quality and cotton lint yield in an irrigated, self-mulching, gray Vertosol in the central-west of New South Wales. *Australian Journal of Soil Research* 36(4): 621-639. {a} Aust. Cotton Res. Inst., NSW Agric., Myall Vale Mail Run, Narrabi, NSW 2390, Australia

Hunsaker, D. J., A. J. Clemmens, et al. (1998). Cotton response to high frequency surface irrigation. *Agricultural Water Management* 37(1): 55-74. {a} USDA, Agric. Res. Serv., U.S. Water Conservation Lab., 4331 E. Broadway Road, Phoenix, AZ 85040, USA

Hunt, P. G., P. J. Bauer, et al. (1998). Nitrogen accumulation in cotton grown continuously or in rotation with peanut using subsurface microirrigation and GOSSYM/COMAX management. *Crop Science* 38(2): 410-415. {a} USDA-ARS, Coastal Plains Soil Water Plant Res. Cent., 2611 W. Lucas St., Florence, SC 29501-1241, USA

Ihara, Y., F. Sakai, et al. (1998). Cloning of cotton homologs of bcsA gene encoding cellulose beta-1,4-glucosyltransferase. *Wood Research*(85): 56-58. Lab. Gene Expression, Wood Res. Inst., Kyoto Univ., Uji, Kyoto 611, Japan

Jain, S. and V. N. Tiwari (1998). Heterosis and inbreeding depression in upland cotton (*G. hirsutum* L.). *Advances in Plant Sciences* 11(1): 95-97. JNKVV Res. Stn., Badnawar, India

Jiang, C. X., R. J. Wright, et al. (1998). Polyploid formation created unique avenues for response to selection in *Gossypium* (cotton). *Proceedings of the National Academy of Sciences of the United States of America* 95(8): 4419-4424. {a} Plant Genome Mapping Lab., Dep. Soil and Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA

Jones, M. A. and R. Wells (1998). Fiber yield and quality of cotton grown at two divergent population densities. *Crop Science* 38(5): 1190-1195. {a} Dep. Crop Sci., N.C. State Univ., Raleigh, NC 27695-7620, USA

Kai, H., Y. Horita, et al. (1998). Synthesis and herbicidal activities of 2-(5-isoxazolyloxy)acetamide derivatives. *Journal of Pesticide Science* 23(3): 255-261. Aburahi Lab., Shionogi and Co. Ltd., 1405 Gotanda, Koka-cho, Koka-gun, Shiga 520-3423, Japan

Kasumov, N. A., Z. I. Abbasova, et al. (1998). Effects of salt stress of the respiratory components of some plants. *Turkish Journal of Botany* 22(6): 389-396. {a} Fac. Biol., Baku State Univ., Baku, Azerbaijan

Kawai, M., S. Aotsuka, et al. (1998). Isolation of a cotton CAP gene: A homologue of adenylyl cyclase-associated protein highly expressed during fiber elongation. *Plant and Cell Physiology* 39(12): 1380-1383. {a} Inst. Mol. Cell. Biosci., Univ. Tokyo, Yayoi 1-1-1, Bunkyo-ku, Tokyo 113-0032, Japan

Kerby, T. A., R. E. Plant, et al. (1998). Environmental and cultivar effects on height-to-node ratio and growth rate in acala cotton. *Journal of Production Agriculture* 11(4): 420-427. {a} Dep. Agronomy Range Sci., Univ. California, Davis, CA 95616, USA

Kinloch, R. A. and J. R. Rich (1998). Responses of cotton yield and *Meloidogyne incognita* soil populations to soil applications of aldicarb and 1,3-D in Florida. *Journal of Nematology*. Dec. 30(4 Suppl): 639-642. {a} University of Florida, IFAS-WFREC, Jay, FL, 32351, USA

Kudachikar, V. B. and B. S. Janagouder (1998). Physiological basis for higher productivity in upland cotton under rainfed conditions. *Journal of Maharashtra Agricultural Universities*. May 23(2): 187-188. {a} Research Institute, Central Food Technological, Mysore, 570 013, India

Kumar, S. and D. Pental (1998). Regeneration of Indian cotton variety MCU-5 through somatic embryogenesis. *Current Science Bangalore* 74(6): 538-540. {a} Dep. Genet., Univ. Delhi South Campus, Benito Juarez Rd., New Delhi 110 021, India

Kumar, S., P. Sharma, et al. (1998). A genetic approach to in vitro regeneration of non-regenerating cotton (*Gossypium hirsutum* L.) cultivars. *Plant Cell Reports* 18(1-2): 59-63. {a} Dep. Genet., Univ. Delhi S. Campus, Benito Juarez Rd., New Delhi 110 021, India

Lacape, M. J., J. Wery, et al. (1998). Relationships between plant and soil water status in five field-grown cotton (*Gossypium hirsutum* L.) cultivars. *Field Crops Research* 57(1): 29-43. {a} CIRAD, Ave. Agropolis, BP 5035, 34032 Montpellier Cedex 1, France

Leys, J. F., F. J. Larney, et al. (1998). Anthropogenic dust and endosulfan emissions on a cotton farm in northern New South Wales, Australia. *Science of the Total Environment* 220(1): 55-70. {a} Dep. Land Water Conservation, Centre Natural Resources, Gunnedah Research Centre, P.O. Box 462, Gunnedah, NSW 2380, Australia

Liakatas, A., D. Roussopoulos, et al. (1998). Controlled-temperature effects on cotton yield and fibre properties. *Journal of Agricultural Science* 130(4): 463-471. {a} Agric. Univ. Athens, Dep. Land Reclam. Agric. Eng., 75 Iera Odos, GR-118 55 Athens, Greece

Liu, C. J., Y. L. Meng, et al. (1998). Cloning and sequencing of a cDNA encoding farnesyl pyrophosphate synthase from *Gossypium arboreum* and its expression pattern in the developing seeds of *Gossypium hirsutum* cv. "Sumian-6". *Acta Botanica Sinica* 40(8): 703-710. {a} Natl. Lab. Plant Mol. Genet., Shanghai Inst. Plant Physiol., Chinese Acad. Sci., Shanghai 200032, China

Loguercio, L. L. and T. A. Wilkins (1998). Structural analysis of a hmg-coA-reductase pseudogene: Insights into evolutionary processes affecting the hmgr gene family in allotetraploid cotton (*Gossypium hirsutum* L.). *Current Genetics* 34(4): 241-249. {a} Dep. Agronomy Range Sci., Univ. Calif., One Shields Ave., Davis, CA 915616-8515, USA

Lu, Z., R. G. Percy, et al. (1998). Stomatal conductance predicts yields in irrigated pima cotton and bread wheat grown at high temperatures. *Journal of Experimental Botany* 49(Spec. Issue): 453-460. {a} Dep. Biol., Univ. California Los Angeles, Los Angeles, CA 90095-1606, USA

Maas, S. J. (1998). Estimating cotton canopy ground cover from remotely sensed scene reflectance. *Agronomy Journal* 90(3): 384-388. {a} USDA-ARS, 17053 Shafter Ave., Shafter, CA 93263, USA

Mahakulkar, B. V., S. S. Wanjari, et al. (1998). Integrated nutrient management in sorghum (*Sorghum bicolor*)-based cropping system. *Indian Journal of Agronomy*. Sept. 43(3): 376-381. {a} Sorghum Research Unit, Punjabrao Krishi Vidyapeeth, Akola, MAH, 444 104, India

Maldes, J. M. and A. Ratnadass (1998). *Crotalaria juncea* (Fabaceae) cited for the first time as alternative host of *Eurystylus oldi* important pest of sorghum and ricinus (Hem., Miridae). *Bulletin de la Societe Entomologique de France* 103(3): 272. CIRAD-AMIS, Protection Cultures, ENTOTROP, BP 5035, F-34032 Montpellier Cedex 1, France

Maliwal, G. L., A. Das, et al. (1998). Water management through land configuration in Bhal region. *Indian Journal of Agricultural Sciences* 68(12): 762-764. {a} National Agricultural Research Project, Gujarat Agricultural University, Arnej, 382 230, India

Marur, C. J. (1998). Photosynthesis and translocation of sugars in cotton plants subjected to drought stress after mepiquat chloride application. *Revista Brasileira de Fisiologia Vegetal* 10(1): 59-64. {a} Inst. Agron. Parana, C.P. 481, Londrina, PR 86001-970, Brazil

Mascarenhas, R. N., D. J. Boethel, et al. (1998). Resistance monitoring to *Bacillus thuringiensis* insecticides for soybean loopers (Lepidoptera: Noctuidae) collected from soybean and transgenic Bt-cotton. *Journal of Economic Entomology* 91(5): 1044-1050. {a} Zeneca Ag Prod., East. Regional Tech. Cent., 16013 Watson Seed Farm Road, Whitakers, NC 27891, USA

Mascarenhas, V. J., J. B. Graves, et al. (1998). Susceptibility of field populations of beet armyworm (Lepidoptera: Noctuidae) to commercial and experimental insecticides. *Journal of Economic Entomology* 91(4): 827-833. La. Agricultural Experiment Station, La. State Univ. Agricultural Cent., Baton Rouge, LA 70803, USA

McAuslane, H. J. and H. T. Alborn (1998). Systemic induction of allelochemicals in glanded and glandless isogenic cotton by *Spodoptera exigua* feeding. *Journal of Chemical Ecology* 24(2): 399-416. {a} Dep. Entom. Nematol., Univ. Florida, PO Box 110620, Gainesville, FL 32611-0620, USA

McCarty, J. C., Jr., J. N. Jenkins, et al. (1998). Introgression of day-neutral genes in primitive cotton accessions: II. Predicted genetic effects. *Crop Science* 38(6): 1428-1431. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

McCarty, J. C., Jr., J. N. Jenkins, et al. (1998). Introgression of day-neutral genes in primitive cotton accessions: I. Genetic variances and correlations. *Crop Science* 38(6): 1425-1428. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

McLean, K. S. and G. W. Lawrence (1998). First report of premature boll rot associated with cotton in Louisiana. *Plant Disease* 82(6): 710. {a} Dep. Agric., Northeast La. Univ., Monroe, LA 71209, USA

Mehetre, S. S., V. R. Patil, et al. (1998). Gamma rays induced branching habit mutant in *Gossypium hirsutum* L. cotton. *Crop Research Hisar* 16(2): 282-283. All India Co-ordinated Cotton Improvement Project, Mahatma Phule Krishi Vidyapeeth, Rahuri-413 722, India

- Meredith, W. R., Jr. (1998). Registration of eight sub-okra, semi-smooth, and nectariless near-isolines of DES 119 cotton germplasm. *Crop Science* 38(6): 1725. {a} USDA-ARS Cotton Physiol. Genet. Res. Unit, Box 314, Stoneville, MS 38776, USA
- Moffett, J. E. and W. B. McCloskey (1998). Effects of soil moisture and yellow nutsedge (*Cyperus esculentus*) density on cotton (*Gossypium hirsutum*). *Weed Science* 46(2): 231-237. {a} Dep. Plant Sci., Univ. Ariz., Tucson, AZ 85721, USA
- Moghal, M. S. and J. C. Schneider (1998). Effects of high temperature on development of *Heliothis virescens* (F) on cotton fruiting structures in the laboratory. *Southwestern Entomologist* 23(2): 123-126. {a} Crop Sci. Res. Laboratory, USDA-ARS, Box 9774, Mississippi State, MS 39762, USA
- Monteiro, R. C., R. A. Zucchi, et al. (1998). Thrips *tabaci* Lind.: Is it a real cotton pest in Brazil? *Anais da Sociedade Entomologica do Brasil* 27(3): 489-494. {a} Dep. Entomol., ESALQ/USP, Caixa Postal 9, 13418-900 Piracicaba, SP, Brazil
- Morre, J. L., H. R. Permingeat, et al. (1998). Multiple shoot induction and plant regeneration from embryonic axes of cotton. *Plant Cell Tissue and Organ Culture* 54(3): 131-136. {a} Centro de Estudios Fotosinteticos y Bioquimicos, CEFOTI (UNR, CONICET, Fundacion Miguel Lillo), Suipacha 531, 2000, Rosario, Argentina
- Mullins, G. L. (1998). Cotton response to the rate and source of sulfur on a sandy coastal plain soil. *Journal of Production Agriculture* 11(2): 214-218. {a} Dep. Agron. Soils, 202 Funchess Hall, Auburn Univ., Auburn, AL 36849-5412, USA
- Munger, P., H. Bleiholder, et al. (1998). Phenological growth stages of the cotton plant (*Gossypium hirsutum* L.): Codification and description according to the BBCH scale. *Journal of Agronomy and Crop Science* 180(3): 143-149. {a} BASF-Landwirtschaftliche Versuchsstn., Carl-Bosch-Str. 64, D-67117 Limburgerhof, Germany
- Muratov, G. A., M. M. Rakhimov, et al. (1998). Enzymatic hydrolysis of cotton fibers. *Prikladnaya Biokhimiya i Mikrobiologiya* 34(5): 525-528. Tashkent State Univ., Tashkent 700095, Uzbekistan
- Nehl, D. B., S. J. Allen, et al. (1998). Slow arbuscular mycorrhizal colonisation of field-grown cotton caused by environmental conditions in the soil. *Mycorrhiza* 8(3): 159-167. {a} Australian Cotton Res. Inst., Myall Vale Mail Run, Narrabri, NSW 2390, Australia
- Nepomuceno, A. L., D. M. Oosterhuis, et al. (1998). Physiological responses of cotton leaves and roots to water deficit induced by polyethylene glycol. *Environmental and Experimental Botany* 40(1): 29-41. {a} Altheimer Lab., Agronomy Dep., Univ. Ark., Fayetteville, AR 72701, USA
- Norton, E. R. and J. C. Silvertooth (1998). Field validation of soil solute profiles in irrigated cotton. *Agronomy Journal* 90(5): 623-630. {a} Dep. Plant Sci., Univ. Ariz., Tucson, AZ 85721, USA
- Pan, X., J. D. Hesketh, et al. (1998). A data base for scaling up from the leaf to the canopy. *Photosynthetica Prague* 35(3): 329-334. {a} Crop Prot. Res. Unit ARS, USDA, 190 EMRL, 1201 West Gregory, Univ. Illinois, Urbana, IL 61801-3838, USA
- Parajulee, M. N., J. E. Slosser, et al. (1998). Seasonal activity of *Helicoverpa zae* and *Heliothis virescens* (Lepidoptera: Noctuidae) detected by pheromone traps in the rolling plains of Texas. *Environmental Entomology* 27(5): 1203-1219. {a} Tex. Agric. Exp. Stn., 11708 Highway 70 South, P.O. Box 1658, Vernon, TX 76385-1658, USA
- Pare, P. W. and J. H. Tumlinson (1998). Cotton volatiles synthesized and released distal to the site of insect damage. *Phytochemistry Oxford* 47(4): 521-526. {a} Cent. Med. Agric. Vet. Entomol., U.S. Dep. Agric. Agric. Res. Serv., 1700 SW 23rd Dr., Gainesville, FL 32608, USA

- Parker, C. D., Jr. and R. G. Luttrell (1998). Oviposition of tobacco budworm (Lepidoptera: Noctuidae) in mixed plantings of non-transgenic and transgenic cottons expressing delta-endotoxin protein of *Bacillus thuringiensis* (Berliner). *Southwestern Entomologist* 23(3): 247-257. Dep. Entomology Plant Pathology, Miss. Agricultural Forestry Experiment Station, Miss. State Univ., Box 9775, Mississippi State, MS 39762, USA
- Patel, B. H., D. M. Patel, et al. (1998). Bio-efficacy of various readymix synthetic insecticides against pest complex of 'H 6' cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 68(12): 780-781. {a} Bansilal Amrutlal College of Agriculture, Gujarat Agricultural University, Campus Anand, Anand, 388 110, India
- Payne, W. A., J. H. Williams, et al. (1998). Crop diversification in the Sahel through use of environmental changes near *faidherbia albida* (Del.) A. Chev. *Crop Science* 38(6): 1585-1591. {a} Columbia Basin Agric. Res. Cent., Oreg. State Univ., P.O. Box 370, Pendleton, OR 97801, USA
- Penna, J. C. V., L. M. Verhalen, et al. (1998). Screening cotton genotypes for seedling drought tolerance. *Genetics and Molecular Biology* 21(4): 545-549. {a} Dep. Agron., Univ. Federal Uberlandia, Caixa Postal 593, 38400-902 Uberlandia, MG, Brazil
- Percy, R. G. and E. L. Turcotte (1998). Registration of extra-long staple cotton germplasm, 89590 and 8810. *Crop Science* 38(5): 1409. {a} USDA-ARS, Maricopa Agric. Cent., 37860 W. Smith-Enke Rd., Maricopa, AZ 85239, USA
- Pettigrew, W. T. and R. B. Turley (1998). Variation in photosynthetic components among photosynthetically diverse cotton genotypes. *Photosynthesis Research* 56(1): 15-25. USDA-ARS, Cotton Physiol. Genet. Res. Unit, PO Box 345, Stoneville, MS 38776, USA
- Pettigrew, W. T. and K. C. Vaughn (1998). Physiological, structural, and immunological characterization of leaf and chloroplast development in cotton. *Protoplasma* 202(1-2): 23-37. {a} Southern Weed Sci. Res. Unit, ARS, U.S. Dep. Agriculture, P.O. Box 350, Stoneville, MS 38776, USA
- Ponte, J. J. D., F. J. Silveira, et al. (1998). Synopsis of the Brazilian literature of the Meloidogyne-cotton association. *Summa Phytopathologica* 24(2): 101-104. {a} UFC, Dep. Fitotecnia, Setor Fitopatol., C.P. 12168, 60356-001 Fortaleza, CE, Brazil
- Prasad, M. (1998). Nitrogen, phosphorus and sulphur requirement of cotton (*Gossypium hirsutum*) -gobhi sarson (*Brassica napus* subsp. *oleifera* var. *annua*) rotation. *Indian Journal of Agronomy*. Dec. 43(4): 612-615. {a} Division of Agronomy, Indian Agricultural Research Institute, New Delhi, 110 012, India
- Prasad, M. and R. Prasad (1998). Nutrient management in cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 43(1): 162-164. Div. Agron., Indian Agric. Res. Inst., New Delhi 110 012, India
- Prior, S. A., H. A. Torbert, et al. (1998). Effects of carbon dioxide enrichment on cotton nutrient dynamics. *Journal of Plant Nutrition* 21(7): 1407-1426. {a} USDA-ARS Natl. Soil Dynamics Lab., POB 3439, Auburn, AL 36831-3439, USA
- Prostko, E. P. and J. M. Chandler (1998). Devil's-claw (*Proboscidea louisianica*) and smellmelon (*Cucumis melo* var. *dudaim*) control in cotton (*Gossypium hirsutum*) with pyriithiobac. *Weed Technology* 12(1): 19-22. {a} Texas A and M Univ., Res. Exten. Cent., Route 2, Box 1, Stephenville, TX 76401, USA
- Ramachandra, R. A., K. R. Reddy, et al. (1998). Interactive effects of elevated carbon dioxide and growth temperature on photosynthesis in cotton leaves. *Plant Growth Regulation* 26(1): 33-40. {a} Sch. Life Sci., Pondicherry University, Pondicherry 605 014, India
- Reddy, K. R., R. R. Robana, et al. (1998). Interactions of CO₂ enrichment and temperature on cotton growth and leaf characteristics. *Environmental and Experimental Botany* 39(2): 117-129. {a} Dep. Plant Soil Sciences, Box 9555, Miss. State Univ., Mississippi State, MS 39762, USA

- Robbins, R. T., V. M. McNeely, et al. (1998). The lance nematode, *Hoplolaimus magnistylus*, on cotton in Arkansas. *Journal of Nematology* Dec. 30(4 Suppl): 590-591. {a} Department of Plant Pathology, University of Arkansas, Fayetteville, AR, 72701, USA
- Rochester, I. J., M. B. Peoples, et al. (1998). Faba beans and other legumes add nitrogen to irrigated cotton cropping systems. *Australian Journal of Experimental Agriculture* 38(3): 253-260. {a} CRC Sustainable Cotton Prod., CSIRO Cotton Res. Unit, P.O. Box 59, Narrabri, NSW 2390, Australia
- Rodriguez, K. R., W. S. Gazaway, et al. (1998). Host suitability of selected tropical legumes and other crops for the reniform nematode, *Rotylenchulus reniformis* Linford & Oliveira, 1940. *Nematropica* . Dec. 28(2): 195-203. {a} Department of Plant Pathology, Auburn University, Auburn, AL, USA
- Rosinski, J. A. and W. R. Atchley (1998). Molecular evolution of the Myb family of transcription factors: Evidence of polyphyletic origin. *Journal of Molecular Evolution* 46(1): 74-83. {a} Cent. Quantitative Genet., Dep. Genet., North Carolina State Univ., Box 7614, Raleigh, NC 27695, USA
- Roussopoulos, D., A. Liakatas, et al. (1998). Controlled-temperature effects on cotton growth and development. *Journal of Agricultural Science* 130(4): 451-462. {a} Agric. Univ. Athens, Dep. Land Reclam. Agric. Eng., 75 Iera Odos, GR-118 55 Athens, Greece
- Rout, D. and M. R. Satapathy (1998). Chemical weed control in rainfed cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 43(2): 348-350. Natl. Agric. Res. Project, Orissa Univ. Agric. Technol., Umerkote 764 073, India
- Ruan, Y. L. and P. S. Chourey (1998). A fiberless seed mutation in cotton is associated with lack of fiber cell initiation in ovule epidermis and alterations in sucrose synthase expression and carbon partitioning in developing seeds. *Plant Physiology Rockville* 118(2): 399-406. {a} Program Plant Molecular Cellular Biol., Dep. Plant Pathol., Univ. Fla., Gainesville, FL 32611-0680, USA
- Russo, V. M. and S. V. Karmarkar (1998). Water extraction of plant tissues for analysis by ion chromatography. *Communications in Soil Science and Plant Analysis* 29(3-4): 245-253. {a} United States Dep. Agriculture, Agricultural Res. Serv., South Central Agricultural Res. Lab., P.O. Box 159, Hwy. 3 West, Lane, OK 74555, USA
- Sachs, E. S., J. H. Benedict, et al. (1998). Expression and segregation of genes encoding CryIA insecticidal proteins in cotton. *Crop Science* 38(1): 1-11. {a} Texas A and M Univ. Res. and Ext. Center, Route 2, Box 589, Corpus Christi, TX 78406, USA
- Sadras, V. O. (1998). Herbivory tolerance of cotton expressing insecticidal proteins from *Bacillus thuringiensis*: Responses to damage caused by *Helicoverpa* spp. and to manual bud removal. *Field Crops Research* 56(3): 287-299. {a} CSIRO Plant Ind., Locked Bag 59, Narrabri, NSW 2390, Australia
- Sadras, V. O. and L. J. Wilson (1998). Recovery of cotton crops after early season damage by thrips (Thysanoptera). *Crop Science* 38(2): 399-409. {a} Univ. Nac. Mar del Plata, Fac. Ciencias Agrarias, CC 276 Balcarce, Argentina
- Sadras, V. O., L. J. Wilson, et al. (1998). Water deficit enhanced cotton resistance to spider mite herbivory. *Annals of Botany London* 81(2): 273-286. {a} Univ. Nacional de Mar del Plata, Fac. Ciencias Agrarias, CC276, Balcarce, Argentina
- Saranga, Y., A. Landa, et al. (1998). Near-infrared analysis of cotton leaves as a guide for nitrogen fertilization. *Agronomy Journal* 90(1): 16-21. {a} Dep. Field Crops Veg. Genet., Fac. Agric. Food Environ. Qual. Sci., Hebrew Univ. Jerusalem, P.O. Box 12, Rehovot 76100, Israel
- Saranga, Y., N. Sass, et al. (1998). Drought conditions induce mote formation in interspecific cotton hybrids. *Field Crops Research* 55(3): 225-234. {a} Hebrew Univ. Jerusalem, Fac. Agricultural, Food Environmental Quality Sci., Dep. Field Crops, Vegetables Genetics, P.O. Box 12, Rehovot 76100, Israel

- Seagull, R. W. (1998). Cytoskeletal stability affects cotton fiber initiation. *International Journal of Plant Sciences* 159(4): 590-598. {a} Dep. Biology, Hofstra Univ., Hempstead, NY 11549, USA
- Seifert, S. and C. E. Snipes (1998). Response of cotton (*Gossypium hirsutum*) to flame cultivation. *Weed Technology* 12(3): 470-473. {a} Miss. State Univ., Dep. Plant Soil Sci., P.O. Box 9555, Mississippi State, MS 39762, USA
- Shappley, Z. W., J. N. Jenkins, et al. (1998). An RFLP linkage map of Upland cotton, *Gossypium hirsutum* L. *Theoretical and Applied Genetics* 97(5-6): 756-761. {a} USDA, ARS Crop Science Res., Lab., Mississippi State, MS 39762, USA
- Sheidai, M., M. V. Tabor, et al. (1998). Cytogenetical studies in *Gossypium hirsutum* L. cultivars and their hybrids. *Cytologia Tokyo* 63(1): 41-48. {a} Biol. Dep. Shahid Beheshti Univ., Tehran, Iran
- Shimony, C., N. Sass, et al. (1998). Microscopic study of mature seed coats and aborted structures of interspecific cotton hybrids. *Botanica Acta* 111(1): 16-21. {a} Hebrew Univ. Jerusalem, Fac. Agric. Food Environ. Quality Sci., Dep. Field Crops Vegetables Genetics, P.O. Box 12, Rehovot 76100, Israel
- Sidhu, G. S., B. K. Kandpal, et al. (1998). Evaluation of soils of Jalgaon district of Maharashtra for cotton (*Gossypium hirsutum*) cultivation. *Indian Journal of Agricultural Sciences* 68(9): 607-610. {a} Regional Centre, Natl. Bur. Soil Survey Land Use Planning, IARI Campus, New Delhi 110 012, India
- Singh, J. and B. S. Sekhon (1998). Population build-up of cotton jassid, *Amrasca biguttula* (Ishida) on different varieties of cotton. *Journal of Insect Science*. [print] March 11(1): 53-55. {a} Department of Entomology, Punjab Agricultural University, Ludhiana, PU, 141 004, India
- Singh, K., B. S. Sandhu, et al. (1998). Anther culture response in cotton. *Annals of Biology Hissar* 14(1): 11-13. Dep. Plant Breeding, Punjab Agric. Univ., Ludhiana 141- 004, India
- Singh, M., R. P. Singh, et al. (1998). Inheritance and linkage relations of genes for yellow seedling in cotton (*Gossypium* spp.). *Indian Journal of Genetics and Plant Breeding* 58(2): 143-147. Div. Genet., Indian Agric. Res. Inst., New Delhi 110 012, India
- Singh, T. (1998). Residual effects of 2, 4-D applied to wheat on the succeeding crop of cotton (*Gossypium hirsutum* L.). *Annals of Biology Hissar* 14(1): 75-76. {a} PAU Regional Res. Station, Bathinda-151 001, India
- Singh, V. and R. Deo (1998). Productivity and economics of different cropping systems under various levels of irrigation. *Indian Journal of Agronomy*. Sept. 43(3): 419-425. {a} Agricultural Research Station, Agronomy Section, Rajasthan Agriculture University, Sriganganagar, 335 001, India
- Singh, V., S. K. Sharma, et al. (1998). Performance of different crop sequences under various irrigation levels. *Indian Journal of Agronomy* 43(1): 38-44. Agric. Res. Stn., Rajasthan Agric. Univ., Sriganganagar 335 001, India
- Smart, L. B., F. Vojdani, et al. (1998). Genes involved in osmoregulation during turgor-driven cell expansion of developing cotton fibers are differentially regulated. *Plant Physiology Rockville* 116(4): 1539-1549. {a} Dep. Agronomy Range Sci., Univ. California, Davis, CA 95616, USA
- Snodgrass, G. L. (1998). Distribution of the tarnished plant bug (Heteroptera: Miridae) within cotton plants. *Environmental Entomology* 27(5): 1089-1093. {a} South. Insect Manage. Res. Unit, USDA-ARS, P.O. Box 346, Stoneville, MS 38776, USA
- Solaiappan, U. and A. A. Dason (1998). Influence of sowing time and mulching practices on physiological parameters of cotton (*Gossypium hirsutum* L.) in rainfed vertisols. *Indian Journal of Agricultural Research*. Dec. 32(4): 243-248. {a} Agricultural Research Station, Tamilnadu Agricultural University, Kovilpatti, 626 701, India

- Solaippan, U. and A. A. Dason (1998). Influence of sowing time, intercropping and mulching on the growth and yield of rainfed cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy*. Dec. 43(4): 616-621. {a} Agricultural Research Station, Tamil Nadu Agricultural University, Kovilpatti, 626 701, India
- Song, F. M. and Z. Zheng (1998). The correlation between inhibition of ethylene production and trifluralin-induced resistance of cotton seedlings against *Fusarium oxysporum* f. sp. *vasinfectum*. *Acta Phytophysiological Sinica* 24(2): 111-118. Dep. Plant Prot., Zhejiang Agric. Univ., Hangzhou 310028, China
- Soomro, A. R., A. M. Memon, et al. (1998). Effect of okra leaf shape on boll rot, earliness and yield of upland cotton *Gossypium hirsutum* L. *Pakistan Journal of Scientific and Industrial Research* 41(1): 36-38. {a} Cotton Research Institute, Sakrand 67210, Pakistan
- Soomro, Z. A., M. S. Kalwar, et al. (1998). Screening elite hybrids of cotton *Gossypium hirsutum* L. *Pakistan Journal of Scientific and Industrial Research*. Oct. 41(5): 260-263. {a} Department of Plant Breeding and Genetics, Sindh Agriculture University, Tando Jam, Pakistan
- Steger, A. J., J. C. Silvertooth, et al. (1998). Upland cotton growth and yield response to timing the initial postplant irrigation. *Agronomy Journal* 90(4): 455-461. {a} Alzheimer Lab., Dep. Agron., Univ. Arkansas, Fayetteville, AR 72704, USA
- Stewart, A. M. and K. L. Edmisten (1998). Cotton response to placement and rate of starter fertilizer. *Journal of Plant Nutrition* 21(5): 967-973. *Crop Sci. Dep., North Carolina State Univ., Raleigh, NC 27695-7620, USA*
- Taskin, T., R. Demir, et al. (1998). The formation of callus cultures from unpollinated pistils in cotton plant (*Gossypium hirsutum* L.). *Biochemical Archives* 14(2): 105-109. {a} Univ. Dicle, Fac. Educ., Dep. Biol., Diyarbakir, Turkey
- Thornton, F. C., J. D. Joslin, et al. (1998). Environmental effects of growing woody crops on agricultural land: First year effects on erosion, and water quality. *Biomass and Bioenergy* 15(1): 57-69. {a} TVA, Atmospheric Sci., Mus. Shoals, AL 35662, USA
- Triplett, B. A. (1998). Stage-specific inhibition of cotton fiber development by adding alpha-amanitin to ovule cultures. *In Vitro Cellular and Developmental Biology Plant* 34(1): 27-33. {a} USDA, ARS, Southern Regional Res. Cent., P.O. Box 19687, New Orleans, LA 70179, USA
- Van, E. G. A., D. T. Bowman, et al. (1998). Changes in the genetic diversity of cotton in the USA from 1970 to 1995. *Crop Science* 38(1): 33-37. {a} Dep. Crop Sci., North Carolina State Univ., Raleigh, NC 27695-8604, USA
- Van, t. H. J. (1998). Production of micronucleoli and onset of cotton fiber growth. *Planta Berlin* 205(4): 561-566. {a} Biol. Dep., Brookhaven Natl. Lab., Upton, NY 11973, USA
- Van, t. H. J. and S. Saha (1998). Growth and mitotic potential of multicelled fibers of cotton (Malvaceae). *American Journal of Botany* 85(1): 25-29. {a} Biol. Dep., Brookhaven Natl. Lab., Upton, NY 11973, USA
- Veenstra, K. H. and D. N. Byrne (1998). The effects of physiological factors and host plant experience on the ovipositional activity of the sweet potato whitefly, *Bemisia tabaci*. *Entomologia Experimentalis et Applicata* 89(1): 15-23. Dep. Entomology, Univ. Ariz., Tucson, AZ 85721, USA
- Venuguoplan, M. V. and P. T. Tarhalkar (1998). Efficiency of foliar fertilization in rainfed upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 68(6): 313-314. {a} Central Inst. Cotton Res., Nagpur, Maharashtra 440 001, India
- Villavaso, E. J., W. L. McGovern, et al. (1998). Components of competitiveness in sterile male boll weevils (Coleoptera: Curculionidae). *Journal of Economic Entomology* 91(3): 631-636. USDA-ARS, P.O. Box 5367, Mississippi State, MS 39762, USA

- Vroh, B. I., J. P. Baudoin, et al. (1998). Cytogenetics of the 'glandless-seed and glanded-plant' trait from *Gossypium sturtianum* Willis introgressed into upland cotton (*Gossypium hirsutum* L.). *Plant Breeding* 117(3): 235-241. Unit. Phytotechnie Regions Intertropicales, Fac. Univ. Sci. Agron., 2 Passage des Deportes, B-5030 Gembloux, Belgium
- Walker, N. R., T. L. Kirkpatrick, et al. (1998). Interaction between *Meloidogyne incognita* and *Thielaviopsis basicola* on cotton (*Gossypium hirsutum*). *Journal of Nematology*. Dec. 30(4): 415-422. {a} Department of Plant Pathology, University of Arkansas, Fayetteville, AR, 72701, USA
- Wang, Z. Z., S. F. Zhang, et al. (1998). Isolation, culture and plant regeneration of protoplasts from an embryogenic callus tissue of *Gossypium hirsutum*. *Acta Botanica Sinica* 40(3): 234-240. {a} Dep. Biol., Xinyang Normal COLL., Xinyang, Henan 464000, China
- Warren, G. F. (1998). Spectacular increases in crop yields in the United States in the Twentieth Century. *Weed Technology* 12(4): 752-760. {a} Purdue Univ., 1130 Cherry Lane, West Lafayette, IN 47906, USA
- Wiedenfeld, R. P. (1998). Previous-crop effects on sugarcane responses to nitrogen fertilization. *Agronomy Journal* 90(2): 161-165. {a} Texas Agric. Exp. Stn., Texas A and M Univ. Res. Ext. Cent., 2415 E. Hwy. 83, Weslaco, TX 78596, USA
- Wu, F., L. Wu, et al. (1998). Chlorophyll meter to predict nitrogen sidedress requirements for short-season cotton (*Gossypium hirsutum* L.). *Field Crops Research* 56(3): 309-314. {a} Dep. Agronomy, Zhejiang Agric. Univ., Hangzhou 310029, China
- Yee, W. L., N. C. Toscano, et al. (1998). Effects of insecticide applications on *Bemisia argentifolii* (Homoptera: Aleyrodidae) densities and honeydew production. *Environmental Entomology* 27(1): 22-32. {a} Amway Corp., Nutrilite Div., 19600 6th St., Lakeview, CA 92567, USA
- Yokoyama, V. Y., E. B. Jang, et al. (1998). Laboratory infestation of cotton bolls by Mediterranean fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 91(1): 293-296. {a} Hortic. Crops Res. Lab., USDA-ARS, 2021 S. Peach Ave., Fresno, CA 93727, USA
- Zak, J. C., B. McMichael, et al. (1998). Arbuscular-mycorrhizal colonization dynamics of cotton (*Gossypium hirsutum* L.) growing under several production systems on the Southern High Plains, Texas. *Agriculture Ecosystems and Environment* 68(3): 245-254. {a} Dep. Biol. Sci., Texas Tech Univ., Lubbock, TX 79409-3131, USA
- Zeringue, H. J., Jr. and B. Y. Shih (1998). Extraction and separation of the bright-greenish-yellow fluorescent material from aflatoxigenic *Aspergillus* spp. infected cotton lint by HPLC-UV/FL. *Journal of Agricultural and Food Chemistry* 46(3): 1071-1075. {a} ARS, USDA, SRRC, PO Box 19687, New Orleans, LA 70179-0687, USA
- Zhang, H. L., A. Nagatsu, et al. (1998). Sesquiterpene glycosides from cotton oil cake. *Phytochemistry Oxford* 48(4): 665-668. {a} Fac. Pharm. Sci., Nagoya City Univ., Tanabe-dori, Mizuho-ku, Nagoya 467, Japan
- Zhao, D. and D. Oosterhuis (1998). Cotton responses to shade at different growth stages: Nonstructural carbohydrate composition. *Crop Science* 38(5): 1196-1203. {a} Dep. Agron., 276 Altheimer Dr., Univ. Arkansas, Fayetteville, AR 72704, USA
- Zhao, D. and D. Oosterhuis (1998). Physiologic and yield responses of shaded cotton to the plant growth regulator PGR-IV. *Journal of Plant Growth Regulation* 17(1): 47-52. {a} Altheimer Lab., Dep. Agronomy, Univ. Arkansas, Fayetteville, AR 72704, USA
- Zhao, D. and D. M. Oosterhuis (1998). Influence of shade on mineral nutrient status of field-grown cotton. *Journal of Plant Nutrition* 21(8): 1681-1695. Dep. Agronomy, Univ. Arkansas, 276 Altheimer Lab., Fayetteville, AR 72704, USA

Zhu, X. F., X. D. Wang, et al. (1998). Assessment of cytoplasmic effects of cytoplasmic male-sterile lines in upland cotton. *Plant Breeding* 117(6): 549-552. {a} Key Lab. Crop Germplasm Breed., Minist. Agric., Dep. Agron., Nanjing Agric. Univ., Nanjing 210095, China

Agrawal, D. C., A. K. Banerjee, et al. (1997). In vitro induction of multiple shoots and plant regeneration in cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 16(9): 647-652. Plant Tissue Culture Div., Natl. Chem. Lab., Pune 411 008, India

Ahmad, R. T., I. A. Khan, et al. (1997). Diallel analysis for seed-cotton yield and its contributing traits in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 67(12): 583-585. Dep. Plant Breed. Genet., Univ. Agric., Faisalabad, Pakistan

Akhtar, L. H., M. U. D. Khan, et al. (1997). Effects of thiamine (vitamin B1) on growth of callus cultures of acala SJ2 (*Gossypium hirsutum* L.). *Sarhad Journal of Agriculture* 13(6): 571-574. {a} Cotton Res. Station, Multan, Pakistan

Akhtar, L. H., M. U. D. Khan, et al. (1997). Effect of nicotinic acid and pyridoxine-HCL on growth of callus cultures of *Gossypium hirsutum* L. *Sarhad Journal of Agriculture* 13(6): 565-570. {a} Cotton Res. Station, Multan, Pakistan

Akhtar, L. H. and M. A. Mirza (1997). Effect of initial callus size/quantity, agar, phytagel and thiamine on further growth of leaf and petiole-derived callus cultures of *Gossypium hirsutum* L. *Pakistan Journal of Scientific and Industrial Research* 40(1-4): 31-33. {a} Cotton Res. Station, Multan, Pakistan

Allen, R. L., C. E. Snipes, et al. (1997). Fruiting response of cotton (*Gossypium hirsutum*) to pyriithobac. *Weed Technology* 11(1): 59-63. {a} Mississippi Agric. For. Exp. Station, Delta Branch, Stoneville, MS 38776, USA

Allender, W. J. (1997). Effect of trifluoperazine and verapamil on herbicide stimulated growth of cotton. *Journal of Plant Nutrition* 20(1): 69-80. Biol. Chem. Res. Inst., NSW Agric., PMB 10, Rydalmere 2116, Australia

Allender, W. J., G. C. Cresswell, et al. (1997). Effect of lithium and lanthanum on herbicide induced hormesis in hydroponically-grown cotton and corn. *Journal of Plant Nutrition* 20(1): 81-95. {a} Biol. Chem. Res. Inst., NSW Agric., PMB 10, Rydalmere, NSW 2116, Australia

Aslam, M., K. Nielson, et al. (1997). Nitrate uptake, efflux, and in vivo reduction by Pima and Acala cotton cultivars. *Crop Science* 37(6): 1795-1801. {a} Dep. Agron. Range Sci., Univ. Calif., Davis, CA 55616-8515, USA

Baird, R. E. and D. E. Carling (1997). First report of *Rhizoctonia solani* AG-7 in Georgia. *Plant Disease* 81(7): 832. {a} Plant Pathol. Dep., Univ. Ga., RDC, Tifton, GA 31793, USA

Baloch, M. J., H. Bhutto, et al. (1997). Combining ability estimates of highly adapted tester lines crossed with pollinator inbreds of cotton, *Gossypium hirsutum* L. *Pakistan Journal of Scientific and Industrial Research* 40(5-12): 95-98. Cotton Res. Inst., Sakrand, Sindh, Pakistan

Batts, R. B. and A. C. York (1997). Weed management in no-till cotton (*Gossypium hirsutum*) with thiazopyr. *Weed Technology* 11(3): 580-585. Dep. Crop Sci., North Carolina State Univ., Raleigh, NC 27695-7620, USA

Bauer, P. J., E. J. Sadler, et al. (1997). Intermittent shade effect on gas exchange of cotton leaves in the humid southeastern USA. *Agronomy Journal* 89(2): 163-166. {a} USDA-ARS Coastal Plains Soil Water and Plant Res. Ctr., 2611 W. Lucas St., Florence, SC 29501-1241, USA

Bejarano, A. J., L. M. A. Blanco, et al. (1997). The influence of verticillium wilt epidemics on cotton yield in southern Spain. *Plant Pathology Oxford* 46(2): 168-178. {a} Dep. Proteccion Veg., CIFA Cordoba, Apdo. 4240, 14080 Cordoba, Spain

- Bi, J. L., J. B. Murphy, et al. (1997). Antinutritive and oxidative components as mechanisms of induced resistance in cotton to *Helicoverpa zea*. *Journal of Chemical Ecology* 23(1): 97-117. {a} Dep. Entomol., Univ. Arkansas, Fayetteville, AR 72701, USA
- Boman, R. K., W. R. Raun, et al. (1997). Long-term nitrogen fertilization in short-season cotton: Interpretation of agronomic characteristics using stability analysis. *Journal of Production Agriculture* 10(4): 580-585. {a} Dep. Agron., Okla. State Univ., Stillwater, OK 74078, USA
- Bondada, B. R., D. M. Oosterhuis, et al. (1997). Cotton leaf age, epicuticular wax, and nitrogen-15 absorption. *Crop Science* 37(3): 807-811. {a} Dep. Plant Soil Sci., Univ. Tennessee, PO Box 1071, Knoxville, TN 37901, USA
- Bourland, F. M., R. E. McGowen, Jr., et al. (1997). Registration of Arkot 8303 germplasm line of cotton. *Crop Science* 37(4): 1397. {a} NEREC, Univ. Arkansas, P.O. Box 48, Keiser, AR 72351, USA
- Bourland, F. M., R. E. McGowen, Jr., et al. (1997). Registration of Arkot 8102, Arkot 8506, and Arkot 8514 germplasm lines of cotton. *Crop Science* 37(4): 1396. {a} NEREC, Univ. Arkansas, P.O. Box 48, Keiser, AR 72351, USA
- Bourland, F. M., R. E. McGowen, Jr., et al. (1997). Registration of Arkot 8110 germplasm line of cotton. *Crop Science* 37(4): 1395-1396. {a} NEREC, Univ. Arkansas, P.O. Box 48, Keiser, AR 72351, USA
- Bowman, D. T. and J. C. McCarty, Jr. (1997). Thrips (Thysanoptera: Thripidae) tolerance in cotton: Sources and heritability. *Journal of Entomological Science* 32(4): 460-471. Dep. Crop Science, Box 8604, N.C. State Univ., Raleigh, NC 27695-8604, USA
- Bradow, J. M., P. J. Bauer, et al. (1997). Quantitation of cotton fiber-quality variations arising from boll and plant growth environments. *European Journal of Agronomy* 6(3-4): 191-204. {a} USDA, ARS, Southern Reg. Res. Cent., P.O. Box 19687, New Orleans, LA 70179, USA
- Bunkelmann, J. R. and R. N. Trelease (1997). Expression of glyoxysomal ascorbate peroxidase in cotton seedlings during postgerminative growth. *Plant Science Shannon* 122(2): 209-216. {a} Box 871601, Molecular Cellular Biol. Program, Dep. Botany, Arizona State Univ., Tempe, AZ 85287-1601, USA
- Busscher, W. J., P. J. Bauer, et al. (1997). Correction of cone index for soil water content differences in a coastal plain soil. *Soil and Tillage Research* 43(3-4): 205-217. {a} Coastal Plains Soil, Water Plant Res. Cent., USDA-ARS, Florence, SC, USA
- Butter, N. S., S. Singh, et al. (1997). Studies on growth and development of *Heliothis armigera* Hubn. on cotton genotypes. *Journal of Entomological Research New Delhi* 21(1): 51-58. {a} Punjab Agric. Univ., Ludhiana-141004, Punjab, India
- Calhoun, D. S. (1997). Inheritance of high glanding, an insect resistance trait in cotton. *Crop Science* 37(4): 1181-1186. Delta Res. Extension Cent., P.O. Box 197, Stoneville, MS 38776, USA
- Calhoun, D. S., J. E. Jones, et al. (1997). Registration of 'H1244' cotton. *Crop Science* 37(3): 1014-1015. {a} Delta Res. Ext. Cent., Stoneville, MS 38776, USA
- Calhoun, D. S., J. E. Jones, et al. (1997). Registration of 'H1220' cotton. *Crop Science* 37(3): 1013-1014. {a} Delta Res. Ext. Cent., Stoneville, MS 38776, USA
- Calhoun, D. S., J. E. Jones, et al. (1997). Registration of 'H1215' cotton. *Crop Science* 37(3): 1013. {a} Delta Res. Ext. Cent., Stoneville, MS 38776, USA
- Candole, B. L. and C. S. Rothrock (1997). Characterization of the suppressiveness of hairy vetch-amended soils to *Thielaviopsis basicola*. *Phytopathology* 87(2): 197-202. {a} Dep. Plant Pathol., Univ. Arkansas, Fayetteville, AR 72701, USA

- Cao, H. M. H. M. R., M. Y. Wang, et al. (1997). Effect of seedling transplantation on the ^{14}C - CO_2 assimilation and the apportionment of ^{14}C assimilations within the cotton plant interplanted in standing wheat. *Acta Botanica Sinica* 39(3): 241-246. {a} Dezhou Tech. Dep., Shandong Agric. Univ., Dezhou 253000, China
- Chakrabarty, P. K., Y. P. Duan, et al. (1997). Cloning and characterization of a member of the *Xanthomonas* avr/pth gene family that evades all commercially utilized cotton R genes in the United States. *Phytopathology* 87(11): 1160-1167. Plant Pathol. Dep., Univ. Florida, Gainesville, FL 32611, USA
- Chand, M., M. Prasad, et al. (1997). Effect of fertilizers on growth, yield and quality of cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 42(3): 528-530. {a} Plant Quarantine Fumigation Stn., 6 Clive Battery, Madras 600 001, India
- Chapman, K. D., R. S. McAndrew, et al. (1997). Cottonseed calnexin: Identification and isolation of a membrane-bound molecular chaperone. *Plant Physiology and Biochemistry Paris* 35(6): 483-490. {a} Div. Biochem. Mol. Biol., Dep. Biol. Sci., Univ. North Texas, Denton, TX 76203-0218, USA
- Chapman, K. D. and A. Sriparameswaran (1997). Intracellular localization of N-acylphosphatidylethanolamine synthesis in cotyledons of cotton (*Gossypium hirsutum* L.) seedlings. *Plant and Cell Physiology* 38(12): 1359-1367. {a} Dep. Biological Sciences, Div. Biochem. Molecular Biol., Univ. North Texas, Denton, TX 76203-5220, USA
- Charles, G. W. (1997). Herbicide strategies for reducing nutgrass (*Cyperus rotundus* L.) density in cotton (*Gossypium hirsutum* L.). *Australian Journal of Experimental Agriculture* 37(2): 231-241. NSW Agric., Aust. Cotton Res. Inst., Narrabri, NSW 2390, Australia
- Chen, J. G., X. M. Du, et al. (1997). Levels of cytokinins in the ovules of cotton mutants with altered fiber development. *Journal of Plant Growth Regulation* 16(3): 181-185. {a} Key Lab. Crop Growth Regulation, Ministry Agric., Dep. Agronomy, Agric. Univ., Nanjing 210095, China
- Chitarra, L. G., J. D. C. Machado, et al. (1997). Performance of cottonseeds (*Gossypium hirsutum* L.) delinted with sulphuric acid. *Ciencia e Agrotecnologia* 21(4): 425-435. {a} Dep. Fitosanidade, UFLA, Caixa Postal 37, CEP. 37.200-000, Lavras-MG, Brazil
- Chung, R. P. T., G. M. Neumann, et al. (1997). Purification and characterization of basic proteins with in vitro antifungal activity from seeds of cotton, *Gossypium hirsutum*. *Plant Science Shannon* 127(1): 1-16. {a} Sch. Biochem., La Trobe Univ., Bandoora, Victoria 3083, Australia
- Colyer, P. D., T. L. Kirkpatrick, et al. (1997). Influence of nematicide application on the severity of the root-knot nematode-fusarium wilt disease complex in cotton. *Plant Disease* 81(1): 66-70. {a} Louisiana Agricultural Experiment Station, Red River Res. Station, P.O. Box 8550, Bossier City, LA 71113-8550, USA
- Cook, C. G., L. N. Namken, et al. (1997). Registration of N220-1-91, N222-1-91, N320-2-91, and N419-91 nematode-resistant cotton germplasm lines. *Crop Science* 37(3): 1028-1029. {a} USDA-ARS Conserv. Prod. Res. Unit, 2413 E. Hwy. 83, Weslaco, TX 78596, USA
- Cook, C. G., L. N. Namken, et al. (1997). Registration of C221-91, C224-91, C300-91, and C306-91 cotton germplasm lines. *Crop Science* 37(3): 1029. {a} USDA-ARS Conserv. Prod. Res. Unit, 2413 E. Hwy. 83, Weslaco, TX 78596, USA
- Cook, C. G., A. F. Robinson, et al. (1997). Tolerance of *Rotylenchulus reniformis* and resistance to *Meloidogyne incognita* race 3 in high-yielding breeding lines of upland cotton. *Journal of Nematology* 29(3): 322-328. {a} USDA, ARS, 2413 E. Hwy 83, Weslaco, TX 77845, USA
- Cortesero, A. M., C. M. D. Moraes, et al. (1997). Comparisons and contrasts in host-foraging strategies of two larval parasitoids with different degrees of host specificity. *Journal of Chemical Ecology* 23(6): 1589-1606. {a} Insect Biol. Population Manage. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Tifton, GA 31793, USA

- Coskuncelebi, K., R. T. Aliyev, et al. (1997). Effect of gibberellic acid on the alterations caused by drought stress in the genome of cotton (*Gossypium hirsutum* L.) seedlings. *Turkish Journal of Biology* 21(2): 175-179. Dep. Biol., Fac. Sci and Arts, Karadeniz Technical Univ., Trabzon, Turkey
- Coyle, G. G. and C. W. Smith (1997). Combining ability for within-boll yield components in cotton, *Gossypium hirsutum* L. *Crop Science* 37(4): 1118-1122. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA
- Culpepper, A. S. and A. C. York (1997). Weed management in no-tillage bromoxynil-tolerant cotton (*Gossypium hirsutum*). *Weed Technology* 11(2): 335-345. Crop Sci. Dep., North Carolina State Univ., Raleigh, NC 27695-7620, USA
- Davis, R. M., J. J. Nunez, et al. (1997). Benefits of cotton seed treatments for the control of seedling diseases in relation to inoculum densities of *Pythium* species and *Rhizoctonia solani*. *Plant Disease* 81(7): 766-768. {a} Dep. Plant Pathol., Univ. Calif., Davis, CA 95616, USA
- Devay, J. E., A. P. Gutierrez, et al. (1997). Inoculum densities of *Fusarium oxysporum* f. sp. *vasinfectum* and *Meloidogyne incognita* in relation to the development of fusarium wilt and the phenology of cotton plants (*Gossypium hirsutum*). *Phytopathology* 87(3): 341-346. {a} Dep. Plant Pathol., Univ. Calif., Davis, CA 95616, USA
- Devay, J. E., B. L. Weir, et al. (1997). Effects of *Verticillium dahliae* infection of cotton plants (*Gossypium hirsutum*) on potassium levels in leaf petioles. *Plant Disease* 81(9): 1089-1092. {a} Dep. Plant Pathol., Univ. Calif., Davis, CA 95616, USA
- Dubery, I. A. and V. Slater (1997). Induced defence responses in cotton leaf disks by elicitors from *Verticillium dahliae*. *Phytochemistry Oxford* 44(8): 1429-1434. {a} Dep. Biochem., Rand Afrikaans Univ., P.O. Box 524, Auckland Park, South Africa
- El, Z. K. M. and P. M. Thaxton (1997). Registration of seven multi-adversity resistant (MAR-5) germplasm lines of upland cotton. *Crop Science* 37(4): 1394-1395. {a} Dep. Soil Crop Sci., Texas Agric. Exp. Stn., Texas A and M Univ. System, Coll. Station, TX 77843-2474, USA
- Ellington, J., M. Southward, et al. (1997). Association among cotton arthropods. *Environmental Entomology* 26(5): 1004-1008. {a} Entomol. Plant Pathol Weed Sci. Dep., N.M. State Univ., Las Cruces, NM 88003, USA
- Ferguson, D. L., R. B. Turley, et al. (1997). Identification of a delta-TIP cDNA clone and determination of related A and D genome subfamilies in *Gossypium* species. *Plant Molecular Biology* 34(1): 111-118. {a} Cotton Physiol. Genetics, USDA-ARS, P.O. Box 345, Stoneville, MS 38776, USA
- Freitas, S. D. S. and M. A. Pizzinatto (1997). Action of rhizobacteria on the *Colletotrichum gossypii* incidence and growth promotion in cotton seedlings (*Gossypium hirsutum*). *Summa Phytopathologica* 23(1): 36-41. {a} Inst. Agronomico, Secao de Microbiol. do Solo, Caixa Postal 28, 13001-970 Campinas, SP, Brazil
- Garber, R. K. and P. J. Cotty (1997). Formation of sclerotia and aflatoxins in developing cotton bolls infected by the S strain of *Aspergillus flavus* and potential for biocontrol with an atoxigenic strain. *Phytopathology* 87(9): 940-945. {a} Southern Regional Res. Cent., Agric. Res. Serv., U.S. Dep. Agric., P.O. Box 19687, New Orleans, LA 70179, USA
- Gerling, D., V. Kravchenko, et al. (1997). Dynamics of common green lacewing (Neuroptera: Chrysopidae) in Israeli cotton fields in relation to whitefly (Homoptera: Aleyrodidae) populations. *Environmental Entomology* 26(4): 815-827. Dep. Zool., George S. Wise Fac. Life Sci., Tel Aviv Univ., Ramat Aviv 69978, Israel
- Giri, A. N. and R. H. Bhosle (1997). Economics of integrated weed management and nitrogen fertilization in cotton (*Gossypium hirsutum*) under rainfed conditions. *Indian Journal of Agronomy* 42(4): 705-708. {a} Cotton Res. Scheme, MAU, Parbhani, Maharashtra 431 402, India

- Gonzalez, B. M. E., J. M. F. C. Carvalho, et al. (1997). Cotton (*Gossypium hirsutum* L.) somatic embryogenesis: A comparative study between two cultivars. *Phytomorphology* 47(4): 375-382. {a} Depto. Biol. Veg., Escuela Tecnica Superior Ingenieros Agron., Ciudad Univ., Universidad Politecnica de Madrid, 28040 Madrid, Spain
- Goulart, A. M. C., M. M. Inomoto, et al. (1997). Host suitability of eight cotton cultivars to *Pratylenchus brachyurus*. *Nematologia Brasileira* 21(2): 111-118. ESALQ/USP, Dep. Zool., 13418-900 Piracicaba, SP, Brazil
- Grantz, D. A., X. J. Zhang, et al. (1997). Ozone deposition to a cotton (*Gossypium hirsutum* L.) field: Stomatal and surface wetness effects during the California Ozone Deposition Experiment. *Agricultural and Forest Meteorology* 85(1-2): 19-31. {a} Dep. Bot. Plant Sci., Statewide Air Pollution Res. Cent., Univ. Calif. Riverside, Kearney Agric. Cent., 9240 S. Riverbend Ave., Parlier, CA 93648, USA
- Gutierrez, J. C. (1997). Registration of 'Maria del Mar' cotton. *Crop Science* 37(4): 1389. Cotton Dep., Centro Investigacion Desarrollo Agrario Las Torres-Tomejil, Alcala del Rio, Seville, Spain
- Gutierrez, J. C. (1997). Registration of 'Victoria' cotton. *Crop Science* 37(4): 1389. Cotton Dep., Cent. Investigacion Desarrollo Agrario Las Torres-Tomejil, Alcala Rio, Seville, Spain
- Hallmann, J., J. W. Kloepper, et al. (1997). Application of the scholander pressure bomb to studies on endophytic bacteria of plants. *Canadian Journal of Microbiology* 43(5): 411-416. {a} Dep. Plant Pathol., Biological Control Inst., Alabama Agricultural Experiment Stn., Auburn, AL 36849-5409, USA
- Harrison, B. D., Y. L. Liu, et al. (1997). Detection and relationships of cotton leaf curl virus and allied whitefly-transmitted geminiviruses occurring in Pakistan. *Annals of Applied Biology* 130(1): 61-75. {a} Dep. Biological Sciences, Univ. Dundee, Dundee DD1 4HN, UK
- Hedin, P. A., J. C. McCarty, Jr., et al. (1997). Effects of foliar applications of carbohydrates on the yield of cotton (*Gossypium hirsutum*) Lint. *Journal of Agricultural and Food Chemistry* 45(7): 2763-2767. {a} Crop Sci. Res. Lab., USDA-ARS, Mississippi State, Mississippi 39762, MS, USA
- Hedin, P. A., J. C. McCarty, Jr., et al. (1997). Caloric analyses of the distribution of energy in ripened cotton (*Gossypium hirsutum* L.). *Journal of Agricultural and Food Chemistry* 45(8): 3258-3261. {a} Crop Sci. Res. Lab., Agricultural Res. Service, U.S. Dep. Agriculture, P.O. Box 5367, Mississippi State, MS 39762, USA
- Heitholt, J. J. (1997). Floral bud removal from specific fruiting positions in cotton: Yield and fiber quality. *Crop Science* 37(3): 826-832. Cotton Physiol. Genet., USDA-ARS, PO Box 345, Stoneville, MS 38776, USA
- Howard, D. D., C. O. Gwathmey, et al. (1997). Potassium fertilization of cotton on two high testing soils under two tillage systems. *Journal of Plant Nutrition* 20(12): 1645-1656. Univ. Tenn., Plant Soil Sci. Dep., 605 Airways Blvd., Jackson, TN 38301, USA
- Hulugalle, N. R. and P. Entwistle (1997). Soil properties, nutrient uptake and crop growth in an irrigated Vertisol after nine years of minimum tillage. *Soil and Tillage Research* 42(1-2): 15-32. {a} Co-operative Res. Cent. Sustainable Cotton Production, Narrabri, NSW, Australia
- Hunt, P. G., P. J. Bauer, et al. (1997). Crop production in a wheat-cotton doublecrop rotation with conservation tillage. *Journal of Production Agriculture* 10(3): 462-465. {a} USDA-ARS Coastal Plains Soil Water Plant Res. Cent., 2611 W. Lucas St., Florence, SC 29501-1241, USA
- Hutchinson, R. L. and D. D. Howard (1997). Response of no-tillage and conventional-tillage cotton to starter fertilization on loess soils. *Journal of Plant Nutrition* 20(7-8): 975-986. {a} Louisiana State Univ. Agricultural Cent., Northeast Res. Station, P.O. Box 438, St. Joseph, LA 71366, USA

- Iqbal, M. J., N. Aziz, et al. (1997). Genetic diversity evaluation of some elite cotton varieties by RAPD analysis. *Theoretical and Applied Genetics* 94(1): 139-144. {a} Natl. Inst. Biotechnol. Genet. Eng., Jhang Road, P.O. Box 577, Faisalabad, Pakistan
- Jain, S. (1997). Studies on combining ability in intra and interspecific crosses of cotton. *Crop Research Hisar* 14(1): 91-95. J. N. K. V. V. Agricultural Res. Station, Badnawar 454 660, India
- Jenkins, J. N., J. C. McCarty, Jr., et al. (1997). Resistance of cotton with delta-endotoxin genes from *Bacillus thuringiensis* var. *kurstaki* on selected Lepidopteran insects. *Agronomy Journal* 89(5): 768-780. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA
- Jeyabalan, D. and K. Murugan (1997). Effect of neem limonoids on feeding and reproduction of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). *Entomon* 22(1): 15-20. {a} Div. Entomology, Dep. Zoology, Bharathiar Univ., Coimbatore 641 046, India
- Ji, Y., D. A. Raska, et al. (1997). Use of meiotic FISH for identification of a new monosome in *Gossypium hirsutum* L. *Genome* 40(1): 34-40. {a} Dep. Soil Crop Sci., Texas A M Univ., College Station, TX 77843, USA
- Johnson, W. C., III, R. D. Wauchope, et al. (1997). Response of peanut to low rates of MSMA. *Weed Science* 45(3): 430-433. {a} USDA-ARS, Coastal Plain Exp. Stn., Tifton, GA 31793, USA
- Jones, M. A. and R. Wells (1997). Dry matter allocation and fruiting patterns of cotton grown at two divergent plant populations. *Crop Science* 37(3): 797-802. {a} Dep. Crop Sci., N.C. State Univ., Raleigh, NC 27695-7620, USA
- Joslin, J. D. and S. H. Schoenholtz (1997). Measuring the environmental effects of converting cropland to short-rotation woody crops: A research approach. *Biomass and Bioenergy* 13(4-5): 301-311. {a} Tenn. Valley Authority, Atmospheric Sci., Norris, TN, USA
- Kalburtji, K. L. and A. Gagianas (1997). Effects of sugar beet as a preceding crop on cotton. *Journal of Agronomy and Crop Science* 178(1): 59-63. {a} Lab. Ecol. and Environ. Prot., Fac. Agric., Aristotelian Univ. Thessaloniki, 540 06 Thessaloniki, Greece
- Keller, G., L. Spatola, et al. (1997). Transgenic cotton resistant to herbicide bialaphos. *Transgenic Research* 6(6): 385-392. {a} Geniva, 585 Science Dr., Madison, WI 57311, USA
- Kerby, T. A., R. E. Plant, et al. (1997). Height-to-node ratio as an index of early season cotton growth. *Journal of Production Agriculture* 10(1): 80-83. {a} Delta Pine Land Co., Scott, MS 38772, USA
- Khalid, S., M. H. Soomro, et al. (1997). Occurrence of cotton leaf curl virus (CLCuV) in Sindh. *Pakistan Journal of Botany* 29(1): 173-174. {a} Crop Diseases Res. Inst., NARC, Park Rd., Islamabad-45500, Pakistan
- Khan, A. A., S. A. Rao, et al. (1997). Estimation of genetic components of variation through diallel analysis of some cotton fibre and seed properties. *Journal of Genetics and Breeding* 51(2): 143-147. {a} Sch. Biol. Sci., Univ. Liverpool, PO Box 147, Liverpool L69 3BX, UK
- Khorsandi, F., M. Y. L. Boone, et al. (1997). Validation of the soil temperature subroutine HEAT in the cotton simulation model GOSSYM. *Agronomy Journal* 89(3): 415-420. {a} Dep. Plant Soil Sci., Miss. State Univ., Mississippi State, MS 39762, USA
- Kjaersgarde, I. V. H., H. M. Jespersen, et al. (1997). Sequence and RT-PCR expression analysis of two peroxidases from *Arabidopsis thaliana* belonging to a novel evolutionary branch of plant peroxidases. *Plant Molecular Biology* 33(4): 699-708. {a} Dep. Protein Chem., Inst. Molecular Biol., Univ. Copenhagen, Oster Farimagsgade 2A, DK 1353 Copenhagen K, Denmark

- Kowsalya, R. and T. S. Raveendran (1997). Comparative influence of Harknessii and Hirsutum plasmons on agronomic traits in cotton hybrids. *Crop Improvement* 24(2): 37-40. Sch. Genet., Tamil Nadu Agric. Univ., Coimbatore-641 003, India
- Laisk, A., V. Oja, et al. (1997). Quantum yields and rate constants of photochemical and nonphotochemical excitation quenching. *Plant Physiology Rockville* 115(2): 803-815. {a} Tartu Ulikooli Molekulaar- ja Rakubioloogia Instituut, Riia tn. 181, Tartu EE2400, Estonia
- Larik, A. S., S. R. Ansari, et al. (1997). Heritability analysis of yield and quality components in *Gossypium hirsutum* L. *Pakistan Journal of Botany* 29(1): 97-101. Dep. Plant Breeding Genetics, Sindh Agric. Univ., Tandojam-70060, Pakistan
- Larson, J. A., R. M. Hayes, et al. (1997). Economic analysis of the harvest-aid decision for cotton in west Tennessee. *Journal of Production Agriculture* 10(3): 385-393. {a} Dep. Agric. Econ. Rural Sociol., Univ. Tennessee Agric. Exp. Stn., Knoxville, TN 37901-1071, USA
- Lege, K. E., J. T. Cothren, et al. (1997). Nitrogen fertility and leaf age effects on ethylene production of cotton in a controlled environment. *Plant Growth Regulation* 22(1): 23-28. {a} Sure-Grow Seed Inc., Route 1 Box 310, Centre, AL 35960, USA
- Leidi, E. O. and J. F. Sairz (1997). Is salinity tolerance related to Na accumulation in Upland cotton (*Gossypium hirsutum*) seedlings? *Plant and Soil* 190(1): 67-75. {a} Dep. de Biologia Vegetal, CSIC, Inst. de Recursos Naturales y Agrobiologia de Sevilla, Apdo. 1052 EP, 41080 Sevilla, Spain
- Liang, Z. L. C., R. Jiang, et al. (1997). New red flower germplasm lines of cotton selected from hybrid of *Gossypium hirsutum* X *G. bickii*. *Science in China Series C Life Sciences* 40(3): 284-292. Inst. Genetics, Chinese Acad. Sci., Beijing 100101, China
- Lin, H., S. S. Salus, et al. (1997). Salt sensitivity and the activities of the H⁺-ATPases in cotton seedlings. *Crop Science* 37(1): 190-197. {a} Dep. Plant Sci., Univ. Arizona, Tucson, Az 85721, USA
- Loera, G. J., D. A. Wolfenbarger, et al. (1997). Toxicity of azinphosmethyl, methyl parathion, and oxamyl against the boll weevil (Coleoptera: Curculionidae) in Texas, Mexico, and Guatemala. *Journal of Agricultural Entomology* 14(4): 355-361. {a} Instituto Nacional de Investigaciones Forestales y Agropecuarias, Apartado Postal 172, Rio Bravo, Tamaulipas, Mexico
- Loganathan, M. and S. Uthamasamy (1997). Effect of types of pheromone traps on moth catches of *Heliothis armigera* Hubner on cotton. *Journal of Entomological Research New Delhi* 21(2): 129-135. Dep. Entomol., Tamil Nadu Agric. Univ., Coimbatore -641 003, Tamil Nadu, India
- Lu, Z., J. Chen, et al. (1997). Photosynthetic rate, stomatal conductance and leaf area in two cotton species (*Gossypium barbadense* and *Gossypium hirsutum*) and their relation with heat resistance and yield. *Australian Journal of Plant Physiology* 24(5): 693-700. {a} UCLA-DOE Lab. Dep. Biol., Univ. California, Los Angeles, CA 90024, USA
- Ludovico, A. and W. R. S. Mattos (1997). Evaluation sugar cane based diets (*Saccharum officinarum* L.) an different whole cottonseed (*Gossypium hirsutum* L.) levels. *Revista Brasileira de Zootecnia* 26(2): 403-410. {a} EMATER-Parana, 86300-000 Cornelio Procopio, PR, Brazil
- Ma, D. P., H. C. Liu, et al. (1997). Cloning and characterization of a cotton lipid transfer protein gene specifically expressed in fiber cells. *Biochimica et Biophysica Acta* 1344(2): 111-114. {a} Dep. Biochemistry Molecular Biology, Box 9650, Mississippi State Univ., Mississippi, MS 39762, USA
- Maas, S. J. (1997). Structure and reflectance of irrigated cotton leaf canopies. *Agronomy Journal* 89(1): 54-59. USDA-ARS, 17053 Shafter Ave., Shafter, CA 93263, USA

- Manjappa, K., A. K. Guggari, et al. (1997). Effect of spacing and fertilizers on yield of cotton (*Gossypium hirsutum*) under irrigated condition. *Indian Journal of Agronomy* 42(1): 184-187. {a} ARS, Paddy, Sirsi 581 401, India
- Mann, J. E., S. G. Turnipseed, et al. (1997). Effects of early-season loss of flower buds on yield, quality, and maturity of cotton in South Carolina. *Journal of Economic Entomology* 90(5): 1324-1331. {a} Edisto Res. Educ. Cent., Clemson Univ., P.O. Box 247, Blackville, SC 29817, USA
- Mansour, M. H., N. M. Zohdy, et al. (1997). The relationship between tannins concentration in some cotton varieties and susceptibility to piercing sucking insects. *Journal of Applied Entomology* 121(6): 321-325. {a} Natl. Res. Cent. Pest Plant Protection Dep., Dokki, Tahrir St., Cairo, Egypt
- May, O. L. and D. S. Howle (1997). Registration of six germplasm lines of upland cotton: PD 93009, PD 93019, PD 93021, PD 93030, PD 93034 and PD 93057. *Crop Science* 37(3): 1030-1031. {a} USDA-ARS, Dep. Agron., Clemson Univ., 2200 Pocket Rd., Florence, SC 29506-9706, USA
- May, O. L. and D. S. Howle (1997). Registration of three germplasm lines of upland cotton: PD 93007, PD 93043, and PD 83046. *Crop Science* 37(3): 1030. {a} USDA-ARS, Dep. Agron., Clemson Univ., 2200 Pocket Rd., Florence, SC 29506-9706, USA
- McAuslane, H. J., H. T. Alborn, et al. (1997). Systemic induction of terpenoid aldehydes in cotton pigment glands by feeding of larval *Spodoptera exigua*. *Journal of Chemical Ecology* 23(12): 2861-2879. {a} Dep. Entomol. Nematol., Univ. Florida, P.O. Box 110620, Gainesville, FL 32611-0620, USA
- McConnell, J. S. and W. H. Baker (1997). Timing of early season nitrogen fertilization of cotton. *Arkansas Agricultural Experiment Station Research Series*(455): 27-29. {a} SEREC, USDA, Booneville, AR, USA
- McConnell, J. S., W. H. Baker, et al. (1997). Irrigation methods and nitrogen fertilization rates in cotton production. *Arkansas Agricultural Experiment Station Research Series*(455): 30-35. {a} SEREC, USDA-Booneville, AR, USA
- McConnell, J. S., W. H. Baker, et al. (1997). Foliar nitrogen fertilization of cotton in Southeast Arkansas. *Arkansas Agricultural Experiment Station Research Series*(455): 22-26. {a} SEREC, USDA-Booneville, Boneville, AR, USA
- Meagher, R. L., Jr., C. W. Smith, et al. (1997). Preference of *Gossypium* genotypes to *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Journal of Economic Entomology* 90(4): 1046-1052. {a} USDA-ARS CMAVE, 1700 S.W. 23rd Drive, Gainesville, FL 32604, USA
- Medeiros, F. S., A. C. Fraga, et al. (1997). The effect of type and harvest time over cotton (*Gossypium hirsutum* L.) fiber quality. *Ciencia e Agrotecnologia* 21(1): 44-49. {a} Eng. Agr. Pesq. Empresa Pesquisa Agropecuaria Rio Grande Norte-Natal-RN, Brazil
- Meredith, W. R., Jr., J. J. Heitholt, et al. (1997). Comparison of obsolete and modern cotton cultivars at two nitrogen levels. *Crop Science* 37(5): 1453-1457. {a} USDA-ARS, Cotton Physiol. Genet., P.O. Box 345, Stoneville, MS 38776, USA
- Milne, M. and G. H. Walter (1997). The significance of prey in the diet of the phytophagous thrips, *Frankliniella schultzei*. *Ecological Entomology* 22(1): 74-81. {a} Dep. Entomol., Univ. Queensl., Brisbane, QLD 4072, Australia
- Morales, L., P. Cena, et al. (1997). Resistance of cotton genotypes to *Anthonomus grandis* Boh., *Frankliniella* sp. and *Aphis gossypii* Glover. *Anais da Sociedade Entomologica do Brasil* 26(1): 93-97. {a} EMATER-Parana, Av. Inglaterra 910, 86046-430 Londrina, PR, Brazil
- Morrison, J. E., Jr., D. F. Williams, et al. (1997). Damage to dry crop seed by red imported fire ant (Hymenoptera: Formicidae). *Journal of Economic Entomology* 90(1): 218-222. {a} Nat. Resources Conservation Res. Unit, USDA-ARS, GSWRL, 808 E. Blackland Road, Temple, TX 76502, USA

- Morton, B. R. and J. A. Levin (1997). The atypical codon usage of the plant psbA gene may be the remnant of an ancestral bias. *Proceedings of the National Academy of Sciences of the United States of America* 94(21): 11434-11438. {a} Dep. Biol. Sci., Barnard Coll., Columbia Univ., 3009 Broadway, New York, NY 10027, USA
- Mullins, G. L. and C. H. Burmester (1997). Starter fertilizer and the method and rate of potassium fertilizer effects on cotton grown on soils with and without winter grazing by cattle. *Communications in Soil Science and Plant Analysis* 28(9-10): 739-746. Dep. Agron. and Soils, Auburn Univ., Auburn, AL 36849-5412, USA
- Mullins, G. L., C. H. Burmester, et al. (1997). Cotton response to in-row subsoiling and potassium fertilizer placement in Alabama. *Soil and Tillage Research* 40(3-4): 145-154. {a} Dep. Agronomy Soils, 202 Funchess Hall, Auburn Univ., Auburn, AL 36849-5412, USA
- Mulrooney, J. E., K. D. Howard, et al. (1997). Application of ultra-low-volume malathion by air assisted ground sprayer for boll weevil (Coleoptera: Curculionidae) control. *Journal of Economic Entomology* 90(2): 639-645. {a} Application Production Technol. Res. Unit, USDA-ARS, Stoneville, MS 38776, USA
- Murphy, G. P. and D. R. Shaw (1997). Field mobility of flumetsulam in three Mississippi soils. *Weed Science* 45(4): 564-567. {a} Dep. Plant Soil Sci., Miss. State Univ., Mississippi State Univ., Mississippi State, MS 39762, USA
- Nadarajan, N. and S. R. S. Rangasamy (1997). Breeding value of a fuzzless-lintless mutant in *Gossypium hirsutum* L. *Indian Journal of Genetics and Plant Breeding* 57(3): 233-237. {a} Dep. Agric. Bot., Agric. Coll. Res. Inst., Madurai 625104, India
- Naranjo, S. E., J. W. Diehl, et al. (1997). Sampling whiteflies in cotton: Validation and analysis of enumerative and binomial plans. *Environmental Entomology* 26(4): 777-788. {a} Western Cotton Res. Lab., USDA-ARS, 4135 East Broadway Road, Phoenix, AZ 85040, USA
- Ogallo, J. L., P. B. Goodell, et al. (1997). Evaluation of NemX, a new cultivar of cotton with high resistance to *Meloidogyne incognita*. *Journal of Nematology* 29(4): 531-537. {a} Dep. Nematol., Univ. California, Riverside, CA 92521, USA
- Oosterhuis, D. M., A. Steger, et al. (1997). Physiological research on plant potassium nutrition at the University of Arkansas. *Arkansas Agricultural Experiment Station Research Series*(455): 18-21. {a} Agronomy Dep., Fayetteville, AR, USA
- Orford, S. J. and J. N. Timmis (1997). Abundant mRNAs specific to the developing cotton fibre. *Theoretical and Applied Genetics* 94(6-7): 909-918. {a} Dep. Genetics, Univ. Adelaide, Adelaide 5005, Australia
- Pannetier, C., M. Giband, et al. (1997). Introduction of new traits into cotton through genetic engineering: Insect resistance as example. *Euphytica* 96(1): 163-166. {a} Lab. de Biologie Cellulaire, INRA, Centre de Versailles, 78026 Versailles Cedex, France
- Pare, P. W. and J. H. Tumlinson (1997). De novo biosynthesis of volatiles induced by insect herbivory in cotton plants. *Plant Physiology Rockville* 114(4): 1161-1167. {a} Vet. Entomol., United States Dep. Agric., Agric. Res. Serv., 1700 Southwest 23rd Drive, Gainesville, FL 32608, USA
- Patel, U. G., J. C. Patel, et al. (1997). Combining ability analysis for seed cotton yield and mean fibre length in upland cotton (*G. hirsutum* (L.)). *Indian Journal of Genetics and Plant Breeding* 57(3): 315-318. Main Cotton Res. Stn., Gujarat Agric. Univ., Surat 395007, India
- Pattinson, G. S., D. I. Warton, et al. (1997). The fungicides Terrazole and Terraclor and the nematicide Fenamiphos have little effect on root colonisation by *Glomus mosseae* and growth of cotton seedlings. *Mycorrhiza* 7(3): 155-159. {a} Sch. Biol. Sci. A12, Univ. Sydney, Sydney, NSW 2006, Australia
- Payton, P., R. D. Allen, et al. (1997). Over-expression of chloroplast-targeted Mn superoxide dismutase in cotton (*Gossypium hirsutum* L., cv. Coker 312) does not alter the reduction of photosynthesis after short

- exposures to low temperature and high light intensity. *Photosynthesis Research* 52(3): 233-244. {a} Dep. Biological Sciences, Texas Tech Univ., Lubbock, TX 79409-3131, USA
- Pettigrew, W. T. and W. R. Meredith, Jr. (1997). Dry matter production, nutrient uptake, and growth of cotton as affected by potassium fertilization. *Journal of Plant Nutrition* 20(4-5): 531-548. {a} USDA-ARS, Cotton Physiol. Genet. Res., P.O. Box 345, Stoneville, MS 38776, USA
- Qadir, M. and M. Shams (1997). Some agronomic and physiological aspects of salt tolerance in cotton (*Gossypium hirsutum* L.). *Journal of Agronomy and Crop Science* 179(2): 101-106. {a} Dep. Soil Sci., Univ. Agric., Faisalabad-38040, Pakistan
- Raina, A. K., D. M. Jackson, et al. (1997). Increased pheromone production in wild tobacco budworm (Lepidoptera: Noctuidae) exposed to host plants and host chemicals. *Environmental Entomology* 26(1): 101-105. {a} Insect Biocontrol Lab., USDA-ARS, Building 306, BARC East, Beltsville, MD 20705, USA
- Rajarajeswari, V. and I. V. Subbarao (1997). Gossypol glands in relation to resistance to bollworm (*Helicoverpa armigera*) in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 67(7): 293-295. {a} Regional Agricultural Res. Station, APAU, Anakapalle 531 001, India
- Ramraj, V. M., B. N. Vyas, et al. (1997). Effects of 28-homobrassinolide on yields of wheat, rice, groundnut, mustard, potato and cotton. *Journal of Agricultural Science* 128(4): 405-413. {a} Res. Cent., Godrej Soaps Ltd., Vikhroli, Bombay 400 079, India
- Reddy, K. R., H. J. Hodges, et al. (1997). Modeling temperature effects on cotton internode and leaf growth. *Crop Science* 37(2): 503-509. {a} Dep. Plant Soil Sci., Box 9555, Mississippi State University, Mississippi State, MS 39762, USA
- Reddy, V. R., K. R. Reddy, et al. (1997). Temperature and aldicarb effects on cotton root growth and development. *Biotronics* 26: 1-11. {a} USDA-ARS, Remote Sensing Modeling Lab., 10300 Baltimore Ave., Bldg. 007, Rm. 008, Beltsville, MD 20705, USA
- Reddy, V. R., Z. Wang, et al. (1997). Growth responses of cotton to aldicarb and temperature. *Environmental and Experimental Botany* 38(1): 39-48. {a} USDA-ARS, Remote Sensing and Modeling Lab., 10300 Baltimore Avenue, Build. 007, Room 008, Beltsville, MD 20705, USA
- Roberts, R. K., D. C. Gerloff, et al. (1997). Economics of foliar-applied potassium for conventional and no-till cotton. *Journal of Production Agriculture* 10(4): 585-589. {a} Dep. Agric. Econ. Rural Sociol., Univ. Tenn., Knoxville, TN 37901, USA
- Robinson, A. F. and A. E. Percival (1997). Resistance of *Meloidogyne incognita* race 3 and *Rotylenchulus reniformis* in wild accessions of *Gossypium hirsutum* and *G. barbadense* from Mexico. *Journal of Nematology* 29(4 Suppl.): 746-755. USDA ARS, Southern Crops Res. Lab., 2765 F and B Road, College Station, TX 77845, USA
- Robinson, M., J. N. Jenkins, et al. (1997). Root-knot nematode resistance of F-2 cotton hybrids from crosses of resistant germplasm and commercial cultivars. *Crop Science* 37(4): 1041-1046. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. BOx 5367, Mississippi State, MS 39762, USA
- Rochester, I. J., G. A. Constable, et al. (1997). Retention of cotton stubble enhances N fertilizer recovery and lint yield of irrigated cotton. *Soil and Tillage Research* 41(1-2): 75-86. {a} CRC Sustainable Cotton Prod., CSIRO Div. Plant Industry, Locked Bag 59, Narrabri, NSW 2390, Australia
- Rosenheim, J. A., L. R. Wilhoit, et al. (1997). Plant compensation, natural biological control, and herbivory by *Aphis gossypii* on pre-reproductive cotton: The anatomy of a non-pest. *Entomologia Experimentalis et Applicata* 85(1): 45-63. {a} Dep. Entomol., Univ. Calif., Davis, CA 95616, USA

Rosolem, C. A. and G. B. Bastos (1997). Mineral deficiency symptoms in cotton cultivar IAC 22. *Bragantia* 56(2): 377-387. {a} Dep. Agric. Melhoramento Vegetal, FCA/UNESP, Caixa Postal 237, 18603-970 Botucatu, SP, Brazil

Ruan, Y. L., P. S. Chourey, et al. (1997). The differential expression of sucrose synthase in relation to diverse patterns of carbon partitioning in developing cotton seed. *Plant Physiology Rockville* 115(2): 375-385. {a} Program Plant Molecular Cellular Biol., Dep. Plant Pathol., Univ. Florida, Gainesville, FL 32611, USA

Ruttledge, T. R. and E. B. Nelson (1997). Extracted fatty acids from *Gossypium hirsutum* stimulatory to the seed-rotting fungus, *Pythium ultimum*. *Phytochemistry Oxford* 46(1): 77-82. {a} Dep. Chem., Earlham Coll., Richmond, IN 47374-4095, USA

Sadras, V. O. (1997). Effects of simulated insect damage and weed interference on cotton growth and reproduction. *Annals of Applied Biology* 130(2): 271-281. CSIRO Plant Industry, Narrabri, NSW 2390, Australia

Sadras, V. O., M. P. Bange, et al. (1997). Reproductive allocation of cotton in response to plant and environmental factors. *Annals of Botany London* 80(1): 75-81. {a} CSIRO Plant Ind., Locked Bag 59, Narrabri, NSW 2390, Australia

Sadras, V. O. and G. P. Fitt (1997). Apical dominance-variability among cotton genotypes and its association with resistance to insect herbivory. *Environmental and Experimental Botany* 38(2): 145-153. {a} CSIRO Plant Ind., Locked Bag 59, Narrabri, NSW 2390, Australia

Sadras, V. O. and L. J. Wilson (1997). Nitrogen accumulation and partitioning in shoots of cotton plants infested with two-spotted spider mites. *Australian Journal of Agricultural Research* 48(4): 525-533. CSIRO Plant Industry, Locked Bag 59, Narrabri, NSW 2390, Australia

Sadras, V. O. and L. J. Wilson (1997). Growth analysis of cotton crops infested with spider mites: II. Partitioning of dry matter. *Crop Science* 37(2): 492-497. {a} C.S.I.R.O., Div. Plant Industry, Cotton Res. Unit, P.O.B. 59, Narrabri 2390, NSW, Australia

Sadras, V. O. and L. J. Wilson (1997). Growth analysis of cotton crops infested with spider mites: I. Light interception and radiation-use efficiency. *Crop Science* 37(2): 481-491. {a} C.S.I.R.O. Div. Plant Industry, Cotton Res. Unit, P.O.B. 59, Narrabri 2390, N.S.W., Australia

Salinas, G. J. R., J. E. Matocha, et al. (1997). Long-term tillage and nitrogen fertilization effects on soil properties of an Alfisol under dryland corn/cotton production. *Soil and Tillage Research* 42(1-2): 79-93. {a} Dep. Soil Crops Sciences, Texas Agricultural Experiment Station, Texas A and M Univ., College Station, TX 77843-2474, USA

Sangiliyandi, G., V. Ganesan, et al. (1997). Differential phytotoxic responses of cotton to dinitroaniline herbicides. *Indian Journal of Plant Physiology* 2(4): 275-278. {a} Sch. Biological Sciences, Madurai Kamaraj Univ., Madurai 625 021, India

Saranga, Y., N. Sass, et al. (1997). Effects of moths on lint quality of interspecific cotton hybrids. *Crop Science* 37(5): 1577-1581. {a} Heb. Univ. Jerusalem, Fac. Agric. Food Environ. Quality Sci., Dep. Field Crops Veg. Genet., P.O. Box 12, Rehovot 76100, Israel

Sarma, D., N. N. Sarma, et al. (1997). Performance off upland cotton (*Gossypium hirsutum*) varieties for early-sowing condition in hill zone of Assam. *Indian Journal of Agronomy* 42(2): 377-380. Regional Agric. Res. Stn., Assam Agric. Univ., Diphu 782 460, India

Sarma, N. N., D. Sarma, et al. (1997). Compatibility of intercropping greengram (*Phaseolus radiatus*) and sesame (*Sesamum indicum*) in rainfed cotton (*Gossypium* species). *Indian Journal of Agronomy* 42(4): 573-575. Regional Agric. Res. Stn., Assam Agric. Univ., Diphu 782 460, India

Sayal, O. U., S. A. Jatoi, et al. (1997). Estimation of combining ability for quantitative traits in *Gossypium hirsutum* L. using Triffing's techniques of diallel. *Scientific Khyber* 10(2): 13-21. Fac. Agric., Gomal Univ., Dera Ismial Khan, Pakistan

Schneider, B., C. Marcone, et al. (1997). Characterization and classification of phytoplasmas from wild and cultivated plants RFLP and sequence analysis of ribosomal DNA. *European Journal of Plant Pathology* 103(8): 675-686. {a} Biol. Bundesanstalt Land- Forstwirtschaft, Inst. Pflanzenschutz Obstbau, D-69221 Dossenheim, Germany

Shetty, P. S., C. K. Rudramuniyappa, et al. (1997). Characterization of mitochondrial proteins from cytoplasmic male sterile (CMS) and fertile lines of cotton, *Gossypium hirsutum* L. *Phytomorphology* 47(4): 383-388. Dep. Bot., Karnatak Univ., Dharwad 580003, Karnataka, India

Shimizu, Y., S. Aotsuka, et al. (1997). Changes in levels of mRNAs for cell wall-related enzymes in growing cotton fiber cells. *Plant and Cell Physiology* 38(3): 375-378. {a} Wood Res. Inst., Kyoto Univ., Gokasho, Uji, Kyoto 611, Japan

Shuman, L. M. and Z. Li (1997). Amelioration of zinc toxicity in cotton using lime or mushroom compost. *Journal of Soil Contamination* 6(4): 425-438. Crop Soil Sci. Dep., Univ. Ga., Ga. Experiment Station, Griffin, GA 30223-1797, USA

Simwat, G. S. and A. K. Dhawan (1997). Field evaluation of different economic threshold levels for chemical control of pink bollworm (*Pectinophora gossypiella*), spotted bollworm (*Earias vittella*, *E. insulana*) and cotton bollworm (*Helicoverpa armigera*) on cotton (*Gossypium* species). *Indian Journal of Agricultural Sciences* 67(6): 259-261. {a} Dep. Entomol., Punjab Agricultural Univ., Ludhiana 141 004, India

Singh, T. and A. S. Sidhu (1997). Effect of foliar application of biozyme on growth and yield of cotton (*Gossypium hirsutum* L.). *Annals of Biology Hissar* 13(2): 317-318. PAU Regional Research Station, Bathinda-151 001, India

Singh, V. P. and A. Arora (1997). Physiological characterization of cotton (*Gossypium hirsutum* L.) genotypes. *Indian Journal of Plant Physiology* 2(4): 312-314. Div. Plant Physiol., Indian Agric. Res. Inst., New Delhi 110 012, India

Smit, F. and I. A. Dubery (1997). Cell wall reinforcement in cotton hypocotyls in response to a *Verticillium dahliae* elicitor. *Phytochemistry Oxford* 44(5): 811-815. {a} Dep. Biochem., RAU-Univ., P.O. Box 524, Auckland Park 2006, South Africa

Smith, C. W. and G. G. Coyle (1997). Association of fiber quality parameters and within-boll yield components in upland cotton. *Crop Science* 37(6): 1775-1779. {a} Dep. Soil Crop Sci., Tex. A and M Univ., College Station, TX 77843-2474, USA

Song, P. and R. D. Allen (1997). Identification of a cotton fiber-specific acyl carrier protein cDNA by differential display. *Biochimica et Biophysica Acta* 1351(3): 305-312. {a} Dep. Biol. Sci., Dep. Plant Soil Sci., Texas Tech Univ., Lubbock, TX 79409, USA

Stapel, J. O., A. M. Cortesero, et al. (1997). Extrafloral nectar, honeydew, and sucrose effects on searching behavior and efficiency of *Microplitis croceipes* (Hymenoptera: Braconidae) in cotton. *Environmental Entomology* 26(3): 617-623. {a} Insect Biol. Population Management Res. Lab., USDA -ARS, Tifton, GA 31793, USA

Stewart, A. M. and K. L. Edmisten (1997). Effect of planting date, starter fertilizer, and fungicide interaction on cotton. *Journal of Plant Nutrition* 20(9): 1225-1230. Crop Sci. Dep., N.C. State Univ., Raleigh, NC 27695-7620, USA

Teixeira, H., J. C. Machado, et al. (1997). *Colletotrichum gossypii* effects and transmissibility upon delinted cotton seeds. *Fitopatologia Brasileira* 22(4): 465-471. {a} Dep. Fitopatol., Univ. Fed. Lavras, Caixa Postal 37, CEP 37200-000 Lavras, MG, Brazil

Teixeira, H., J. D. C. Machado, et al. (1997). Evaluation of the effects of chemical and biological treatments upon *Colletotrichum gossypii* South. transmission in cotton seeds (*Gossypium hirsutum* L.). *Ciencia e Agrotecnologia* 21(4): 413-418. {a} Dep. Fitossanidade-Fitopatologia, UFLA, Lavras, MG, Brazil

Thanisawanyangkura, S., H. Sinoquet, et al. (1997). Leaf orientation and sunlit leaf area distribution in cotton. *Agricultural and Forest Meteorology* 86(1-2): 1-15. {a} INRA Bioclimatol.-PIAF, Domaine de Crouelle, 63039 Clermont-Ferrand Cedex 02, France

Tomar, R. S. S., R. K. Sharma, et al. (1997). Studies on cotton plus legumes intercropping systems in relation to planting patterns. *Crop Research Hisar* 14(3): 415-420. Coll. Agric., Indore, India

Unruh, B. L. and J. C. Silvertooth (1997). Planting and irrigation termination timing effects on the yield of Upland and Pima cotton. *Journal of Production Agriculture* 10(1): 74-79. {a} Dep. Plant Sciences, University Arizona, Tucson, AZ 85721, USA

Van, E. G. A., D. T. Bowman, et al. (1997). Pedigrees and distinguishing characteristics of Upland and Pima cotton germplasm lines released between 1972 and 1996. *North Carolina Agricultural Research Service Technical Bulletin*(312): 1-80. {a} Crop Sci. Dep., NC State Univ., Raleigh, NC, USA

Van, T. H. J. and S. Saha (1997). Cotton fibers can undergo cell division. *American Journal of Botany* 84(9): 1231-1235. {a} Biol. Dep., Brookhaven Natl. Lab., Upton, NY 11973, USA

Wang, H. and R. M. Davis (1997). Susceptibility of selected cotton cultivars to seedling disease pathogens and benefits of chemical seed treatments. *Plant Disease* 81(9): 1085-1088. {a} Dep. Plant Pathol., Univ. Calif., Davis 95616-8680, USA

Wauchope, R. D., H. R. Sumner, et al. (1997). A measurement of the total mass of spray and irrigation mixtures intercepted by small whole plants. *Weed Technology* 11(3): 466-472. Nematodes, Weeds Crops Res. Unit, United States Dep. Agriculture, Agricultural Res. Service, Tifton, GA 31794, USA

Webster, T. M. and H. D. Coble (1997). Purple nutsedge (*Cyperus rotundus*) management in corn (*Zea mays*) and cotton (*Gossypium hirsutum*) rotations. *Weed Technology* 11(3): 543-548. {a} Dep. Horticulture Crop Sci., Ohio Agricultural Res. Development Cent., Ohio State Univ., Wooster, OH 44691, USA

Webster, T. M. and H. D. Coble (1997). Changes in the weed species composition of the southern United States: 1974 to 1995. *Weed Technology* 11(2): 308-317. {a} Dep. Horticulture Crop Sci., Ohio Agricultural Res. Development Cent., Ohio State Univ., Wooster, OH 44691, USA

Wei, Y. A., D. L. Hendrix, et al. (1997). Diglucomelezitose, a novel pentasaccharide in silverleaf whitefly honeydew. *Journal of Agricultural and Food Chemistry* 45(9): 3481-3486. {a} West. Cotton Res. Lab., USDA-ARS, 4135 East Broadway Road, Phoenix, AZ 85040, USA

Weil, R. R., C. D. Foy, et al. (1997). Influence of soil moisture regimes on subsequent soil manganese availability and toxicity in two cotton genotypes. *Agronomy Journal* 89(1): 1-8. {a} Agronomy Dep., Univ. Maryland, College Park, MD 20742, USA

Wheeler, T. A., J. R. Gannaway, et al. (1997). Influence of tillage, seed quality, and fungicide seed treatments on cotton emergence and yield. *Journal of Production Agriculture* 10(3): 394-400. {a} Texas A and M Univ., Rt. 3, Box 219, Lubbock, TX 79401, USA

Wilcut, J. W., D. L. Jordan, et al. (1997). Weed management in cotton (*Gossypium hirsutum*) with soil-applied and post-directed herbicides. *Weed Technology* 11(2): 221-226. {a} Crop Sci. Dep., Box 7620, North Carolina State Univ., Raleigh, NC 27695-7620, USA

Yong, J. W. H., S. C. Wong, et al. (1997). Stomatal responses to changes in vapour pressure difference between leaf and air. *Plant Cell and Environment* 20(10): 1213-1216. {a} Environ. Biol. Group, Res. Sch. Biol. Sci., Inst. Advanced Studies, Australian Natl. Univ., GPO Box 475, ACT 2601, Canberra, Australia

- Yuan, Z. F. and S. Y. Hu (1997). Ultrastructural changes in hydrating and germinating pollen grain of cotton, with special reference to endoplasmic reticulum and coated vesicles. *Acta Botanica Sinica* 39(12): 1085-1090. {a} Coll. Life Sci., Peking Univ., Beijing 100871, China
- Zhang, J. S. Y. H. Y., L. Zhu, et al. (1997). Ultracytochemical localization of calcium in the pollen tube track of cotton gynoecium. *Acta Botanica Sinica* 39(2): 121-125. {a} Lab. Plant Reprod. Biol., Coll. Life Sci., Wuhan Univ., Wuhan 430072, China
- Zhang, T. Z., J. J. Pan, et al. (1997). Interaction of virescent genes in upland cotton (*Gossypium hirsutum* L.): Chlorophyll content. *Crop Science* 37(4): 1123-1126. {a} USDA-ARS, Southern Crops Res. Lab., Crop Germplasm Res. Unit, 2765 F and B Rd., College Station, TX 77845, USA
- Zhao, D. and D. Oosterhuis (1997). Physiological response of growth chamber-grown cotton plants to the plant growth regulator PGR-IV under water-deficit stress. *Environmental and Experimental Botany* 38(1): 7-14. {a} Dep. Agronomy, Univ. Arkansas, Fayetteville, AR 72701, USA
- Zholkevich, V. N., N. K. Zubkova, et al. (1997). Interaction of heat shock and water stress in plants: 2. Osmoregulation in cotton leaves exposed to subsequent short-term hyperthermia and soil drought. *Fiziologiya Rastanii* 44(4): 613-623. {a} K.A. Timiryazev Inst. Plant Physiol., Russ. Acad. Sci., Botanicheskaya ul. 35, Moscow 127276, Russia
- Adrian, J. L., Jr., P. A. Duffy, et al. (1996). Turfgrass-sod: A viable farm enterprise. *Journal of Production Agriculture* 9(2): 276-283. {a} 308A Comer Hall, Dep. Agric. Economic Rural Sociol., Auburn University, AL 36849, USA
- Alborn, H. T., U. S. R. Rose, et al. (1996). Systemic induction of feeding deterrents in cotton plants by feeding of *Spodoptera* spp. larvae. *Journal of Chemical Ecology* 22(5): 919-932. {a} Dep. Chem. Ecol., Goteborg Univ., S-413 19 Goteborg, Sweden
- Ali, A. and S. Y. Young (1996). Activity of *Bacillus thuringiensis* Berliner against different ages and stages of *Helicoverpa zea* (Lepidoptera: Noctuidae) on cotton. *Journal of Entomological Science* 31(1): 1-8. {a} Jaffer Brothers Ltd., Ex. 9C Shamsabad Colony, Khanewal Road, Multan, Pakistan
- Babu, R., M. V. H. Rao, et al. (1996). Cotton (*Gossypium hirsutum*)-based multiple cropping systems for sustained yield. *Indian Journal of Agricultural Sciences* 66(10): 577-580. {a} Dep. Agron., Coll. Agric., UAS, Dharwad 580 005, India
- Baird, R. E., D. E. Carling, et al. (1996). Characterization and comparison of isolates of *Rhizoctonia solani* AG-7 from Arkansas, Indiana, and Japan, and select AG-4 isolates. *Plant Disease* 80(12): 1421-1424. {a} Plant Pathol. Dep., Univ. Georgia, RDC, PO Box 1209, Tifton, GA 31793, USA
- Baird, R. E., R. F. Davis, et al. (1996). Frequency and geographical distribution of plant-parasitic nematodes on cotton in Georgia. *Journal of Nematology* 28(4 Suppl.): 661-667. Univ. Georgia, Dep. Plant Pathol., Cooperative Extension Service, Rural Development Center, P.O. Box 1209, Tifton, GA 31793, USA
- Baker, W. H., J. S. McConnell, et al. (1996). Cotton response to supplemental foliar nitrogen in the central Arkansas delta. *Arkansas Agricultural Experiment Station Research Series*(450): 78-80.
- Banuelos, G. S., B. Mackey, et al. (1996). Response of cotton and kenaf to boron-amended water and soil. *Crop Science* 36(1): 158-164. {a} USDA, ARS, Water Manage. Res. Lab., 2021 S. Peach Ave., Fresno, CA, USA
- Basu, A. K. (1996). Current genetic research in cotton in India. *Genetica Dordrecht* 97(3): 279-290. The Cotton Corp. of India Ltd., Air India Bldg. 12th Flr., Nariman Point, Bombay 400 021, India
- Bauer, P. J. and J. M. Bradow (1996). Cotton genotype response to early-season cold temperatures. *Crop Science* 36(6): 1602-1606. {a} USDA-ARS, 2611 W. Lucas St., Florence, SC 29501-1241, USA

Bauer, P. J. and W. J. Busscher (1996). Winter cover and tillage influences on coastal plain cotton production. *Journal of Production Agriculture* 9(1): 50-54. {a} USDA-ARS Coastal Plains Soil Water Plant Res. Cent., 2611 W. Lucas St., Florence, SC 29501-1241, USA

Bauer, P. J. and C. C. Green (1996). Evaluation of F-2 genotypes of cotton for conservation tillage. *Crop Science* 36(3): 655-658. {a} USDA-ARS Coastal Plains Soil, Water, Plant Res. Cent., 2611 W. Lucas St., Florence, SC 29501-1241, USA

Baumhardt, R. L. and R. J. Lascano (1996). Rain infiltration as affected by wheat residue amount and distribution in ridged tillage. *Soil Science Society of America Journal* 60(6): 1908-1913. {a} Texas Agric. Exp. Stn., Route 3, Box 219, Lubbock, TX 79401, USA

Bednarz, C. W. and D. M. Oosterhuis (1996). Partitioning of potassium in the cotton plant during the development of a potassium deficiency. *Journal of Plant Nutrition* 19(12): 1629-1638. {a} Miss. State Univ., Delta Res. Extension Center, P.O. Box 69, Stonesville, MS 38776, USA

Bednarz, C. W. and D. M. Oosterhuis (1996). Partitioning of potassium in field-grown cotton plants under various levels of available potassium. *Arkansas Agricultural Experiment Station Research Series*(450): 75-77.

Behle, R. W., M. R. McGuire, et al. (1996). Extending the residual toxicity of *Bacillus thuringiensis* with casein-based formulations. *Journal of Economic Entomology* 89(6): 1399-1405. {a} Bioactive Agents Res. Unit, Natl. Center Agric. Utilization Res., USDA-ARS, 1815 N. University, Peoria, IL 61604, USA

Bejarano, A. J., L. M. A. Blanco, et al. (1996). Etiology, importance, and distribution of verticillium wilt of cotton in southern Spain. *Plant Disease* 80(11): 1233-1238. {a} Instituto Agricultura Sostenible, C.S.I.C., Apdo. 4084, 14080-Cordoba, Spain

Bevers, S. J., J. E. Slosser, et al. (1996). Yield and income response to planting dates and harvest aid chemicals in Texas cotton production. *Journal of Economic Entomology* 89(2): 343-346. {a} Tex. Agric. Extension Serv., P.O. Box 2159, Vernon, TX 76385-2159, USA

Bondada, B. R., D. M. Oosterhuis, et al. (1996). Effect of water stress on the epicuticular wax composition and ultrastructure of cotton *Gossypium hirsutum* (L.) leaf, bract, and boll. *Environmental and Experimental Botany* 36(1): 61-69. {a} Univ. Florida, Citrus Res. Education Cent., 700 Experiment Station Rd., Lake Alfred, FL 33850, USA

Bondada, B. R., D. M. Oosterhuis, et al. (1996). Canopy photosynthesis, growth, yield, and boll 15N accumulation under nitrogen stress in cotton. *Crop Science* 36(1): 127-133. {a} Citrus Res. Educ. Cent., Univ. Fla., 700 Experiment Stn. Rd., Lake Alfred, FL 33850, USA

Bourland, F. M. (1996). Registration of 'H1330' cotton. *Crop Science* 36(3): 813. Dep. Agronomy, 115 Plant Sci. Build., Univ. Arkansas, Fayetteville, AR 72701, USA

Bowman, D. T., O. L. May, et al. (1996). Genetic base of upland cotton cultivars released between 1970 and 1990. *Crop Science* 36(3): 577-581. {a} Dep. Crop Sci., Raleigh, NC 27695-8604, USA

Brown, B. A., R. M. Hayes, et al. (1996). Effect of long-term vetch (*Vicia villosa*) cover crop and tillage system on fluometuron dissipation from surface soil. *Weed Science* 44(1): 171-175. {a} P.O. Box 1071, Knoxville, TN 37901, USA

Brubaker, C. L., C. G. Benson, et al. (1996). Occurrence of terpenoid aldehydes and lysigenous cavities in the 'glandless' seeds of Australian *Gossypium* species. *Australian Journal of Botany* 44(5): 601-612. {a} Cent. Plant Biodiversity Res., CSIRO Div. Plant Ind., GPO Box 1600, Canberra, ACT 2601, Australia

Bunkelmann, J. R. and R. N. Trelease (1996). Ascorbate peroxidase: A prominent membrane protein in oilseed glyoxysomes. *Plant Physiology Rockville* 110(2): 589-598. {a} Dep. Botany, Arizona State Univ., Tempe, AZ 85287-1601, USA

Burla, M., M. Goverde, et al. (1996). Influence of biocontrol organisms on root pathogenic fungi and on the plant symbiotic micro-organisms *Rhizobium phaseoli* and *Glomus mosseae*. *Zeitschrift fuer Pflanzenkrankheiten und Pflanzenschutz* 103(2): 156-163. {a} Inst. Microbiol., ETH, Schmelzbergstr. 7, CH-8094 Zurich, Switzerland

Butter, N. S. and S. Singh (1996). Ovipositional response of *Helicoverpa armigera* to different cotton genotypes. *Phytoparasitica* 24(2): 97-102. {a} Directorate Extension Education, Punjab Agricultural Univ., Ludhiana 141 004, Punjab, India

Chapman, K. D. and W. B. Sprinkle (1996). Developmental, tissue-specific, and environmental factors regulate the biosynthesis of N-acylphosphatidyl-ethanolamine in cotton (*Gossypium hirsutum* L.). *Journal of Plant Physiology* 149(3-4): 277-284. {a} Dep. Biol. Sci., Div. Biochem. Mol. Biol., Univ. North Texas, PO Box 5128, Denton, TX 76203, USA

Chen, J. G., X. M. Du, et al. (1996). Fluctuation in levels of endogenous plant hormones in ovules of normal and mutant cotton during flowering and their relation to fiber development. *Journal of Plant Growth Regulation* 15(4): 173-177. {a} Dep. Agronomy, Nanjing Agricultural Univ., Nanjing 210095, China

Clark, L. E., T. R. Moore, et al. (1996). Response of cotton to cropping and tillage systems in the Texas Rolling Plains. *Journal of Production Agriculture* 9(1): 55-60. {a} Texas Agric. Exp. Stn., P.O. 1658, Vernon, TX 76385-1658, USA

Dahatonde, B. N. (1996). Efficacy of pressmud-cake with and without azotobacter seed inoculation for manuring of hybrid cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 41(4): 664-665. Dep. Agron., Dr Punjabrao Deshmukh Krishi Vidyapeeth, Akola, Maharashtra 444 104, India

Davidonis, G. H., A. Johnson, et al. (1996). Influence of low-weight seeds and motes on the fiber properties of other cotton seeds. *Field Crops Research* 48(2-3): 141-153. {a} USDA, ARS, Southern Regional Res. Cent., PO Box 19687, New Orleans, LA 70179-0687, USA

Davis, E. M., J. Tsuji, et al. (1996). Purification of (+)-delta-cadinene synthase, a sesquiterpene cyclase from bacteria-inoculated cotton foliar tissue. *Phytochemistry Oxford* 41(4): 1047-1055. {a} Dep. Biochem. Molecular Biol., Okla. State Univ., Stillwater, OK 74078-3035, USA

Dhumale, D. B., G. L. Ingole, et al. (1996). Interspecific hybridization through embryo culture in cotton, *Gossypium arboreum* and *G. hirsutum*. *Indian Journal of Experimental Biology* 34(3): 288-289. Tissue Culture Lab., Dep. Bot., Punjabrao Krishi Vidyapeeth, Akola 444 104, India

Dotray, P. A., J. W. Keeling, et al. (1996). Palmer amaranth (*Amaranthus palmeri*) and devil's-claw (*Proboscidea louisianica*) control in cotton (*Gossypium hirsutum*) with pyrithiobac. *Weed Technology* 10(1): 7-12. {a} Texas Tech Univ., Lubbock, TX 79409, USA

Dubery, I. A. and R. Meyer (1996). Specific binding of a *Verticillium dahliae* phytotoxin to protoplasts of cotton, *Gossypium hirsutum*. *Plant Cell Reports* 15(10): 777-780. {a} Dep. Biochem., Rand Afrikaans Univ., P.O. Box 524, Auckland Park 2006, South Africa

El, Z. K. M. and P. M. Thaxton (1996). Registration of 'Tamcot Sphinx' cotton. *Crop Science* 36(4): 1074. {a} Dep. Soil Crop Sci., Texas Agric. Exp. Stn., Texas A and M Univ. System, College Station, TX 77843-2474, USA

Eldon, S. and R. J. Hillocks (1996). The effect of reduced phytoalexin production on the resistance of upland cotton (*Gossypium hirsutum*) to *verticillium* and *fusarium* wilts. *Annals of Applied Biology* 129(2): 217-225. {a} Sch. Natural Sci., Halton Coll., Kingsway, Widnes, Cheshire WA8 7QQ, UK

Entry, J. A., C. C. Mitchell, et al. (1996). Influence of management practices on soil organic matter, microbial biomass and cotton yield in Alabama's "Old Rotation". *Biology and Fertility of Soils* 23(4): 353-358. {a} Dep. Agronomy Soils, 202 Funchness Hall, Auburn University, Auburn, AL 36849-5412, USA

Everitt, J. H., D. E. Escobar, et al. (1996). Using spatial information technologies for detecting and mapping whitefly and harvester ant infestations in South Texas. *Southwestern Entomologist* 21(4): 421-432. United States Dep. Agric., Agric. Research Serv., 2413 East Highway 83, Weslaco, TX 78596-8344, USA

Faver, K. L. and T. J. Gerik (1996). Foliar-applied methanol effects on cotton (*Gossypium hirsutum* L.) gas exchange and growth. *Field Crops Research* 47(2-3): 227-234. Blackland Res. Center, 808 E. Blackland Road Temple, TX 76502, USA

Faver, K. L., T. J. Gerik, et al. (1996). Late season water stress in cotton: II. Leaf gas exchange and assimilation capacity. *Crop Science* 36(4): 922-928. {a} Blackland Res. Cent., 808 E. Blackland Rd., Temple, TX 76502, USA

Ferguson, D. L., R. B. Turley, et al. (1996). Comparison of protein profiles during cotton (*Gossypium hirsutum* L.) fiber cell development with partial sequences of two proteins. *Journal of Agricultural and Food Chemistry* 44(12): 4022-4027. {a} Cotton Physiol. Genetics Res., Agric. Res. Service, U.S. Dep. Agric., P.O. Box 345, Stoneville, MS 38776-0345, USA

Fernandez, C. J., J. T. Cothren, et al. (1996). Partitioning of biomass in water- and nitrogen-stressed cotton during pre-bloom stage. *Journal of Plant Nutrition* 19(3-4): 595-617. {a} Texas A and M Univ. Agric. Res. Extension Cent., 1619 Garner Field Rd., Uvalde, TX 78801, USA

Fernandez, C. J., K. J. McInnes, et al. (1996). Water status and leaf area production in water- and nitrogen-stressed cotton. *Crop Science* 36(5): 1224-1233. {a} Texas A and M Univ. Res. Ext. Center, Uvalde, Tx 78801, USA

Flint, H. M., J. E. Leggett, et al. (1996). Effects of the plant growth regulator mepiquat chloride on silverleaf whitefly (Homoptera: Aleyrodidae) infestations on cotton. *Journal of Entomological Science* 31(1): 112-122. {a} West. Cotton Res. Lab., USDA-ARS, Phoenix, AZ 85040, USA

Flint, H. M., S. E. Naranjo, et al. (1996). Cotton water stress, arthropod dynamics, and management of *Bemisia tabaci* (Homoptera: Aleyrodidae). *Journal of Economic Entomology* 89(5): 1288-1300. Western Cotton Res. Lab., USDA-ARS, 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

Gerik, T. J., K. L. Faver, et al. (1996). Late season water stress in cotton: I. Plant growth, water use, and yield. *Crop Science* 36(4): 914-921. {a} Blackland Res. Cent., 808 E. Blackland Rd., Temple, TX 76502, USA

Gorham, J. (1996). Glycinebetaine is a major nitrogen-containing solute in the Malvaceae. *Phytochemistry Oxford* 43(2): 367-369. Centre Arid Zone Studies, Sch. Biol. Sci., Univ. Wales, Bangor LL57 2UW, UK

Gossett, D. R., S. W. Banks, et al. (1996). Antioxidant response to NaCl stress in a control and an NaCl-tolerant cotton cell line grown in the presence of paraquat, buthionine sulfoximine, and exogenous glutathione. *Plant Physiology Rockville* 112(2): 803-809. {a} Dep. Biological Sci., Louisiana State Univ.-Shreveport, One University Place, Shreveport, LA 71115, USA

Gridi, P. I. L., E. Cia, et al. (1996). Progeny evaluation for resistance to the root borer in cotton. *Bragantia* 55(1): 75-81. {a} Secao Algodao, Inst. Agron., Caixa Postal 28, 13001-970 Campinas, SP, Brazil

Guertal, E. A., A. O. Abaye, et al. (1996). Sources of boron for foliar fertilization of cotton and soybean. *Communications in Soil Science and Plant Analysis* 27(15-17): 2815-2828. {a} Dep. Agronomy and Soils, Auburn Univ., Auburn, AL 36849, USA

Halcomb, J. L., J. H. Benedict, et al. (1996). Survival and growth of bollworm and tobacco budworm on nontransgenic and transgenic cotton expressing a CryIA insecticidal protein (Lepidoptera: Noctuidae). *Environmental Entomology* 25(2): 250-255. {a} Texas A and M Univ. Agricultural Res. Extension Cent., Route 2, Box 589, Corpus Christi, TX 78410, USA

- Hanson, R. E., F. M. N. Islam, et al. (1996). Distribution of 5S and 18S-28S rDNA loci in a tetraploid cotton (*Gossypium hirsutum* L.) and its putative diploid ancestors. *Chromosoma Berlin* 105(1): 55-61. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA
- Harrison, M. A., R. M. Hayes, et al. (1996). Environment affects cotton and velvetleaf response to pyriithiobac. *Weed Science* 44(2): 241-247. {a} Dep. Plant Soil Sci., P.O. Box 1071, Knoxville, TN 37901, USA
- He, J. S. C., R. Jiang, et al. (1996). Genetic analysis of a dwarf mutant in cotton. *Acta Genetica Sinica* 23(5): 397-402. {a} Inst. Genetics, Acad. Sinica, Beijing 100101, China
- Headrick, D. H., T. S. Bellows, Jr., et al. (1996). Behaviors of female *Eretmocerus* sp. nr. *californicus* (Hymenoptera: Aphelinidae) attacking *Bemisia argentifolii* (Homoptera: Aleyrodidae) on cotton, *Gossypium hirsutum*, (Malvaceae) and melon, *Cucumis melo* (Cucurbitaceae). *Biological Control* 6(1): 64-75. Dep. Entomol., Univ. California, Riverside, Riverside, CA 92521, USA
- Heitholt, J. J., W. R. Meredith, Jr., et al. (1996). Comparison of cotton genotypes varying in canopy characteristics in 76-cm vs. 102-cm rows. *Crop Science* 36(4): 955-960. {a} USDA-ARS, Cotton Physiol. Genetics, P.O. Box 345, Stoneville, MS 38776, USA
- Henneberry, T. J., D. L. Hendrix, et al. (1996). *Bemisia argentifolii* (Homoptera: Aleyrodidae) honeydew sugars and relationships to sticky cotton. *Environmental Entomology* 25(3): 551-558. {a} Western Cotton Res. Lab., USDA-ARS, Pacific West Area, 4135 E. Broadway, Phoenix, AZ 85040-8830, USA
- Holt, J. S. and D. R. Orcutt (1996). Temperature thresholds for bud sprouting in perennial weeds and seed germination in cotton. *Weed Science* 44(3): 523-533. Dep. Botany Plant Sci., Univ. California, Riverside, CA 92521, USA
- Hood, G., R. B. Turley, et al. (1996). Ribosomal protein RL44 is encoded by two subfamilies in upland cotton (*Gossypium hirsutum* L.). *Biochemical and Biophysical Research Communications* 226(1): 32-36. {a} Dep. Biol., Delta State Univ., Cleveland, MS, USA
- Hudspeth, R. L., S. L. Hobbs, et al. (1996). Characterization and expression of chitinase and 1,3-beta-glucanase genes in cotton. *Plant Molecular Biology* 31(4): 911-916. {a} J.G. Boswell Cotton Seed Breeding Co., P.O. Box 787, 850 Plymouth Ave., Corcoran, CA 93212, USA
- Hudspeth, R. L., S. L. Hobbs, et al. (1996). Characterization and expression of metallothionein-like genes in cotton. *Plant Molecular Biology* 31(3): 701-705. Phytogen, 101 Waverly Dr., Pasadena, CA 91105, USA
- Hulme, P. J., D. C. McKenzie, et al. (1996). An evaluation of controlled traffic with reduced tillage for irrigated cotton on a Vertisol. *Soil and Tillage Research* 38(3-4): 217-237. {a} NSW Agriculture Biological, Chemical Res. Inst., Rydalmere, NSW 2116, Australia
- Hulugalle, N. R. and P. Entwistle (1996). Effect of sowing cowpea on properties of an irrigated Vertisol and growth and yield of succeeding cotton. *Australian Journal of Soil Research* 34(4): 529-544. Co-operative Res. Centre Sustainable Cotton Production, Aust. Cotton Res. Inst., New South Wales Dep. Agric., PMB, Myall Vale Mail Run, Narrabri, NSW 2390, Australia
- Hulugalle, N. R., D. L. Larsen, et al. (1996). Effect of broad beds and dolichos residue management on properties of an irrigated vertisol. *Soil Technology* 8(4): 275-286. {a} Australian Cotton Res. Inst., New South Wales Dep. Agric., PMB, Myall Vale Mail Run, Narrabri, N.S.W. 2390, Australia
- Hutmacher, R. B., J. E. Ayars, et al. (1996). Uptake of shallow groundwater by cotton: Growth stage, groundwater salinity effects in column lysimeters. *Agricultural Water Management* 31(3): 205-223. {a} USDA-ARS, Water Manage. Res. Lab., 2021 South Peach Ave., Fresno, CA 93727-5951, USA

John, M. E. (1996). Structural characterization of genes corresponding to cotton fiber mRNA, E6: Reduced E6 protein in transgenic plants by antisense gene. *Plant Molecular Biology* 30(2): 297-306. Fiber Technol. Div., Agracetus Inc., 8520 University Green, Middleton, WI 53562, USA

Johnson, D. H. and R. E. Talbert (1996). Cotton (*Gossypium hirsutum*) response to imazaquin and imazethapyr soil residues. *Weed Science* 44(1): 156-161. {a} American Cyanamid Co., P.O. Box 400, Princeton, NJ 08543-0400, USA

Jones, M. A., R. Wells, et al. (1996). Cotton response to seasonal patterns of flower removal: II. Growth and dry matter allocation. *Crop Science* 36(3): 639-645. {a} Dep. Crop Sci., North Carolina State Univ., Raleigh, NC 27695-7620, USA

Jones, M. A., R. Wells, et al. (1996). Cotton response to seasonal patterns of flower removal: I. Yield and fiber quality. *Crop Science* 36(3): 633-638. {a} Dep. Crop Sci., North Carolina State Univ., Raleigh, NC 27695-7620, USA

Karlen, D. L., P. G. Hunt, et al. (1996). Fertilizer ¹⁵Nitrogen recovery by corn, wheat, and cotton grown with and without pre-plant tillage on Norfolk loamy sand. *Crop Science* 36(4): 975-981. {a} USDA-ARS, Natl. Soil Tilth Lab., 2150 Pammel Dr., Ames, IA 50011, USA

Keeling, J. W., P. A. Dotray, et al. (1996). Effects of repeated applications of trifluralin and pendimethalin on cotton (*Gossypium hirsutum*). *Weed Technology* 10(2): 295-298. Texas Agric. Exp. Stn., Lubbock, TX 79401, USA

Keeling, J. W., A. G. Matches, et al. (1996). Comparison of interseeded legumes and small grains for cover crop establishment in cotton. *Agronomy Journal* 88(2): 219-222. {a} Tex. A and M Agric. Exp. Stn., Rt. 3, Box 219, Lubbock, TX 79401, USA

Keeton, A., E. C. Murdock, et al. (1996). Chemical control systems for coffee Senna (*Cassia occidentalis*) in cotton (*Gossypium hirsutum*). *Weed Technology* 10(3): 550-555. Dep. Agron., Clemson Univ., Clemson, SC 29634, USA

Keino, J. K., C. A. Beyrouy, et al. (1996). Effect of foliar potassium on kinetic uptake parameters of cotton. *Arkansas Agricultural Experiment Station Research Series*(450): 72-74.

Kenchana, G. P. V., B. H. Katarki, et al. (1996). Genetics of yield, yield attributes and their implications in breeding of cotton (*Gossypium hirsutum* L.). *Indian Journal of Genetics and Plant Breeding* 56(2): 147-151. {a} Dep. Genet. Plant Breeding, Univ. Agric. Sci., Dharwad 580 005, India

Khan, R. A. and W. T. Molin (1996). Photosynthetic electron transport in thylakoids from cotton cultivars (*Gossypium* sp.) differing in tolerance to prometryn. *Plant Cell Reports* 15(12): 969-973. {a} Dep. Plant Sci., Univ. Arizona, Tucson, AZ 85721, USA

Khurana, A. D. (1996). Bioefficacy of fenvalerate when tank mixed with systemic insecticides for the control of bollworms on *Hirsutum* cotton. *Journal of Insect Science* 9(2): 193-195. {a} Haryana Agric. Univ. Cotton Res. Station, Sirsa 125 055, India

Kowsalya, R. and T. S. Raveendran (1996). Genetic variability and D-2 analysis in upland cotton. *Crop Research Hisar* 12(1): 36-42. Sch. Genetics, Tamil Nadu Agricultural Univ., Coimbatore-641 003, India

Kudlicka, K., J. H. Lee, et al. (1996). A comparative analysis of in vitro cellulose synthesis from cell-free extracts of mung bean (*Vigna radiata*, Fabaceae) and cotton (*Gossypium hirsutum*, Malvaceae). *American Journal of Botany* 83(3): 274-284. {a} Dep. Bot., University Texas Austin, Austin, TX 78713, USA

Kuliev, R. A. (1996). New methods of efficiency increasing of experimental mutagenesis in the cotton-plant selection. *Tsitologiya i Genetika* 30(6): 39-43. Baku State Univ., Baku, Azerbaijan

Lasota, L. R. (1996). Evaluation by the United States Environmental Protection Agency of pesticidal substances produced in plants. *Field Crops Research* 45(1-3): 181-186. Office Prevention, Pesticides Toxic Substances Programs, United States Environmental Protection Agency, Washington, DC 20460, USA

Leavitt, S. W., E. A. Paul, et al. (1996). Carbon isotopes and carbon turnover in cotton and wheat FACE experiments. *Plant and Soil* 187(2): 147-155. {a} Lab. Tree-Ring Res., Univ. Arizona, Tucson, AZ 85721, USA

Li, J., S. I. Zingen, et al. (1996). Induction of resistance of cotton plants to *Verticillium* wilt and of tomato plants to *Fusarium* wilt by 3-aminobutyric acid and methyl jasmonate. *Zeitschrift fuer Pflanzenkrankheiten und Pflanzenschutz* 103(3): 288-299. Universitaet Hohenheim, Inst. Phytomedizin, D-70593 Stuttgart, Germany

Lordello, A. I. L. and R. R. A. Lordello (1996). Identification of *Meloidogyne incognita* races in some plants. *Summa Phytopathologica* 22(1): 43-45. {a} EMBRAPA/IAC, Secale Nematol., 13.020-902 Campinas, SP, Brazil

Lozovaya, V. V., X. Luo, et al. (1996). Rapid recovery of photosynthetic cell suspension cultures following heterotrophic bleaching. *In Vitro Cellular and Developmental Biology Plant* 32(4): 295-298. {a} Univ. Ill., Dep. Crop Sciences, Edward R. Madigan Lab., 1201 W. Gregory, Urbana, IL 61801, USA

Marur, C. J., P. Mazzafera, et al. (1996). Carbon assimilation and export in leaves of cotton plants under water deficit. *Revista Brasileira de Fisiologia Vegetal* 8(3): 181-186. {a} Inst. Agronomico Parana, IAPAR, P.O. Box 481, Londrina, 86001-970 Parana, Brazil

Mascagni, H. J., Jr. and D. J. Boquet (1996). Starter fertilizer and planting date effects on corn rotated with cotton. *Agronomy Journal* 88(6): 975-982. {a} La. State Univ. Agric. Cent., Northeast Res. Stn., St. Joseph, LA 71366, USA

May, O. L., D. S. Howle, et al. (1996). Registration of PD-314 germplasm line of upland cotton with high yield and fiber quality. *Crop Science* 36(6): 1718. {a} USDA-ARS, Clemson Univ., 2200 Pocket Rd., Florence, SC 29501-9706, USA

McCarty, J. C., Jr., J. N. Jenkins, et al. (1996). Genetic analysis of primitive cotton germplasm accessions. *Crop Science* 36(3): 581-585. {a} USDA-ARS, Crop Sci. Res. Lab., Box 5367, Mississippi State, MS 39762, USA

McConnell, J. S., W. H. Baker, et al. (1996). Distribution of residual nitrate-nitrogen in long-term fertilization studies of an alfisol cropped to cotton. *Journal of Environmental Quality* 25(6): 1389-1394. {a} Southeast Res. Ext. Cent., Univ. Arkansas, P.O. Box 3508, Monticello, AR 71656, USA

McConnell, J. S., W. H. Baker, et al. (1996). Irrigation methods and nitrogen fertilization rates in cotton production. *Arkansas Agricultural Experiment Station Research Series*(450): 67-71.

McConnell, J. S., W. H. Baker, et al. (1996). Foliar nitrogen fertilization of cotton in southeastern Arkansas. *Arkansas Agricultural Experiment Station Research Series*(450): 63-66.

McConnell, J. S., W. H. Baker, et al. (1996). Cotton yield response to irrigation, reduced tillage and cover crops. *Arkansas Agricultural Experiment Station Research Series*(449): 60-64. {a} Dep. Agron., Arkansas Agric. Exp. Stn., Fayetteville, AR 72701, USA

McInroy, J. A., G. Musson, et al. (1996). Masking of antibiotic-resistance upon recovery of endophytic bacteria. *Plant and Soil* 186(2): 213-218. {a} Dep. Plant Pathol., Biol. Control Inst., Ala. Agric. Exp. Station, Auburn Univ., Auburn, AL 36849-5409, USA

McMichael, B. L., D. R. Upchurch, et al. (1996). Soil temperature derived prediction of root density in cotton. *Environmental and Experimental Botany* 36(3): 303-312. {a} USDA-ARS, Cropping Systems Res. Lab., Rt. 3, Box 215, Lubbock, TX 79401, USA

- Medeiros, F. S., A. C. Fraga, et al. (1996). Effect of storage on the physiological quality of cotton seeds. *Ciencia e Agrotecnologia* 20(3): 284-292. Univ. Fed. Lavras, 37.200-000 Lavras, MG, Brazil
- Mehetre, S. S. and R. K. J. Narayan (1996). Evolutionary DNA variation and genome differentiation in *Gossypium* L. *Proceedings of the Indian National Science Academy Part B Biological Sciences* 63(1): 63-72. {a} Mahatma Phule Krishi Vidyapeeth, Coll. Agric., Vidyanagar, Kolhapur 416004, Maharashtra, India
- Meredith, W. R., Jr., W. T. Pettigrew, et al. (1996). Sub-okra, semi-smoothness, and nectariless effect on cotton lint yield. *Crop Science* 36(1): 22-25. {a} USDA, ARS, Cotton Physiology Genetics, P.O. Box 345, Stoneville, MS 38776-0345, USA
- Millar, A. A. and E. S. Dennis (1996). The alcohol dehydrogenase genes of cotton. *Plant Molecular Biology* 31(4): 897-904. {a} CSIRO Div. Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia
- Millar, A. A. and E. S. Dennis (1996). Protein synthesis during oxygen deprivation in cotton. *Australian Journal of Plant Physiology* 23(3): 341-348. {a} CSIRO, Div. Plant Industry, Canberra, GPO Box 1600, Canberra, ACT 2601, Australia
- Mitchell, C. C., G. Traxler, et al. (1996). Measuring sustainable cotton production using total factor productivity. *Journal of Production Agriculture* 9(2): 289-297. {a} Dep. Agron. Soils, Auburn University, AL 36849, USA
- Molin, W. T. and R. A. Khan (1996). Differential tolerance of cotton (*Gossypium* sp.) cultivars to the herbicide prometryn. *Pesticide Biochemistry and Physiology* 56(1): 1-11. {a} Dep. Plant Sci., Univ. Arizona, Tucson, AZ 85721, USA
- Monga, D. and S. Raj (1996). Varietal screening against root rot of cotton in sick field. *Crop Research Hisar* 12(1): 82-86. {a} Central Inst. Cotton Res., Regional Station, Sirsa-125 005, Haryana, India
- Monks, C. D., M. G. Patterson, et al. (1996). Effect of bensulfuron-methyl on cotton (*Gossypium hirsutum* L.) growth and development. *Weed Technology* 10(4): 851-855. Dep. Agron. Soils, Alabama Exp. Stn., Auburn Univ., Auburn, AL 36849, USA
- Moreshet, S., M. Fuchs, et al. (1996). Water transport characteristics of cotton as affected by drip irrigation layout. *Agronomy Journal* 88(5): 717-722. {a} Dep. Environmental Physics Irrigation, Inst. Soils Water, Agric. Res. Org., Volcani Center, PO Box 6, Bet Dagan, Israel
- Myaka, F. A. and J. C. B. Kabissa (1996). Fitting short duration cowpea into a cotton-based cropping system in Tanzania: Effect of planting pattern, time of planting cowpea, and insecticide application to the cotton. *Experimental Agriculture* 32(2): 225-230. {a} Ilonga Agricultural Res. Training Inst., P.O. Box Ilonga, Kilosa, Tanzania
- Nadjimov, U. K., M. S. Mirakhmedov, et al. (1996). Effects of fusaric acid on intact cotton plants (*Gossypium hirsutum* L.). *Journal of Plant Growth Regulation* 15(3): 129-131. {a} Inst. Biol. Sci. Univ. Wales, Aberystwyth, Dyfed SY23 3DA, UK
- Nandeshwar, S. B. (1996). In-vitro shoot organogenesis in cotton. *Advances in Plant Sciences* 9(1): 181-182. Tissue Culture Lab, Crop Improvement Div., Central Inst. Cotton Res., Post bag No. 225, GPO, Nagpur, India
- Naranjo, S. E., C. C. Chu, et al. (1996). Economic injury levels for *Bemisia tabaci* (Homoptera: Aleyrodidae) in cotton: Impact of crop price, control costs, and efficacy of control. *Crop Protection* 15(8): 779-788. {a} Western Cotton Res. Lab., USDA-ARS, 4135 E. Broadway Rd., Phoenix, AZ 85040, USA
- Narkhede, W. N., G. M. Bharad, et al. (1996). Effect of plant density and drip irrigation on pre-monsoon cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 41(1): 127-131. Cotton Res. Station, Degloor Rd., Nanded, Maharashtra 431 604, India

- Nasira, K. and M. A. Maqbool (1996). Description of *Tylenchorhynchus gossypii* N.SP. (Nematoda: Dolichodoridae) with observation on *T. elegans* Siddiqi, 1961 from Pakistan. *Pakistan Journal of Nematology* 14(1): 33-40. National Nematological Res. Cent., Univ. Karachi, Karachi 75270, Pakistan
- Nateshan, H. M., V. Muniyappa, et al. (1996). Host range, vector relations and serological relationships of cotton leaf curl virus from southern India. *Annals of Applied Biology* 128(2): 233-244. {a} Dep. Plant Pathol., Univ. Agric. Sci., Hebbal, Bangalore 560 024, India
- Oliveira, E. D., J. D. C. Machado, et al. (1996). Occurrence and survival of *Botryodiplodia theobromae* Pat. in cotton seeds (*Gossypium hirsutum* L.) produced in the state of Minas Gerais, Brazil. *Ciencia e Agrotecnologia* 20(1): 31-38. {a} CNPMS/EMBRAPA, Sete Lagos, MG, Brazil
- Palomo, G. A. (1996). Distribution, collection and use of wild cotton species in Mexico. *Ciencia Mexico City* 47(4): 359-369.
- Patil, D. B., K. T. Naphade, et al. (1996). Effect of nitrogen and phosphorus levels on gossypol and oil content of cotton genotypes (*Gossypium hirsutum* L.). *Agricultural Science Digest* 16(3-4): 163-166. Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola-444 104, India
- Pear, J. R., Y. Kawagoe, et al. (1996). Higher plants contain homologs of the bacterial *celA* genes encoding the catalytic subunit of cellulose synthase. *Proceedings of the National Academy of Sciences of the United States of America* 93(22): 12637-12642. {a} Calgene Inc., 1920 Fifth St., Davis, CA 95616, USA
- Perumal, N. K. (1996). Shedding of fruiting parts in relation to plant traits and yield in rainfed cotton (*Gossypium hirsutum* L.). *Indian Journal of Plant Physiology* 1(3): 218-219. Central Inst. Cotton Reserch, Nagpur 440001, India
- Pettigrew, W. T., J. J. Heitholt, et al. (1996). Genotypic interactions with potassium and nitrogen in cotton of varied maturity. *Agronomy Journal* 88(1): 89-93. {a} USDA-ARS, Cotton Physiol. Genetics Res., P.O. Box 345, Stoneville, MS 38776, USA
- Pizzinatto, M. A. and S. S. Freitas (1996). Effect of fluorescent *Pseudomonas* spp. on seed health and germination and on cotton plant growth. *Summa Phytopathologica* 22(1): 9-14. {a} Inst. Agron., Secao Fitopatol., Caixa Postal 28, Campinas, SP. 13.020-902, Brazil
- Porcelli, F. and F. A. Palmieri (1996). *Oxycareus hyalinipennis* (Costa 1838) (Rhynchota, Lygaeidae) on Kenaf in Basilicata. *Entomologica Bari* 30: 197-205. Dip. Innovazione, Settore Biotecnol. Agric., Div. Agric. Ecocompatibile, Centro Ricerche Energia Trisaia, Rotondella, Italy
- Porter, P. M., M. J. Sullivan, et al. (1996). Cotton cultivar response to planting date on the Southeastern Coastal Plain. *Journal of Production Agriculture* 9(2): 223-227. {a} Dep. Entomol., Edisto REC, Clemson Univ., P.O. Box 247, Blackville, SC 29817, USA
- Press, C. M., W. F. Mahaffee, et al. (1996). Organic by-product effects on soil chemical properties and microbial communities. *Compost Science and Utilization* 4(2): 70-80. {a} Dep. Plant Pathol., Biol. Control Inst., Auburn Univ., Auburn, AL, USA
- Qamar, F., S. A. Khan, et al. (1996). Comparative studies of different nematicides against plant parasitic nematodes in cotton (*Gossypium hirsutum* L.) in Pakistan. *Pakistan Journal of Scientific and Industrial Research* 39(5-8): 144-145. PCSIR Lab. Complex, Off University Road, Karachi 75280, Pakistan
- Quisenberry, J. E. and B. L. McMichael (1996). Screening cotton germplasm for root growth potential. *Environmental and Experimental Botany* 36(3): 333-337. {a} USDA-ARS, Cropping Systems Res. Lab., Plant Stress Water Conservation Res. Unit, Route 3, Box 215, Lubbock, TX 79401, USA
- Rajasegar, G., S. R. S. Rangasamy, et al. (1996). Callus induction, somatic embryoid formation and plant regeneration in cotton (*Gossypium hirsutum* L.). *Journal of Phytological Research* 9(2): 145-147. {a} Dep. Botany, National Univ. Singapore, Singapore-119260, Singapore

- Rajasekaran, K. (1996). Regeneration of plants from cryopreserved embryogenic cell suspension and callus cultures of cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 15(11): 859-864. J.G. Boswell Company, 850 Plymouth Avenue, P.O. Box 787, Corcoran, CA 93212-0787, USA
- Rajasekaran, K., J. W. Grula, et al. (1996). Selection and characterization of mutant cotton (*Gossypium hirsutum* L.) cell lines resistant to sulfonylurea and imidazolinone herbicides. *Plant Science Shannon* 119(1-2): 115-124. {a} 608 Woodbluff Avenue, Duarte, CA 91010-1462, USA
- Rajasekaran, K., J. W. Grula, et al. (1996). Herbicide-resistant Acala and Coker cottons transformed with a native gene encoding mutant forms of acetohydroxyacid synthase. *Molecular Breeding* 2(4): 307-319. {a} USDA, ARS, SRRC, 1100 Robert E. Lee Blvd., New Orleans, LA 70124-4305, USA
- Raveendran, T. S., N. Shanthi, et al. (1996). Combining ability analysis in cotton (*Gossypium* species) hybrids based on cytoplasmic-genic male-sterility system. *Indian Journal of Agricultural Sciences* 66(2): 101-104. Dep. Cotton, Sch. Genet., Tamil Nadu Agric. Univ., Coimbatore 641 003, India
- Reddy, A. R., K. R. Reddy, et al. (1996). Mepiquat chloride (PIX)-induced changes in photosynthesis and growth of cotton. *Plant Growth Regulation* 20(3): 179-183. {a} Sch. Life Sci., Pondicherry Univ., Pondicherry 605 014, India
- Reyes, V. M. H., Y. Ji, et al. (1996). ISH-facilitated analysis of meiotic bivalent pairing. *Genome* 39(4): 784-792. {a} Dep. Soil Crop Sci., Texas A M Univ., College Station, TX 77843, USA
- Riegel, C., F. A. Fernandez, et al. (1996). Meloidogyne incognita infested soil amended with chicken litter. *Journal of Nematology* 28(3): 369-378. Dep. Plant Pathol., Univ. Georgia, Athens, GA 30602-7274, USA
- Rochester, I., G. Constable, et al. (1996). Effective nitrification inhibitors may improve fertilizer recovery in irrigated cotton. *Biology and Fertility of Soils* 23(1): 1-6. {a} CSIRO Div. Plant Ind., Aust. Cotton Res. Inst., Locked Bag 59, Narrabri, NSW 2390, Australia
- Rodriguez, G. D. (1996). The influence of the boron in the retention of the cotton's bolls on the "Marisma del Guadalquivir". *Investigacion Agraria Produccion y Proteccion Vegetales* 11(1): 59-67. CIDA "Las Torres", Apdo. de Correos 41200, Alcala del Rio, Sevilla, Spain
- Rogers, G. S., P. J. Milham, et al. (1996). Interactions between rising CO₂ concentration and nitrogen supply in cotton. I. Growth and leaf nitrogen concentration. *Australian Journal of Plant Physiology* 23(2): 119-125. {a} Sch. Horticulture, Univ. Western Sydney, Hawkesbury, Locked Bag No. 1, PO Richmond, NSW 2753, Australia
- Rogers, J. B., D. S. Murray, et al. (1996). Ivyleaf morningglory (*Ipomoea hederacea*) interference with cotton (*Gossypium hirsutum*). *Weed Technology* 10(1): 107-114. {a} DowElanco, P.O. Box 205, Lemont, PA 16851, USA
- Rose, U. S. R., A. Manukian, et al. (1996). Volatile semiochemicals released from undamaged cotton leaves. *Plant Physiology Rockville* 111(2): 487-495. {a} Insect Attractants, Behavior Basic Biol. Res. Lab., United States Dep. Agriculture, Agricultural Res. Serv., 1700 Southwest 23rd Drive, Gainesville, FL 32604, USA
- Sachs, E. S., J. H. Benedict, et al. (1996). Pyramiding CryIA(b) insecticidal protein and terpenoids in cotton to resist tobacco budworm (Lepidoptera: Noctuidae). *Environmental Entomology* 25(6): 1257-1266. {a} Monsanto Co., Build. BB3J, 700 Chesterfield Village Parkway, St. Louis, MO 63198, USA
- Sadras, V. O. (1996). Cotton responses to simulated insect damage: Radiation-use efficiency, canopy architecture and leaf nitrogen content as affected by loss of reproductive organs. *Field Crops Research* 48(2-3): 199-208. CSIRO Div. Plant Industry, Cotton Res. Unit, PO Box 59, Narrabri 2390, NSW, Australia
- Sadras, V. O. and L. J. Wilson (1996). Effects of timing and intensity of spider mite infestation on the oil yield of cotton crops. *Australian Journal of Experimental Agriculture* 36(5): 577-580. CSIRO, Div. Plant Ind., Cotton Res. Unit, Locked Bag 59, Narrabri, NSW 2390, Australia

Samarakoon, A. B. and R. M. Gifford (1996). Water use and growth of cotton in response to elevated CO₂ in wet and drying soil. *Australian Journal of Plant Physiology* 23(1): 63-74. {a} CSIRO, Div. Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia

Sandhu, B. S. (1996). Opportunities and challenges in hybrid cotton. *Crop Improvement* 23(1): 135-138. Dep. Plant Breeding, Punjab Agric. Univ., Ludhiana-141004, India

Sassenrath, C. G. F., G. Lu, et al. (1996). Photon flux density versus leaf senescence in determining photosynthetic efficiency and capacity of *Gossypium hirsutum* L. leaves. *Environmental and Experimental Botany* 36(4): 439-446. {a} USDA-ARS, Crop Simulation Res. Unit, PO Box 5367, Mississippi State, MS 39762-5367, USA

Savage, M. J., J. T. Ritchie, et al. (1996). Lower limit of soil water availability. *Agronomy Journal* 88(4): 644-651. {a} Dep. Agronomy, Univ. Natal, Pietermaritzburg 3201, South Africa

Sayal, O. U. and M. Z. Sulemani (1996). Comparison of gene action controlling the qualitative traits in some early maturing cultivars of American cotton *Gossypium hirsutum* L. *Sarhad Journal of Agriculture* 12(6): 653-661. {a} Dep. Plant Breeding Genet., Fac. Agric. Gomal Univ., N.W.F.P., Dera Ismail Khan, Pakistan

Sayal, O. U. and M. Z. Sulemani (1996). Comparison of gene action controlling the qualitative traits in some early maturing cultivars of American cotton *Gossypium hirsutum* L. *Sarhad Journal of Agriculture* 12(2): 137-145. {a} Dep. Plant Breed. Genet., Fac. Agric., Gomal Univ., Dera Ismail Khan, Pakistan

Schmutz, A., A. J. Buchala, et al. (1996). Changing the dimensions of suberin lamellae of green cotton fibers with a specific inhibitor of the endoplasmic reticulum-associated fatty acid elongases. *Plant Physiology* Rockville 110(2): 403-411. {a} Inst. Botanische Biol., Univ. Freiburg, A. Gockelstrasse 3, CH-1700 Freiburg, Switzerland

Scott, W. P., G. L. Snodgrass, et al. (1996). Impact of insecticides applied with/without bromoxynil herbicide on various cotton pests in laboratory bioassays. *Journal of Entomological Science* 31(4): 365-370. {a} USDA, ARS, South. Insect Manage. Lab., Stoneville, MS 38776, USA

Seifert, S. and C. E. Snipes (1996). Influence of flame cultivation on mortality of cotton (*Gossypium hirsutum*) pests and beneficial insects. *Weed Technology* 10(3): 544-549. Fac. Hort. Agric., Humboldt Univ., Berlin, Germany

Setatou, H. B. and A. D. Simonis (1996). Effect of time and rate of nitrogen application on cotton. *Fertilizer Research* 43(1-3): 49-53. {a} Soil Sciences Institute, 541 10 Thessaloniki, Greece

Shankle, M. W., R. M. Hayes, et al. (1996). MSMA and pyriithiobac effects on cotton (*Gossypium hirsutum*) development, yield and quality. *Weed Science* 44(1): 137-142. {a} P.O. Box 1071, Knoxville, TN 37901, USA

Shappley, Z. W., J. N. Jenkins, et al. (1996). Establishment of molecular markers and linkage groups in two F-2 populations of upland cotton. *Theoretical and Applied Genetics* 92(8): 915-919. {a} USDA, ARS, CSRL, PO Box 5367, Mississippi State, MS 39762, USA

Shaver, T. N. and J. D. Lopez, Jr. (1996). Ovipositional response of the tobacco budworm and bollworm to chickpea. *Southwestern Entomologist* 21(3): 277-282. Southern Crops Res. Lab., USDA-ARS Areawide Pest Management Res. Unit, College Station, TX 77845, USA

Shaw, D. R. (1996). Development of stale seedbed weed control programs for southern row crops. *Weed Science* 44(2): 413-416. Dep. Plant Soil Sci., Miss. State Univ., Mississippi State, MS 39762, USA

Shepherd, R. L., J. C. McCarty, Jr., et al. (1996). Registration of nine cotton germplasm lines resistant to root-knot nematode. *Crop Science* 36(3): 820. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

Shishehbor, P. and P. A. Brennan (1996). Life history traits of castor whitefly, *Trialeurodes ricini misra* (Hom., Aleyrodidae), on eight host plant species. *Journal of Applied Entomology* 120(9): 519-522. {a} Dep. Environ. Resour. Management, Univ. Coll. Dublin, Belfield, Dublin 4, Ireland

Siddiqui, M. A. (1996). A study of variability and heritability of some quantitative character in *hirsutum* cotton. *Journal of Maharashtra Agricultural Universities* 21(2): 256-258. {a} Marathwada Agric. Univ., Parbhani 431 402, India

Silva, F. P. D., F. I. O. Melo, et al. (1996). Okra-leaf and normal cotton isolines with different backgrounds. *Brazilian Journal of Genetics* 19(3): 453-457. {a} Departamento de Fitotecnia, Centro de Ciencias Agrarias, Universidade Federal do Ceara, Caixa Postal 12168, 60451-970 Fortaleza, CE, Brazil

Silva, J. B., J. A. R. Matos, et al. (1996). The effect of cotton seed bacterization on the control of *Rhizoctonia solani*. *Fitopatologia Brasileira* 21(3): 342-348. Dep. Agron.-Area Fitossanidade, Univ. Federal Rural de Pernambuco, 52171-900 Recife, PE, Brazil

Sims, S. R., S. A. Berberich, et al. (1996). Analysis of expressed proteins in fiber fractions from insect-protected and glyphosate-tolerant cotton varieties. *Crop Science* 36(5): 1212-1216. {a} CREEGEN, Unit Monsanto Co., 700 Chesterfield Parkway North, Chesterfield, MO 63198, USA

Simwat, G. S. and A. K. Dhawan (1996). Efficacy of dimethoate for controlling sucking pests of upland cotton (*Gossypium hirsutum*) under irrigation. *Indian Journal of Agricultural Sciences* 66(4): 263-264. Punjab Agricultural Univ., Ludhiana 141 004, India

Singh, I. and B. L. Nagda (1996). Effect of sowing date, spacing and nitrogen level on upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 66(5): 293-294. {a} Dep. Agron., Agricultural Research Substation, Rajasthan Agriculture University, Sumerpur 306 902, India

Singh, S. B., D. Singh, et al. (1996). Variation in physio-morphological characters related to drought tolerance in cotton (*Gossypium* species). *Indian Journal of Agricultural Sciences* 66(6): 357-359. {a} Central Inst. Cotton Res., Nagpur, Maharashtra 440 001, India

Skinner, R. H. (1996). Leaf temperature effects on *Bemisia argentifolii* (Homoptera: Aleyrodidae) oviposition. *Environmental Entomology* 25(6): 1371-1375. Great Plains Systems Research, USDA-ARS, Fort Collins, CO 80522, USA

Skinner, R. H. (1996). Response of *Bemisia argentifolii* (Homoptera: Aleyrodidae) to water and nutrient stressed cotton. *Environmental Entomology* 25(2): 401-406. Western Cotton Res. Lab., USDA-ARS, Phoenix, AZ 85040, USA

Snipes, C. E. and R. L. Allen (1996). Interaction of graminicides applied in combination with pyriithiobac. *Weed Technology* 10(4): 889-892. Mississippi Agric. For. Exp. Stn., Delta Branch, Stoneville, MS 38776, USA

Snodgrass, G. L. (1996). Glass-vial bioassay to estimate insecticide resistance in adult tarnished plant bugs (Heteroptera: Miridae). *Journal of Economic Entomology* 89(5): 1053-1059. South. Insect Manage. Lab., USDA-ARS, P.O. Box 346, Stoneville, MS 38776, USA

Staggenborg, S. A., R. J. Lascano, et al. (1996). Determining cotton water use in a semiarid climate with the GOSSYM cotton simulation model. *Agronomy Journal* 88(5): 740-745. {a} Northeast Area Ext. Office, Kansas State Univ., Manhattan, KS 66502, USA

Stephenson, P., B. A. Collins, et al. (1996). Localization of ubiquitin to differentiating vascular tissues. *American Journal of Botany* 83(2): 140-147. {a} Dep. Biol. Sci., Smith Coll., Northampton, MA 01063, USA

- Stevens, W. E., J. J. Varco, et al. (1996). Evaluating cotton nitrogen dynamics in the GOSSYM simulation model. *Agronomy Journal* 88(2): 127-132. {a} Univ. Mo.-Delta Cent., P.O. Box 160, Portageville, MO 63873, USA
- Tanaka, M. A. D. S., J. O. M. Menten, et al. (1996). Growth habit of *Colletotrichum gossypii* and *C. gossypii* var. *cephalosporioides* on cotton seeds. *Bragantia* 55(1): 95-104. {a} Secao Fitopatol., Inst. Agron., Caixa Postal 28, 13001-970 Campinas, SP, Brazil
- Tang, B., J. N. Jenkins, et al. (1996). Evaluation of genetic variances, heritabilities, and correlations for yield and fiber traits among cotton F-2 hybrid populations. *Euphytica* 91(3): 315-322. {a} Dep. Plant Soil Sci., Mississippi State Univ., Mississippi State, MS 39762, USA
- Tatineni, V., R. G. Cantrell, et al. (1996). Genetic diversity in elite cotton germplasm determined by morphological characteristics and RAPDs. *Crop Science* 36(1): 186-192. {a} Depl Agron. Hortic., N.M. State Univ., La Cruces, NM 88033-8003, USA
- Thaker, V. S., S. Saroop, et al. (1996). Physiological and biochemical changes associated with cotton fibre development: VI. Acid and alkaline phosphatases. *Acta Physiologiae Plantarum* 18(2): 111-116. {a} Dep. Biosci., Saurashtra Univ., Rajkot 360 005, India
- Thimmaiah, K. K., G. Robbani, et al. (1996). Role of private sector in crop improvement. *Crop Improvement* 23(Special): 99-111. Mahenbdra Hybrid Seeds Co. Ltd., A-10/11, Old MIDC Area, Jalna 431 203, MS, India
- Tort, N. (1996). Effects of light, different growth media, temperature and salt concentrations on germination of cotton seeds (*Gossypium hirsutum* L. cv. Nazilli 87). *Journal of Agronomy and Crop Science* 176(4): 217-221. Ege Univ. Fen Fak., Botanik Anabilim Dali, Bornova-Izmir, Turkey
- Trainin, T., M. Shmuel, et al. (1996). In vitro prenylation of the small GTPase Rac13 on cotton. *Plant Physiology Rockville* 112(4): 1491-1497. {a} Dep. Plant Sci., Inst. Life Sci., Hebrew Univ., Jerusalem 91904, Israel
- Trelease, R. N., M. S. Lee, et al. (1996). C-terminal polypeptides are necessary and sufficient for in vivo targeting of transiently-expressed proteins to peroxisomes in suspension-cultured plant cells. *Protoplasma* 195(1-4): 156-167. {a} Dep. Botany, Arizona State Univ., Box 871601, Tempe, AZ 85287-1601, USA
- Triplett, G. B., Jr., S. M. Dabney, et al. (1996). Tillage systems for cotton on silty upland soils. *Agronomy Journal* 88(4): 507-512. {a} Dep. Plant Soil Sci., Mississippi State Univ., Box 9555, Mississippi State, MS 39762, USA
- Turley, R. B. and D. L. Ferguson (1996). Changes of ovule proteins during early fiber development in a normal and a fiberless line of cotton (*Gossypium hirsutum* L.). *Journal of Plant Physiology* 149(6): 695-702. United States Dep. Agriculture, Agricultural Res. Serv., Cotton Physiology Genetics Res., P.O. Box 345, Stoneville, MS 38776, USA
- Unruh, B. L. and J. C. Silvertooth (1996). Comparisons between an Upland and a Pima cotton cultivar: II. Nutrient uptake and partitioning. *Agronomy Journal* 88(4): 589-595. {a} Dep. Plant Sci, Univ. Arizona, Tucson, AZ 85721, USA
- Unruh, B. L. and J. C. Silvertooth (1996). Comparisons between an Upland and a Pima cotton cultivar: I. Growth and yield. *Agronomy Journal* 88(4): 583-589. {a} Dep. Plant Sci., Univ. Arizona, Tucson, AZ 85721, USA
- Van, I. M. W. and D. M. Oosterhuis (1996). Drought effects on the water relations of cotton fruits, bracts, and leaves during ontogeny. *Environmental and Experimental Botany* 36(1): 51-59. {a} Dep. Horticulture, Univ. Georgia, Georgia Experiment Station, Griffin, GA 30223, USA

Venzon, M. and F. A. Pallini (1996). Occurrence of *Taedia stigmosa* (Berg.) (Hemiptera: Miridae) in cotton fields in "Triangulo Mineiro", State of Minas Gerais, Brazil (short communication). *Cientifica Jaboticabal* 24(1): 225-227. {a} Dep. Entomol., EPAMIG, Caixa Postal 351, 38001-970 Uberaba, MG, Brazil

Viereck, R., M. Kirsch, et al. (1996). Down-regulation of plant V-type H⁺-ATPase genes after light-induced inhibition of growth. *FEBS Letters* 384(3): 285-288. {a} Bot. Inst., Ruprecht-Karls-Univ., Im Neuenheimer Feld 360, D-69120 Heidelberg, Germany

Vigil, E. L., W. S. Anthony, et al. (1996). Fine structural aspects of cotton fiber attachment to the seed coat: Morphological factors affecting saw ginning of lint cotton. *International Journal of Plant Sciences* 157(1): 92-102. {a} USDA-ARS, Climate Stress Lab. B-046A, Beltsville, MD 20705, USA

Vigil, E. L., A. L. Fleming, et al. (1996). Comparative cytological and biochemical analysis of protein storage vacuoles from cotyledons and radicles of cotton seeds. *Seed Science Research* 6(1): 31-37. {a} Climate Stress Lab., U.S. Dep. Agric., ARS, Beltsville, MD 20705, USA

Volokita, M., A. Abeliovich, et al. (1996). Denitrification of groundwater using cotton as energy source. *Water Science and Technology* 34(1-2): 379-385. Lab. Applied Environ. Microbiol., J. Blaustein Inst. Desert Res., Ben-Gurion Univ. Negev, Sde Boker 94993, Israel

Vroh, B. I., L. Harvengt, et al. (1996). Improved RAPD amplification of recalcitrant plant DNA by the use of activated charcoal during DNA extraction. *Plant Breeding* 115(3): 205-206. {a} Unite Biol. Vegetale, Fac. Sciences Agronomiques, B-5030 Gembloux, Belgium

Waldrop, M. P., T. M. Sterling, et al. (1996). Fate of prometryn in prometryn-tolerant and -susceptible cotton cultivars. *Pesticide Biochemistry and Physiology* 56(2): 111-122. {a} Dep. Entomol. Plant Pathol. Weed Sci., N. M. State Univ., Las Cruces, NM 88003, USA

Wanjura, D. F., J. R. Mahan, et al. (1996). Irrigation starting time effects on cotton under high-frequency irrigation. *Agronomy Journal* 88(4): 561-566. {a} USDA-ARS, Cropping Systems Res. Lab., Route 3, Box 215, Lubbock, TX 79401, USA

Webster, E. P. and D. R. Shaw (1996). Carryover potential of pyriithiobac to rotational crops on a Mississippi Black Belt region clay soil. *Weed Technology* 10(1): 140-144. {a} Dep. Plant Soil Sciences, Mississippi State Univ., Mississippi State, MS 39762, USA

Wei, Y. A., D. L. Hendrix, et al. (1996). Isolation of a novel tetrasaccharide, bemisiotetrose, and glycine betaine from silverleaf whitefly honeydew. *Journal of Agricultural and Food Chemistry* 44(10): 3214-3218. {a} Western Cotton Res. Lab., ARS, U.S. Dep. Agric., 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

Willis, T. M. and A. S. Black (1996). Irrigation increases groundwater recharge in the Macquarie Valley. *Australian Journal of Soil Research* 34(6): 837-847. Sch. Sci. Agric., Charles Sturt University, PO Box 588, Wagga Wagga, NSW 2678, Australia

Wool, D. and D. F. Hales (1996). Previous infestation affects recolonization of cotton by *Aphis gossypii*: Induced Resistance or plant damage? *Phytoparasitica* 24(1): 39-48. {a} Dep. Zool., George S. Wise Fac. Life Sci., Tel-Aviv Univ., Tel Aviv 69978, Israel

Yadav, K. S. and R. L. Rajput (1996). Effect of hybrid, spacing and nitrogen levels on growth and yield of upland cotton (*Gossypium hirsutum*). *Agricultural Science Digest* 16(1): 49-51. JNKVV-Zonal Agric. Res. Stn., Morena-476 001, MP, India

Yee, W. L., N. C. Toscano, et al. (1996). *Bemisia argentifolii* (Homoptera: Aleyrodidae) action thresholds and cotton photosynthesis. *Environmental Entomology* 25(6): 1267-1273. {a} Dep. Entomol., Univ. California, Riverside, CA 92521, USA

- Zeringue, H. J., Jr. (1996). Possible involvement of lipoxygenase in a defense response in aflatoxigenic *Aspergillus*-cotton plant interactions. *Canadian Journal of Botany* 74(1): 98-102. USDA, Agric. Res. Serv., Southern Regional Res. Center, P.O. Box 19687, New Orleans, LA 70179-0687, USA
- Zhang, B. H. L. X. L., F. L. Li, et al. (1996). Development of abnormal plantlets and their normalization during cotton tissue culture. *Acta Botanica Sinica* 38(11): 845-852. Cotton Res. Inst., Chinese Acad. Agric. Sci., Anyang, Henan 455112, China
- Zhang, Z., M. L. Pierce, et al. (1996). Detection and differentiation of pectic enzyme activity in vitro and in vivo by capillary electrophoresis of products from fluorescent-labeled substrate. *Electrophoresis* 17(2): 372-378. {a} Dep. Biochem. Mol. Biol., Oklahoma State Univ., Stillwater, OK 74078-3035, USA
- Abbas, H. K., T. Tanaka, et al. (1995). Susceptibility of various crop and weed species to AAL-toxin, a natural herbicide. *Weed Technology* 9(1): 125-130. {a} U.S. Dep. Agric., Agric. Res. Serv., South Weed Sci. Lab., Stoneville, MS 38776, USA
- Abdurakhmanova, Z. N., A. Abdullaev, et al. (1995). Photosynthesis and activity of carboxylating enzymes in the cotton leaves of different layers. *Fiziologiya Rastenii Moscow* 42(2): 201-205. {a} Inst. Plant Physiol. Biophys., Taj. Acad. Sci., ul. S. Aini 299/2, 734063 Dushanbe, Tajikistan
- Ahmad, M., A. Rauf, et al. (1995). Studies on salt tolerance of cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research* 29(1-2): 64-68. {a} Central Cotton Res. Inst., Multan, Pakistan
- Allen, R. L. and C. E. Snipes (1995). Interactions of foliar insecticides applied with pyriithiobac. *Weed Technology* 9(3): 512-517. Miss. Agric. For. Exp. Stn., Delta Branch, Stoneville, MS 38776, USA
- Allen, S. J. (1995). Plant quarantine and diseases of cotton. *Australasian Plant Pathology* 24(1): 70-73. N.S.W. Agric., Aust. Cotton Res. Inst., Narrabri, NSW 2390, Australia
- Amor, Y., C. H. Haigler, et al. (1995). A membrane-associated form of sucrose synthase and its potential role in synthesis of cellulose and callose in plants. *Proceedings of the National Academy of Sciences of the United States of America* 92(20): 9353-9357. {a} Dep. Botany, Inst. Life Sci., Hebrew Univ. Jerusalem, Jerusalem 91904, Israel
- Anand, S. C., S. R. Koenning, et al. (1995). Effect of temporal deployment of different sources of resistance to soybean cyst nematode. *Journal of Production Agriculture* 8(1): 119-123. {a} Dep. Agron., Univ. Missouri Delta Cent., Portageville, MO 63873, USA
- Arshad, M., A. Hussain, et al. (1995). Effect of soil applied L-tryptophan on growth and chemical composition of cotton. *Journal of Plant Nutrition* 18(2): 317-329. {a} Dep. Soil Environ. Sci., Univ. Calif., Riverside, CA 92521-0424, USA
- Babu, R., M. V. H. Rao, et al. (1995). Effect of chemical defoliant on earliness, seed-cotton yield and quality of upland cotton (*Gossypium hirsutum*) under irrigated condition. *Indian Journal of Agronomy* 40(1): 157-159. Agric. Res. Stn., Univ. Agric. Sci., Arabhavi 591 318, India
- Baird, R. E., T. B. Brenneman, et al. (1995). First report of *Rhizoctonia* sp. CAG-5 on cotton in Georgia. *Plant Disease* 79(3): 320. {a} Plant Pathol. Dep., Tifton, GA 31794, USA
- Baird, R. E., T. B. Brenneman, et al. (1995). Influence of crop rotation and flutolanil on the diversity of fungi on peanut shells. *Phytoprotection* 76(3): 101-113. {a} Plant Pathol. Dep., Univ. Georgia, Rural Dev. Cent., P.O. Box 1209, Tifton, GA 31793, USA
- Baird, R. E. and D. E. Carling (1995). First report of *Rhizoctonia solani* AG-7 in Indiana. *Plant Disease* 79(3): 321. {a} Pathol. Dep., Univ. Georgia, RDC, Tifton, GA 31793, USA

Baker, R. J., J. L. Longmire, et al. (1995). Organization of repetitive elements in the upland cotton genome (*Gossypium hirsutum*). *Journal of Heredity* 86(3): 178-185. {a} Dep. Biol. Sci., Texas Tech Univ., Lubbock, TX 79409, USA

Baloch, M. J., H. Bhutto, et al. (1995). Combining ability estimates in 5X5 diallel intrahirsutum crosses. *Pakistan Journal of Botany* 27(1): 121-126. {a} Cotton Res. Inst., Sakrand, Sindh, Pakistan

Banerjee, R. N. and T. K. Paul (1995). Malvaceae of west Dinajpur District, West Bengal. *Journal of Economic and Taxonomic Botany* 19(2): 313-316. Bot. Survey India, Howrah-711 103, India

Barrett, D. J. and R. M. Gifford (1995). Acclimation of photosynthesis and growth by cotton to elevated CO₂: Interactions with severe phosphate deficiency and restricted rooting volume. *Australian Journal of Plant Physiology* 22(6): 955-963. {a} CSIRO Div. Plant Ind., GPO Box 1600, Canberra, ACT 2601, Australia

Bednarz, C. W. and D. M. Oosterhuis (1995). Partitioning of potassium during the development of deficiency symptoms in cotton. *Arkansas Agricultural Experiment Station Research Series*(443): 91-95. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA

Bejarano, A. J., V. J. M. Melero, et al. (1995). Influence of inoculum density of defoliating and nondefoliating pathotypes of *Verticillium dahliae* on epidemics of verticillium wilt of cotton in southern Spain. *Phytopathology* 85(12): 1474-1481. {a} Inst. Agric. Sostenible, CSIC, Apdo. 4084, 14080 Cordoba, Spain

Bhatade, S. S., K. G. Nandanwankar, et al. (1995). Phenotypic stability of newly developed genotypes of upland cotton (*Gossypium hirsutum*) in Marathwada region of Maharashtra. *Indian Journal of Agricultural Sciences* 65(4): 295-297. Cotton Res. Stn., Marathwada Agric. Univ., Nanded, Maharashtra 431 604, India

Bhatnagar, G. S. and P. C. Chaplot (1995). Production potential and economics of different crop sequences in Mahi-command area of south-east Rajasthan. *Indian Journal of Agricultural Sciences* 65(4): 279-280. {a} Agric. Res. Stn., Rajasthan Agric. Univ., Banswara 327 001, India

Bhuva, K. S., N. M. Sukhadia, et al. (1995). Intercropping in upland cotton (*Gossypium hirsutum*) with pulse and oilseed crops under rainfed condition. *Indian Journal of Agronomy* 40(1): 95-97. Dep. Agronomy, Gujart Agric. Univ., Campus Junagadh 362 001, India

Blanchard, J. L. and G. W. Schmidt (1995). Pervasive migration of organellar DNA to the nucleus in plants. *Journal of Molecular Evolution* 41(4): 397-406. {a} Dep. Botany, Univ. Georgia, Athens, GA 30602, USA

Bloodworth, L. H. and J. R. Johnson (1995). Cover crops and tillage effects of cotton. *Journal of Production Agriculture* 8(1): 107-112. {a} USDA-SCS Jamie L. Whitten Plant Materials Cent., Coffeetown, MS 38922, USA

Bondada, B. R., D. M. Oosterhuis, et al. (1995). Cotton leaf growth and foliar nitrogen absorption. *Arkansas Agricultural Experiment Station Research Series*(443): 67-71. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA

Bowman, D. T. (1995). Registration of three glabrous cotton germplasm lines with high lint percentage: NC 88-90, NC 88-91, and NC 88-95. *Crop Science* 35(2): 600. Crop Sci. Dep., Box 8604, N.C. State Univ., Raleigh, NC 27695-8604, USA

Bunkelmann, J., F. J. Corpas, et al. (1995). Four putative, glyoxysome membrane proteins are instead immunologically-related protein body membrane proteins. *Plant Science Limerick* 106(2): 215-226. {a} Dep. Botany, Arizona State Univ., Tempe, AZ 85287-1601, USA

Burke, J. J. and D. R. Upchurch (1995). Cotton rooting patterns in relation to soil temperatures and the thermal kinetic window. *Agronomy Journal* 87(6): 1210-1216. {a} USDA-ARS Cropping Systems Res. Lab., Route 3, Box 215, Lubbock, TX 79401, USA

- Cantrell, R. G., C. L. Roberts, et al. (1995). Registration of 'Acala 1517-95' cotton. *Crop Science* 35(4): 1227-1228. {a} Dep. Agron. Hortic., New Mexico State Univ., Las Cruces, NM 88003-0003, USA
- Carvalho, L. P. D., C. D. Cruz, et al. (1995). Diallel analysis of yield and other traits in cotton. *Revista Brasileira de Genetica* 18(1): 93-97. {a} Rua Osvaldo Cruz 1143, Caixa Postal 174, 58107-720 Campina Grande, PB, Brazil
- Chapman, K. D., I. Lin, et al. (1995). Metabolism of Cottonseed Microsomal N-Acylphosphatidylethanolamine. *Archives of Biochemistry and Biophysics* 318(2): 401-407. {a} Dep. Biol. Sci., Div. Biochem. Mol. Biol., Univ. N. Texas, Denton, TX 76203, USA
- Charles, G. W. (1995). Nutgrass (*Cyperus rotundus* L.) control in cotton (*Gossypium hirsutum* L.). *Australian Journal of Experimental Agriculture* 35(5): 633-639. NSW Agric., Aust. Cotton Res. Inst., Narrabri, NSW 2390, Australia
- Chu, C. C., T. J. Henneberry, et al. (1995). *Bemesia argentifolii* (Homoptera: Aleyrodidae): Host preference and factors affecting oviposition and feeding site preference. *Environmental Entomology* 24(2): 354-360. {a} Western Cotton Res. Lab., USDA-ARS, 4151 Highway 86, Brawley, CA 92227, USA
- Chu, C. C., T. J. Henneberry, et al. (1995). Effect of irrigation frequency on cotton yield in short-season production systems. *Crop Science* 35(4): 1069-1073. {a} USDA-ARS Western Cotton Res. Lab., Brawley, CA 92227, USA
- Cohen, Y., Z. Plaut, et al. (1995). Deficit irrigation of cotton for increasing groundwater use in clay soils. *Agronomy Journal* 87(5): 808-814. {a} Inst. Soils and Water, Agric. Res. Organization, Volcani Cent., Bet Dagan 50250, Israel
- Cook, C. G. and L. N. Namken (1995). Registration B411, B416, and B417 parental lines of cotton. *Crop Science* 35(5): 1518. {a} USDA-ARS, Subtrop. Agric. Res. Lab., 2413 E. Highway 83, Weslaco, TX 78596, USA
- Cook, C. G. and A. W. Scott, Jr. (1995). Registration of C21781-2 cotton germplasm. *Crop Science* 35(2): 599. {a} USDA-ARS, Subtrop. Agric. Res. Lab., 2413 E. Hwy. 83, Weslaco, TX 78596, USA
- Creech, R. G., J. N. Jenkins, et al. (1995). Cotton resistance to root-knot nematode: I. Penetration and reproduction. *Crop Science* 35(2): 365-368. {a} Mississippi State Univ., P.O. Box 9555, Mississippi State, MS 39762, USA
- Daayf, F., M. Nicole, et al. (1995). Differentiation of *Verticillium dahliae* populations on the basis of vegetative compatibility and pathogenicity on cotton. *European Journal of Plant Pathology* 101(1): 69-79. {a} Lab. Phytopathol. Tropicale, ORSTOM, 911, Avenue Agropolis, 34032 Montpellier, France
- Davidonis, G. (1995). Changes in polyamine distribution during cotton fiber and seed development. *Journal of Plant Physiology* 145(1-2): 108-112. USDA, ARS, Southern Regional Res. Cent., P.O. Box 19687, New Orleans, LA 70119, USA
- Davila, H. G., H. Hamada, et al. (1995). Cadinane-type sesquiterpenes induced in *Gossypium* cotyledons by bacterial inoculation. *Phytochemistry Oxford* 39(3): 531-536. {a} Dep. Biochem. Molecular Biol., Okla. State Univ., Stillwater, OK 74089-0454, USA
- Davis, G. D. and M. Essenberg (1995). (+)-delta-cadinene is a product of sesquiterpene cyclase activity in cotton. *Phytochemistry Oxford* 39(3): 553-567. {a} Dep. Biochem. Molecular Biol., Okla. State Univ., Stillwater, OK 740780-0454, USA
- De, C. L. P., C. D. Cruz, et al. (1995). Genetic divergence in Brazilian cotton, *Gossypium hirsutum* var. *latifolium* Hutch. *Revista Brasileira de Genetica* 18(3): 439-443. {a} EMPRAPA, Centro Nacional Pesquisa Algodao, 58107-720 Campina Grande, PB, Brazil

Delgado, J. A. and M. C. Amacher (1995). Modeling the uptake of sulfur by crops on two alluvial soils of Louisiana: Cotton. *Communications in Soil Science and Plant Analysis* 26(11-12): 1921-1934. USDA-ARS, Soil-Plant Nutrient Res. Unit, Ft. Collins, CO 80522, USA

Delmer, D. P., J. R. Pear, et al. (1995). Genes encoding small GTP-binding proteins analogous to mammalian rac are preferentially expressed in developing cotton fibers. *Molecular and General Genetics* 248(1): 43-51. {a} Calgene Inc., 1920 Fifth St., Davis, CA 95616, USA

Echkewu, C. A. and S. O. Alabi (1995). Genetic effects of yield and its components in interspecific crosses of cotton. *Discovery and Innovation* 7(4): 395-399. Inst. Agric. Res., Ahmadu Bello Univ., PMB, 1044 Samaru, Zaria, Nigeria

Eghball, B. and J. F. Power (1995). Fractal description of temporal yield variability of 10 crops in the United States. *Agronomy Journal* 87(2): 152-156. {a} Dep. Agronomy, USDA -ARS, Univ. Nebraska, Lincoln, NE 68583, USA

Essington, M. E., D. D. Tyler, et al. (1995). Fluometuron behavior in long-term tillage plots. *Soil Science* 160(6): 405-414. {a} Dep. Plant Soil Sci., Univ. Tenn., P.O. Box 1071, Knoxville, TN 37901-1071, USA

Farias, F. J. C., M. A. P. Ramalho, et al. (1995). Comparison between stability methods for yield in upland cotton cultivars. *Ciencia e Pratica* 19(3): 252-255. {a} EMBRAPA-CNPA, CP 174, CEP 58107-720, Campina Grande, PB, Brazil

Faye, D. D., W. C. Wang, et al. (1995). Phytoinhibition-phytostimulation potentials of systemic insecticides on tissue-cultured cotton (*Gossypium hirsutum*) cells. *Southwestern Entomologist* 20(2): 151-156. {a} Sphere Corp., Route 4, Box 172, Georgetown, TX 78628, USA

Flint, H. M., T. J. Henneberry, et al. (1995). The effects of transgenic cotton, *Gossypium hirsutum* L., containing *Bacillus thuringiensis* toxin genes for the control of the pink bollworm *Pectinophora gossypiella* (Saunders) and other arthropods. *Southwestern Entomologist* 20(3): 281-292. {a} Western Cotton Res. Lab., USDA, ARS, Phoenix, AZ 85040-8830, USA

Flint, H. M., N. J. Parks, et al. (1995). Tests of garlic oil for control of the silverleaf whitefly, *Bemisia argentifolii* bellows and perring (Homoptera: Aleyrodidae) in cotton. *Southwestern Entomologist* 20(2): 137-150. Western Cotton Res. Lab., USDA-ARS, Phoenix, AZ 85040-8830, USA

Flint, H. M., J. W. Radin, et al. (1995). The effects of drip or furrow irrigation of cotton on *Bemisia argentifolii* (Homoptera: Aleyrodidae). *Journal of Agricultural Entomology* 12(1): 25-32. Western Cotton Res. Lab., Phoenix, AZ 85040, USA

Foy, C. D., R. R. Weil, et al. (1995). Differential manganese tolerances of cotton genotypes in nutrient solution. *Journal of Plant Nutrition* 18(4): 685-706. {a} Climate Stress Lab., Nat. Resource Inst., USDA, ARS, Beltsville, MD 02705-2350, USA

Gorham, J. and J. Bridges (1995). Effects of calcium on growth and leaf ion concentrations of *Gossypium hirsutum* grown in saline hydroponic culture. *Plant and Soil* 176(2): 219-227. Centre Arid Zone Studies, Sch. Biol. Sci., Memorial Build., Univ. Wales, Bangor LL57 2UW, UK

Gruha, J. W., R. L. Hudspeth, et al. (1995). Organization, inheritance and expression of acetohydroxyacid synthase genes in the cotton allotetraploid *Gossypium hirsutum*. *Plant Molecular Biology* 28(5): 837-846. {a} PhytoGen, 101 Waverly Dr., Pasadena, CA 91105, USA

Gururajan, K. N. and S. Henry (1995). Genetics of certain boll characters in cotton (*Gossypium hirsutum* L.). *Indian Journal of Genetics and Plant Breeding* 55(4): 374-378. Regional Station, Central Inst. Cotton Res., Coimbatore, Tamil Nadu, India

Guzhov, Y. L., N. G. Simongulyan, et al. (1995). Heterosis and production of hybrid seeds of cotton on the basis of cytoplasmic male sterility in crosses of *Gossypium hirsutum* L. X *G. barbadense* L. *Izvestiya*

Akademii Nauk Seriya Biologicheskaya Moscow(6): 674-684. {a} Russ. People's Friendship Univ., ul. Miklukho-Maklaya 6, Moscow 117198, Russia

Haeger, K. P., H. Braun, et al. (1995). Evolution of seed storage protein genes: Legumin genes of *Ginkgo biloba*. *Journal of Molecular Evolution* 41(4): 457-466. {a} Dep. Plant Ecol., Univ. Bayreuth, Univ. 30, D-95440 Bayreuth, Germany

Haitas, V. C., S. E. Kotoula, et al. (1995). Influence of propaquizafop application rate and time on *Sorghum halepense* (L.) Pers. control and cotton (*Gossypium hirsutum*) yield. *Weed Research* 35(1): 1-6. {a} Efthimiadis S.A., 57022 Thessaloniki, Plant Protect. Inst., 57001 Thessaloniki, Greece

Han, X. Y. F. (1995). Stylet penetration behaviour of soybean aphid, *Aphis glycines*, on host and non-host plants. *Acta Entomologica Sinica* 38(3): 278-283. Inst. Zool., Academia Sinica, Beijing 100080, China

Hanson, R. E., M. S. Zwick, et al. (1995). Fluorescent in situ hybridization of a bacterial artificial chromosome. *Genome* 38(4): 646-651. {a} Dep. Soil Crop Sciences, Texas A and M Univ., College Station, TX 77843-2474, USA

Harris, J. R., B. J. Gossett, et al. (1995). Growth characteristics of selected dinitroaniline-resistant and -susceptible goosegrass (*Eleusine indica*) population. *Weed Technology* 9(3): 561-567. {a} Ala. Agric. Exp. Stn., Marion Junction, AL 36759, USA

Hasan, H. A. H. (1995). Efficacy of Monceren and Monceren-Combi on mycoflora and improvement of germination of cotton seed. *Dirasat Series B Pure and Applied Sciences* 22(1): 89-95. Botany Dep., Faculty Science, Assiut Univ., Assiut, Egypt

Hasenfratz, M. P., C. L. Tsou, et al. (1995). Expression of two related vacuolar H⁺-ATPase 16-kilodalton proteolipid genes is differentially regulated in a tissue-specific manner. *Plant Physiology Rockville* 108(4): 1395-1404. {a} Dep. Agronomy Range Science, Univ. California, Davis, CA 95616-8515, USA

Hedin, P. A., F. E. Callahan, et al. (1995). Total sterols in root-knot nematode *Meloidogyne incognita* infected cotton *Gossypium hirsutum* (L.) plant roots. *Comparative Biochemistry and Physiology B* 111(3): 447-452. {a} USDA-ARS, PO Box 5367, Mississippi State, MS 39762, USA

Hedin, P. A. and J. C. McCarty, Jr. (1995). Boll weevil *Anthonomus grandis* Boh. oviposition is decreased in cotton *Gossypium hirsutum* L. lines lower in anther monosaccharides and gossypol. *Journal of Agricultural and Food Chemistry* 43(10): 2735-2739. {a} Crop Sci. Res. Lab., Agricultural Res. Serv., U.S. Dep. Agric., Mississippi State, MS 39762, USA

Heitholt, J. J. (1995). Cotton flowering and boll retention in different plant configurations and leaf shapes. *Agronomy Journal* 87(5): 994-998. USDA-ARS, Cotton Physiol. Genetics, P.O. Box 345, Stoneville, MS 38776, USA

Hicks, S. K. and R. J. Lascano (1995). Estimation of leaf area index for cotton canopies using the LI-COR LAI-2000 plant canopy analyzer. *Agronomy Journal* 87(3): 458-464. {a} Texas A and M Univ. Res. Ext. Cent., Rt. 3, Box 219, Lubbock, TX 79401-9757, USA

Howard, D. D. and C. O. Gwathmey (1995). Influence of surfactants on potassium uptake and yield response of cotton to foliar potassium nitrate. *Journal of Plant Nutrition* 18(12): 2669-2680. Plant Soil Sci. Dep., Univ. Tenn., Agric. Experiment Station, West Tenn. Experiment Station, 605 Airways Blvd., Jackson, TN 38301, USA

Jafri, A. Z. and R. Ahmad (1995). Effect of soil salinity on leaf development, stomatal size and its distribution in cotton (*Gossypium hirsutum* L.). *Pakistan Journal of Botany* 27(2): 297-303. {a} Dep. Botany, Islamia Degree Coll., Multan, Pakistan

Jayalalitha, K. and A. Narayanan (1995). The effects of sulphur deficiency on growth and mineral composition of hydroponically grown cotton plants. *Indian Journal of Plant Physiology* 38(4): 309-312. Dep. PlantPhysiol., Agric. Coll., Bapatla -522 101, India

Jenkins, J. N., R. G. Creech, et al. (1995). Cotton resistance to root-knot nematode: II. Post-penetration development. *Crop Science* 35(2): 369-373. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

John, M. E. (1995). Characterization of a cotton (*Gossypium hirsutum* L.) fiber mRNA (Fb-B6). *Plant Physiology Rockville* 107(4): 1477-1478. Agracetus Inc., 8520 University Green, Middleton, WI 53562, USA

John, M. E. and G. Keller (1995). Characterization of mRNA for a proline-rich protein of cotton fiber. *Plant Physiology Rockville* 108(2): 669-676. {a} Agracetus Inc., Fiber Technol. Div., 8520 Univ. Green, Middleton, WI 53562, USA

Johnson, D. H., J. D. Beaty, et al. (1995). Effects of rotational crop herbicides on rice (*Oryza sativa*). *Weed Science* 43(4): 648-654. {a} American Cyanamid Co., Agric. Res. Div., P.O. Box 400, Princeton, NJ 08543-0400, USA

Johnson, D. H., R. E. Talbert, et al. (1995). Carryover potential of imazaquin to cotton, grain sorghum, wheat, rice, and corn. *Weed Science* 43(3): 454-460. American Cyanamid Co., P.O. Box 400, Princeton, NJ 08543-0440, USA

Joost, O., G. Bianchini, et al. (1995). Differential induction of 3-hydroxy-3-methylglutaryl CoA reductase in two cotton species following inoculation with *Verticillium*. *Molecular Plant Microbe Interactions* 8(6): 880-885. {a} Dep. Plant Pathol. and Microbiol., Tex. A and M Univ., College Station, TX 77843, USA

Kadapa, S. N. (1995). Composite-crossing methodology pays rich dividend in cotton breeding: A resume. *Indian Journal of Genetics and Plant Breeding* 55(3): 290-301. Res., Swananda, Attikalla Behind Railway Goods Shed, Dharwad 5850007, India

Kalsy, H. S., H. R. Garg, et al. (1995). Genetic divergence and heterosis in American cotton. *Crop Improvement* 22(2): 232-236. PAU Regional Res. Stn., Faridkot 151 203, India

Karimov, K. K. and E. A. Meshcheryakova (1995). Mobile forms of nitrogen compounds in the source-sink systems including cotton ovaries. *Fiziologiya Rastanii Moscow* 42(1): 74-78. {a} Inst. Plant Physiol. Biophys., Taj. Acad. Sci., ul. Aini 299/2, 734063 Dushanbe, Tajikistan

Keerio, M. D., M. S. Kalwar, et al. (1995). Genetics of seed cotton yield and its primary components in *Gossypium hirsutum* L. *Pakistan Journal of Botany* 27(2): 425-429. Dep. Plant Breeding Genetics, Sindh Agric. Univ., Tandojam, Sindh, Pakistan

Keisling, T. C., R. F. Ford, et al. (1995). Tillage systems for cotton on Mississippi River Delta and loessial plains soils. *Communications in Soil Science and Plant Analysis* 26(3-4): 441-452. {a} Univ. Arkansas NEREC, P.O. Box 48, Keiser, AR 72351, USA

Keisling, T. C., N. J. Mascagni, Jr., et al. (1995). Using cotton petiole nitrate-nitrogen concentration for prediction of cotton nitrogen nutritional status on a clayey soil. *Journal of Plant Nutrition* 18(1): 35-45. {a} Univ. Arkansas, NEREC, P.O. Box 48, Keiser, AR 72351, USA

Khan, A. N., R. H. Qureshi, et al. (1995). Selection of cotton cultivars for salinity tolerance at seedling stage. *Sarhad Journal of Agriculture* 11(2): 153-159. {a} Dep. Soil Sci., Fac. Agric., Gomal Univ., Dera Ismail Khan, Pakistan

Kirkpatrick, T. L., I. M. W. Van, et al. (1995). Influence of *Meloidogyne incognita* on the water relations of cotton grown in microplots. *Journal of Nematology* 27(4): 465-471. {a} Univ. Ark. Southwest Res. and Extension Cent., Hope, AR 71801-9729, USA

- Kloth, R. H. (1995). Quantitative trait loci affecting cotton fiber and linked to the t-1 locus in upland cotton. *Theoretical and Applied Genetics* 91(5): 762-768. U.S. Dep. Agric., Jamie Whitten Delta States Res. Cent., PO Box 345, Stoneville, MS 38776, USA
- Kloth, R. H. (1995). Interaction of two loci that affect trichome density in upland cotton. *Journal of Heredity* 86(1): 78-80. United States Dep. Agric., Cotton Physiol. Genetics Res. Unit, PO Box 345, Stoneville, MS 38776-0345, USA
- Kudlicka, K., R. M. Brown, Jr., et al. (1995). Beta-Glucan synthesis in the cotton fiber. IV. In vitro assembly of the cellulose 1 allomorph. *Plant Physiology Rockville* 107(1): 111-123. {a} Dep. Botany, Univ. Texas, Austin, TX 78713-7640, USA
- Kumar, V., L. S. Yadav, et al. (1995). Development and reproductive potential of *Helicoverpa armigera* (Hubner) on various plant parts of cotton cultivars/hybrid. *Insect Science and its Application* 16(2): 199-205. {a} Dep. Entomology, CCS Haryana Agricultural Univ., Hisar-125 004, India
- Kushaeva, F. K., A. A. Umarov, et al. (1995). Uptake, migration and distribution of tritiated tetranyl in cotton plants. *Fiziologiya Rastanii Moscow* 42(6): 961-963. {a} Inst. Chem. Plant Subst., Uzb. Acad. Sci., ul. Akad. Kh. Abdullaeva 88, 700170 Tashkent, Uzbekistan
- Lan, S., Z. Xu, et al. (1995). Pollen wall structure using a new stripping-sputtering device for scanning electron microscopy. *Grana* 34(5): 325-331. {a} General Lab., Huazhong Agric. Univ., Wuhan 430070, China
- Lege, K. E., J. T. Cothren, et al. (1995). Phenolic acid and condensed tannin concentrations of six cotton genotypes. *Environmental and Experimental Botany* 35(2): 241-249. Soil Crop Sci. Dep., Texas A and M Univ., College Station, TX 77843-2474, USA
- Liu, F., J. D. Thatcher, et al. (1995). Bifunctional glyoxylate cycle protein of *Caenorhabditis elegans*: A developmentally regulated protein of intestine and muscle. *Developmental Biology* 169(2): 399-414. {a} Dep. Neurol. Baylor Coll. Med., Houston, TX 77030, USA
- Liu, R. J. (1995). Effect of vesicular-arbuscular mycorrhizal fungi on verticillium wilt of cotton. *Mycorrhiza* 5(4): 293-297. Dep. Horticulture Lai-Yang Agric. Coll., Lai-Yang, Shandong 265200, China
- Liu, R. J., H. F. Li, et al. (1995). Detection of pathogenesis-related proteins in cotton plants. *Physiological and Molecular Plant Pathology* 47(6): 357-363. {a} Hortic. Dep., Lai-Yang Agric. Coll., Lai-Yang, Shandong 265200, China
- Lopez, M., J. C. Gutierrez, et al. (1995). Selection and characterization of cotton cultivars for dryland production in the south-west of Spain. *European Journal of Agronomy* 4(1): 119-126. {a} Inst. de Recursos Naturales Agrobiol. de Sevilla, CSIC, Apdo. 1052, E-41080 Sevilla, Spain
- Loughrin, J. H., A. Manukian, et al. (1995). Volatiles emitted by different cotton varieties damaged by feeding beet armyworm larvae. *Journal of Chemical Ecology* 21(8): 1217-1227. {a} Insect Attractants, Behavior Basic Biol. Res. Lab., Agric. Res. Serv., United States Dep. Agric., Gainesville, FL 32604, USA
- Mandeshwar, S. B. (1995). Regeneration of cotton *Gossypium hirsutum* from shoot tip culture. *Advances in Plant Sciences* 8(1): 89-94. Tissue Culture Lab., Crop Improvement Div., C.I.C.R., Post Bag No. 225, GPO, Nagpur, India
- Mann, J. E., J. A. Durant, et al. (1995). Alternatives to pyrethroids for control of *Helicoverpa zea* (Boddie) and *Heliothis virescens* (F.) on cotton in South Carolina. *Journal of Agricultural Entomology* 12(2-3): 127-136. Clemson Univ., Dep. Entomol., Long Hall, Box 340365, Clemson, SC 29634-0365, USA
- Mao, Z., G. Huang, et al. (1995). Genetic analysis of the complete maintainer for genetic-male-sterility in upland cotton. *Science in China Series B Chemistry Life Sciences and Earth Sciences* 38(5): 546-554. Cotton Res. Inst., Sichuan Acad. of Agricultural Sciences, Jianyang 641400, China

Marur, C. J. and L. Sodek (1995). Microwave drying of plant material for biochemical analysis. *Revista Brasileira de Fisiologia Vegetal* 7(1): 111-114. {a} Inst. Agromomico do Parana, Londrina, PR 86001-970, Brazil

May, O. L., D. T. Bowman, et al. (1995). Genetic diversity of U.S. upland cotton cultivars released between 1980 and 1990. *Crop Science* 35(6): 1570-1574. {a} USDA-ARS, Dep. Agronomy, 500 Pocket Road, Florence, SC 29501-9603, USA

May, O. L. and B. C. Bridges, Jr. (1995). Breeding cottons for conventional and late-planted production systems. *Crop Science* 35(1): 132-136. USDA-ARS Clemson Univ., P.O. Box 3039, Florence, SC 29502-3039, USA

McCarty, J. C., Jr. and J. N. Jenkins (1995). Performance of commercial cotton cultivars under elevated densities of tobacco budworm. *Agronomy Journal* 87(5): 804-807. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

McClelland, M. R., M. C. Smith, et al. (1995). Weed control in conservation-tillage cotton in Arkansas. *Arkansas Agricultural Experiment Station Research Series*(442): 49-55. {a} Agronomy, Arkansas Agric. Exp. Stn., Fayetteville, AR 72701, USA

McConnell, J. S., W. H. Baker, et al. (1995). Response of cotton grown on compacted soils to phosphorus and potassium fertilization and deep tillage: Final report. *Arkansas Agricultural Experiment Station Research Series*(443): 96-100. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA

McConnell, J. S., W. H. Baker, et al. (1995). Irrigation methods and nitrogen fertilization rates in cotton production. *Arkansas Agricultural Experiment Station Research Series*(443): 76-82. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA

McConnell, J. S., W. H. Baker, et al. (1995). Cotton yield response to irrigation, reduced tillage and cover crops. *Arkansas Agricultural Experiment Station Research Series*(442): 20-24. {a} Agronomy, Arkansas Agric. Exp. Stn., Fayetteville, AR 72701, USA

McConnell, J. S., R. E. Glover, et al. (1995). Nitrogen fertilization and plant development of cotton as determined by nodes above white flower. *Journal of Plant Nutrition* 18(5): 1027-1036. {a} Univ. Arkansas, Southeast Res. Ext. Cent., Monticello, AR 71655, USA

McInroy, J. A. and J. W. Kloepper (1995). Population dynamics of endophytic bacteria in field-grown sweet corn and cotton. *Canadian Journal of Microbiology* 41(10): 895-901. Dep. Plant Pathol. Alabama Agric. Experiment Stn., Biological Control Inst., Auburn Univ., Auburn, AL 36849-5409, USA

McInroy, J. A. and J. W. Kloepper (1995). Survey of indigenous bacterial endophytes from cotton and sweet corn. *Plant and Soil* 173(2): 337-342. Dep. Plant Pathol., Biol. Control Inst., Ala. Agric. Experiment Station, Auburn Univ., AL 36849-5409, USA

McPherson, G. R., J. N. Jenkins, et al. (1995). Combining ability analysis of root-knot nematode resistance in cotton. *Crop Science* 35(2): 373-375. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

McSorley, R. and D. W. Dickson (1995). Effect of tropical rotation crops on *Meloidogyne incognita* and other plant-parasitic nematodes. *Journal of Nematology* 27(4 Suppl.): 535-544. {a} Dep. Entomol. Nematol., Univ. Florida, Gainesville, FL 32611-0620, USA

Melero, V. J. M., L. M. A. Blanco, et al. (1995). Control of *Verticillium* wilt of cotton by means of soil solarization and tolerant cultivars in southern Spain. *Plant Pathology Oxford* 44(2): 250-260. {a} Dep. Agronomia, ETSIAM, Univ. Cordoba, Cordoba, Spain

Mohan, P., P. Singh, et al. (1995). Gossypol-gland density and free gossypol content in seed and cotyledonary leaf of upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 65(1): 66-68. {a} Div. Crop Improvement, Cent. Inst. Cotton Res., Nagpur, Maharashtra 440 001, India

Morales, R. J. A., K. R. Summy, et al. (1995). Estimating parasitism by *Catolaccus grandis* (Hymenoptera: Pteromalidae) after inundative releases against the boll weevil (Coleoptera: Curculionidae). *Environmental Entomology* 24(6): 1718-1725. Subtropical Agric. Res. Lab., USDA-ARS, Weslaco, TX 78596, USA

Moreira, J. D. A. N., E. C. Freire, et al. (1995). Use of numerical taxonomy to compare "Moco" cotton with other Cotton species and races. *Revista Brasileira de Genetica* 18(1): 99-103. {a} Centro Nacional Pesquisa do Algodao, EMBRAPA, Caixa Postal 174, 58107-720 Campina Grande, PB, Brazil

Morris, D. A. and N. J. Larcombe (1995). Phloem transport and conjugation of foliar-applied 1-aminocyclopropane-1-carboxylic acid in cotton (*Gossypium hirsutum* L.). *Journal of Plant Physiology* 146(4): 429-436. {a} Dep. Biol., Sch. Biol. Sci., Biomed. Sci. Build., Univ. Southampton, Bassett Crescent East, Southampton SO16 7PX, UK

Multani, D. S. and B. R. Lyon (1995). Genetic fingerprinting of Australian cotton cultivars with RAPD markers. *Genome* 38(5): 1005-1008. {a} Sch. Biol. Sci., Univ. Sydney, NSW 2006, Australia

Naranjo, S. E. and H. M. Flint (1995). Spatial distribution of adult *Bemisia tabaci* (Homoptera: Aleyrodidae) in cotton and development and validation of fixed-precision sampling plans for estimating population density. *Environmental Entomology* 24(2): 261-270. Western Cotton Res. Lab., USDA-ARS, 4135 East Broadway Road, Phoenix, AZ 85040, USA

Naranjo, S. E., H. M. Flint, et al. (1995). Comparative analysis of selected sampling methods for adult *Bemisia tabaci* (Homoptera: Aleyrodidae) in cotton. *Journal of Economic Entomology* 88(6): 1666-1678. Western Cotton Res. Lab., USDA-ARS, 4135 East Broadway Road, Phoenix, AZ 85040, USA

Natwick, E. T., C. C. Chu, et al. (1995). Pima and Upland cotton susceptibility to *Bemisia argentifolii* under desert conditions. *Southwestern Entomologist* 20(4): 429-438. {a} Univ. California Cooperative Extension, Holtville, CA 92250, USA

Nimbal, C. I., D. R. Shaw, et al. (1995). Response of MSMA-resistant and -susceptible common cocklebur (*Xanthium strumarium*) biotypes to cotton (*Gossypium hirsutum*) herbicides and cross-resistance to arsenicals and membrane disruptors. *Weed Technology* 9(3): 440-445. {a} Hortic. Dep., Univ. Ky., Lexington, KY 40546, USA

Nirval, B. G., A. A. Chavan, et al. (1995). Management of sowing date in rainy-season crops for sustainable crop yield under dryland condition. *Indian Journal of Agricultural Sciences* 65(3): 170-174. Dryland Agric. Res. Cent., Marathwada Agric. Univ., Parbhani, Maharashtra 431-402, India

Nolte, K. D., D. L. Hendrix, et al. (1995). Sucrose synthase localization during initiation of seed development and trichome differentiation in cotton ovules. *Plant Physiology Rockville* 109(4): 1285-1293. {a} Hortic. Sci. Dep., 1151 Fifield Hall, Univ. Florida, Gainesville, FL 32611, USA

Obroucheva, N. V., O. V. Antipova, et al. (1995). Relationship between initiation of cell elongation and cell division in radicles of germinating seeds. *Plant and Soil* 173(2): 311-316. Inst. Plant Physiol., Russ. Acad. Sci., Botanical St. 35, 127276 Moscow, Russia

Ogle, H. J., A. M. Stirling, et al. (1995). Some factors affecting the development and biocontrol of cotton seedling disease. *Australian Journal of Experimental Agriculture* 35(6): 771-776. Dep. Agric., Univ. Queensland, Brisbane, QLD 4072, Australia

Oliveira, E. D., J. D. C. Machado, et al. (1995). Survival of *Botryodiplodia theobromae* Pap. associated with cotton (*Gossypium hirsutum* L.) seeds in soil. *Ciencia e Pratica* 19(3): 297-301. {a} CNPMS/EMBRAPA-Sete Lagoas, MG

Owen, D. F. and J. R. Gannaway (1995). Registration of 12 glandless cotton germplasm lines with improved fiber strength: TX-8702gl to TX-8713gl. *Crop Science* 35(5): 1515. {a} Texas Agric. Experiment Station, Rt. 3, Box 219, Lubbock, TX 79401, USA

Pandey, S. C., M. K. Saxena, et al. (1995). Studies on genetic variability in upland cotton (*Gossypium hirsutum* L.). *Crop Research Hisar* 10(3): 277-278. J. N. K. V. V. Coll. Agric., Indore-452 001, India

Panwar, R. S., R. K. Malik, et al. (1995). Influence of tank mixture of herbicides on weed control in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 40(2): 347-348. Dep. Agronomy, Chaudhary Charan Singh Haryana Agric. Univ., Hisar 125 004, India

Paplomatas, E. J., K. Elena, et al. (1995). First report of *Phytophthora boehmeriae* causing boll rot of cotton. *Plant Disease* 79(8): 860. Benaki Phytopahtol. Inst., 145 61 Kifissia-Athens, Greece

Patel, I. S. and D. N. Yadav (1995). Susceptibility of *Amrasca biguttula biguttula* and *Chrysopa scelestes* in cotton (*Gossypium* species) to three systemic insecticides. *Indian Journal of Agricultural Sciences* 65(4): 308-309. {a} Main Castor Res. Stn., GAU, Sardar Krushinagar 385 506, India

Perera, N. H., E. Hartmann, et al. (1995). Regulation of cotton photosynthesis during moderate chilling. *Plant Science Limerick* 111(2): 133-143. {a} Dep. Biological Sciences, Texas Tech. Univ., Lubbock, TX 79409-3131, USA

Pettigrew, W. T. (1995). Source-to-sink manipulation effects on cotton fiber quality. *Agronomy Journal* 87(5): 947-952. USDA-ARS, Cotton Physiol. Genetics Res., P.O. Box 345, Stoneville, MS 38776, USA

Plant, R. E. and T. A. Kerby (1995). CPM: Software for cotton final plant mapping. *Agronomy Journal* 87(6): 1143-1147. {a} Dep. Agron. Range Sci., Univ. Calif., Davis, CA 95616, USA

Porter, P. M. and A. Khalilian (1995). Wheat response to row spacing in relay intercropping systems. *Agronomy Journal* 87(5): 999-1003. {a} Dep. Agronomy and Plant Genetics, Southwest Exp. Stn., Univ. Minnesota, Box 428, Lamberton, MN 56152, USA

Priolo, N. S., M. C. Arribere, et al. (1995). Isolation and partial characterization of a lipase from primary roots of *Gossypium hirsutum*. *Acta Farmaceutica Bonaerense* 14(1): 31-39. LIPROVE, Departamento de Ciencias Biologicas, Facultad de Ciencias Exactas, Universidad Nacional de La Plata, Casilla de Correo 711, 1900 La Plata, Argentina

Prior, S. A., H. H. Rogers, et al. (1995). Free-air carbon dioxide enrichment of cotton: Root morphological characteristics. *Journal of Environmental Quality* 24(4): 678-683. {a} Natl. Soil Dyn. Lab., USDA-Agric. Res. Serv., P.O. Box 3439, Auburn, IL 36831, USA

Qi, X., B. X. Behrens, et al. (1995). Solubilization and partial characterization of extensin fragments from cell walls of cotton suspension cultures: Evidence for a covalent cross-link between extensin and pectin. *Plant Physiology Rockville* 108(4): 1691-1701. {a} Departments Biochemistry and Molecular Biology, Oklahoma State Univ., Stillwater, OK 74078, USA

Queiroz, V. R. B. (1995). Anatomical alterations in cotton plants with reddish withering symptoms. *Bragantia* 54(1): 39-46. Secao Bot. Econ., Inst. Agron., Caixa Postal 28, 13001-970 Campinas, SP, Brazil

Ragazzi, A., S. Moricca, et al. (1995). Infection of cotton by *Fusarium oxysporum* f. sp. *vasinfectum* as affected by water stress. *Phytoparasitica* 23(4): 315-321. {a} Ist. Patol. Zool. Froestale Agraria, Univ. Firenze, 50144-Firenze, Italy

Reddy, K. R., M. L. Boone, et al. (1995). Developing and validating a model for a plant growth regulator. *Agronomy Journal* 87(6): 1100-1105. {a} Dep. Plant Soil Sci., Mississippi State Univ., Mississippi State, MS 39762, USA

Reddy, V. R. (1995). Modeling ethephon-temperature interactions in cotton. *Computers and Electronics in Agriculture* 13(1): 27-35. Syst. Res. Lab., USDA-ARS, Build. 007, Room 008, BARC-West, Beltsville, MD 20705, USA

Reddy, V. R., K. R. Reddy, et al. (1995). Carbon dioxide and temperature interactions on stem extension, node initiation, and fruiting in cotton. *Agriculture Ecosystems and Environment* 55(1): 17-28. {a} USDA/ARS Syst. Res. Lab., 10300 Baltimore Ave., Build. 007, Room 008, BARC-West, Beltsville, MD 20705, USA

Reddy, V. R., K. R. Reddy, et al. (1995). Carbon dioxide enrichment and temperature effects on cotton canopy photosynthesis, transpiration, and water-use efficiency. *Field Crops Research* 41(1): 13-23. {a} Dep. Plant Soil Sci., Mississippi State Univ., Mississippi State, MS 39762, USA

Reeves, D. W. and G. L. Mullins (1995). Subsoiling and potassium placement effects on water relations and yield of cotton. *Agronomy Journal* 87(5): 847-852. {a} USDA-ARS Natl. Soil Dynamics Lab., P.O. Box 3439, Auburn, AL 36831-3439, USA

Reinhardt, D. H. and T. L. Rost (1995). On the correlation of primary root growth and tracheary element size and distance from the tip in cotton seedlings grown under salinity. *Environmental and Experimental Botany* 35(4): 575-588. {a} EMBRAPA/CNP-Mandioca e Fruticultura, Caixa Postal 007, 44380 Cruz das Almas, BA, Brazil

Reinhardt, D. H. and T. L. Rost (1995). Salinity accelerates endodermal development and induces an exodermis in cotton seedling roots. *Environmental and Experimental Botany* 35(4): 563-574. {a} EMBRAPA/CNPMF, Caixa Postal 7, 44380 Cruz das Almas, Bahia, Brazil

Reinhardt, D. H. and T. L. Rost (1995). Primary and lateral root development of dark- and light-grown cotton seedlings under salinity stress. *Botanica Acta* 108(5): 457-465. {a} EMBRAPA, CNPMF, Caixa Postal 7, 44380 Cruz das Almas, Bahia, Brazil

Reinhardt, D. H. and T. L. Rost (1995). Developmental changes of cotton root primary tissues induced by salinity. *International Journal of Plant Sciences* 156(4): 505-513. EMBRAPA/CNPMF, Caixa Postal 7, 44380 Cruz de Almas, Bahia, Brazil

Rothrock, C. S., T. L. Kirkpatrick, et al. (1995). The influence of winter legume cover crops on soilborne plant pathogens and cotton seedling diseases. *Plant Disease* 79(2): 167-171. {a} Dep. Plant Pathol., Univ. Arkansas, Fayetteville, AR 72701, USA

Russomanno, O. M. R., L. N. Coutinho, et al. (1995). Pathogenicity preservation of *Verticillium dahliae* cultures maintained in distilled water. *Summa Phytopathologica* 21(2): 178-181. {a} Inst. Biol.-Caixa Postal 7119, 0164-970, Sao Paulo, SP, Brazil

Sabino, N. P., J. I. Kondo, et al. (1995). Effect of the cotton fiber strength on yarn properties. *Bragantia* 54(1): 121-130. Secao Tecnol. Fibras, Inst. Agron., Caixa Postal 28, 13001-970 Campinas, SP, Brazil

Safiyazov, J. S., R. N. Mannanov, et al. (1995). The use of bacterial antagonists for the control of cotton diseases. *Field Crops Research* 43(1): 51-54. {a} Inst. Microbiol., Uzbek Acad. Sci., 7b A. Kodiry Str., Tashkent 700128, Uzbekistan

Sandoval, J. A., Z. H. Huang, et al. (1995). N-Acylphosphatidylethanolamine in dry and imbibing cottonseeds. Amounts, molecular species, and enzymatic synthesis. *Plant Physiology Rockville* 109(1): 269-275. {a} Dep. Biol. Sci., Div. Biochem., Univ. North Texas, Denton, TX 76203-0218, USA

Sarkar, R. K., A. Chakraborty, et al. (1995). Effect of intercropping oilseed and pulse crops in upland cotton (*Gossypium hirsutum*) for total productivity and monetary advantage in the system. *Indian Journal of Agricultural Sciences* 65(4): 246-249. {a} Univ. Coll. Agric., Calcutta Univ., Calcutta, W. Bengal 700 019, India

- Sassenrath, C. G. F. (1995). Dependence of canopy light distribution on leaf and canopy structure for two cotton (*Gossypium*) species. *Agricultural and Forest Meteorology* 77(1-2): 55-72. USDA -ARS Crop Simulation Res. Unit, Dep. Plant Soil Sci., P.O. Box 5367, Mississippi State University, MS 39762-5367, USA
- Sethar, M. A., D. L. Laidman, et al. (1995). Hardening potential in cotton (*Gossypium hirsutum* L.) at high temperature stress. *Pakistan Journal of Botany* 27(1): 165-168. {a} Dep. Physiol. Biochem., Sindh Agric. Univ., Tandojam, Pakistan
- Sharma, R. K. and R. S. S. Tomar (1995). Effect of spacings and levels of nitrogen on cotton cultivars (*G. hirsutum* L.). *Agricultural Science Digest* 15(1-2): 5-7. Natl. Agric. Res. Project, JNKVV-Zonal Agric. Res. Sta., Morena-476001, India
- Sharma, R. K., R. S. S. Tomar, et al. (1995). Response of upland cotton (*Gossypium hirsutum*) hybrids to varying levels of nitrogen with and without neemcake under irrigated condition of Malwa tract of Madhya Pradesh. *Indian Journal of Agronomy* 40(2): 344-346. {a} KVK Coll. Agric., Khandwa, Madhya Pradesh 450 001, India
- Shennan, C., S. R. Grattan, et al. (1995). Feasibility of Cyclic Reuse of Saline Drainage in a Tomato-Cotton Rotation. *Journal of Environmental Quality* 24(3): 476-486. {a} Dep. Vegetable Corps, Univ. California, Davis, CA 95616, USA
- Shimizu, T., N. Hashimoto, et al. (1995). A novel isourazole herbicide, fluthiacet-methyl, is a potent inhibitor of protoporphyrinogen oxidase after isomerization by glutathione S-transferase. *Plant and Cell Physiology* 36(4): 625-632. {a} Life Sci. Res. Inst., Kumiai Chem. Ind. Co. Ltd., Kikugawa-cho, Ogasa-gun, Shizuoka 439, Japan
- Shutov, A. D., I. A. Kakhovskaya, et al. (1995). Legumin-like and vicilin-like seed storage proteins: Evidence for a common single-domain ancestral gene. *Journal of Molecular Evolution* 41(6): 1057-1069. {a} Inst. Plant Genet., Crop Plant Res., Corrensstr. 3, D-06466 Gatersleben, Germany
- Simwat, G. S. and A. K. Dhawan (1995). Population distribution of cotton jassid (*Amrasca biguttula biguttula*) and whitefly (*Bemisia tabaci*) on upland cotton (*Gossypium hirsutum*) at different times during day and levels of plant canopy. *Indian Journal of Agricultural Sciences* 65(1): 84-86. Dep. Entomol., Punjab Agric. Univ., Ludhiana 141 004, India
- Singh, B., G. S. Chahal, et al. (1995). Efficiency of different selection criteria for the improvement of seed cotton yield in early segregating generations of *Gossypium hirsutum* L. *Crop Improvement* 22(1): 61-64. {a} Biotechnol. Centre, Punjab Agric. Univ., Ludhiana, India
- Singh, K. N. and D. P. Sharma (1995). Effect of subsurface drain spacing and nitrogen level on yield and growth of upland cotton (*Gossypium hirsutum*) and pearl millet (*Pennisetum glaucum*) in saline soil. *Indian Journal of Agricultural Sciences* 65(1): 27-33. Cent. Soil Salinity Res. Inst., Karnal, Haryana 132 001, India
- Singh, V. V., S. S. Narayanan, et al. (1995). Evaluation of selected germplasm of tetraploid cultivated cottons (*Gossypium* species) for amino acid profile. *Indian Journal of Agricultural Sciences* 65(1): 24-26. {a} Div. Crop Improvement, Cent. Inst. Cotton Res., Nagpur, Maharashtra 440 001, India
- Siwach, S. S. and B. P. S. Lather (1995). Gain through varietal improvement in upland cotton. *Crop Improvement* 22(2): 264-265. Cotton Sect., Dep. Plant Breeding, CCS, HAU, Hissar, India
- Smith, C., M. McClelland, et al. (1995). Cover crops and herbicide systems in no-till cotton. *Arkansas Agricultural Experiment Station Research Series*(442): 56-62. {a} Agronomy, Arkansas Agric. Exp. Stn., Fayetteville, AR 72701, USA
- Sohu, R. S. and G. S. Chahal (1995). Efficiency of different selection criteria to improve yield and earliness in cotton. *Crop Improvement* 22(2): 210-213. Dep. Plant Breeding, Punjab Agric. Univ., Ludhiana 141 004, India

- Solaiappan, U., A. Muthusankaranarayanan, et al. (1995). Effect of humic acid on rainfed upland cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 40(1): 156-157. Agric. Res. Stn., Tamil Nadu Agric. Univ., Kovilpatti 627 701, India
- Song, F. M. Z. Z. (1995). Involvement of cell-wall hydroxyproline and free proline in the resistance of cotton seedlings to *Fusarium* wilt. *Acta Phytophysiologica Sinica* 21(3): 235-241. Dep. Plant Protection, Zhejiang Agricultural Univ., Hangzhou 310029, China
- Soomro, Z. A., M. S. Kalwar, et al. (1995). Genetic analysis of yield and yield components in intraspecific crosses of *Gossypium hirsutum* L. *Pakistan Journal of Botany* 27(2): 431-434. Dep. Plant Breeding Genetics, Sindh Agric. Univ., Tandojam, Sindh, Pakistan
- Subbarao, K. V., A. Chassot, et al. (1995). Genetic relationships and cross pathogenicities of *Verticillium dahliae* isolates from cauliflower and other crops. *Phytopathology* 85(10): 1105-1112. {a} Dep. Plant Pathol., Univ. Calif., Davis, c/o U.S. Agric. Res. Stn., Salinas, CA 93905, USA
- Subbian, P., P. Gopalasundaram, et al. (1995). Energetics and water-use efficiency of intensive cropping system. *Indian Journal of Agronomy* 40(3): 398-401. Dep. Agronomy, Tamil Nadu Agric. Univ., Coimbatore 641 003, India
- Sumner, D. R., C. C. Dowler, et al. (1995). Conservation tillage and seedling diseases in cotton and soybean double-cropped with triticale. *Plant Disease* 79(4): 372-375. {a} Dep. Plant Pathol., Univ. Ga., Tifton, GA, USA
- Tarhalkar, P. P. and M. V. Venugopalan (1995). Effect of organic recycling of fodder legumes in stabilizing productivity of rainfed cotton (*Gossypium hirsutum* L.) on marginal lands. *Tropical Agriculture* 72(1): 73-75. Div. Crop. Production, Central Inst. Cotton Res., Nagpur 440 001, India
- Tariq, M., M. A. Khan, et al. (1995). Inheritance of lint percent, seed and lint indices and fibre length in upland cotton (*G. hirsutum* L.). *Sarhad Journal of Agriculture* 11(5): 607-617. {a} Millet Res. Station, Rawalpindi, Pakistan
- Tariq, M., M. A. Khan, et al. (1995). Inheritance of boll weight, boll number and yield of seed cotton in upland cotton (*G. hirsutum* L.). *Sarhad Journal of Agriculture* 11(5): 599-605. {a} Millet Res. Station, Rawalpindi, Pakistan
- Tewolde, H., C. J. Fernandez, et al. (1995). Critical petiole nitrate-nitrogen for lint yield and enhanced maturity in Pima cotton. *Agronomy Journal* 87(2): 223-227. {a} Texas A and M Univ. Agric. Res. and Ext. Cent., 1619 Garner Field Road, Uvalde, TX 78801, USA
- Thomas, J. C., D. G. Adams, et al. (1995). Protease inhibitors of *Manduca sexta* expressed in transgenic cotton. *Plant Cell Reports* 14(12): 758-762. {a} Dep. Biochem., Univ. Ariz., Tucson, AZ 85721, USA
- Timpa, J. D., A. M. Striegel, et al. (1995). Rapid, inexpensive method for determining glucose concentrations in cotton bolls and other plant tissues. *Crop Science* 35(1): 274-278. {a} USDA-ARS, Southern Regional Research Center, P.O. Box 19687, New Orleans, LA 70179, USA
- Tiwari, S. C. and T. A. Wilkins (1995). Cotton (*Gossypium hirsutum*) seed trichomes expand via diffuse growing mechanism. *Canadian Journal of Botany* 73(5): 746-757. {a} Dep. Agronomy Range Sci., Univ. California, Davis, CA 95616-8515, USA
- Tomar, S. K., O. S. Rana, et al. (1995). Effect of sowing date on yield of cotton (*Gossypium* species). *Indian Journal of Agronomy* 40(2): 342-343. Cotton Res. Stn., Govind Ballabh Pant Univ. Agric. Technol., Bulandshahr, Uttar Pradesh 203 001, India
- Tonkikh, A. K., T. A. Babaev, et al. (1995). Mechanism of the auxin-like action of thyroxine. *Fiziologiya Rastanii Moscow* 42(2): 195-200. {a} Fac. Biol., Tashk. State Univ., Vuzgorodok, GSP, 700095 Tashkent, Uzbekistan

Torbert, H. A., S. A. Prior, et al. (1995). Elevated atmospheric carbon dioxide effects on cotton plant residue decomposition. *Soil Science Society of America Journal* 59(5): 1321-1328. {a} USDA-ARS Blackland, Soil Water Res. Lab., 808 East Blackland Rd., Temple, TX 76502, USA

Torbert, H. A. and D. W. Reeves (1995). Interactions of traffic and tillage applied to cotton on N movement below the root zone of a subsequent wheat crop. *Soil and Tillage Research* 33(1): 3-16. {a} USDA-ARS Blackland, Soil Water Res. Lab., 808 E. Blackland Rd., Temple, TX 76502, USA

Turley, R. B., D. L. Ferguson, et al. (1995). A cDNA encoding ribosomal protein S4e from cotton (*Gossypium hirsutum* L.). *Plant Physiology Rockville* 108(1): 431-432. {a} United States Dep. Agric., Agricultural Res. Service, Cotton Physiol. Genetics Res., PO Box 345, Stoneville, MS 38776, USA

Van, I. M. W., J. J. Heitholt, et al. (1995). Foliar methanol applications to cotton in the southeastern United States: Leaf physiology, growth, and yield components. *Agronomy Journal* 87(6): 1157-1160. {a} Dep. Hortic., Georgia Stn., Univ. Ga., Griffin, GA 30223, USA

Van, I. M. W. and D. M. Oosterhuis (1995). Diurnal water relations of expanding and full-sized cotton fruits and subtending leaves. *Plant Cell and Environment* 18(7): 807-812. {a} Dep. Plants Soils Biometeorol., Utah State Univ., Logan, UT 84322-480, USA

Van, L. M. W., W. M. Harris, et al. (1995). Phloem in developing cotton fruits: 6(5)-carboxyfluorescein as a tracer for functional phloem. *Journal of Experimental Botany* 46(284): 321-328. {a} Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA

Varghese, S., K. V. Patel, et al. (1995). Heterosis for oil and gossypol content in hybrids of upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 65(10): 760-762. Main Cotton Res. Stn., Gujarat Agric. Univ., Surat 395 007, India

Vasil, e. A. A. (1995). Optimization model of stomatal regulation in C-3-plants: 4. Response of stomata to changes in oxygen concentration. *Fiziologiya Rastenii Moscow* 42(4): 598-602. Inst. Plant Physiol., Russ. Acad. Sci., ul. Botanicheskaya 35, 127276 Moscow, Russia

Venugopalan, M. V., P. P. Tarhalkar, et al. (1995). Efficacy of phosphate carriers as foliar fertilizer on rainfed upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 65(5): 328-333. Div. Crop Production, Central Inst. Cotton Res., Nagpur, Maharashtra 440 001, India

Vietor, D. M., J. M. Chandler, et al. (1995). Should public funds support biotechnology development? A case about herbicide-resistant cotton. *Journal of Natural Resources and Life Sciences Education* 24(2): 173-178. {a} Soil Crop Sci. Dep., Texas A and M Univ., Coll. Station, TX 77843-2474, USA

Vigil, E. L. and T. K. Fang (1995). Protease activities and elongation growth of excised cotton seed axes during the first 24 hours of imbibition. *Seed Science Research* 5(4): 201-207. {a} Climate Stress Lab., USDA/ARS, B-046A, BARC-West, Beltsville, MD 20705, USA

Vigil, E. L. and T. K. Fang (1995). Comparative biochemical and morphological changes in imbibed cotton seed hypocotyls and radicles in situ and in vitro - Protein breakdown and elongation growth. *Seed Science Research* 5(1): 41-51. {a} Climate Stress Lab., USDA/ARS, Bld 046A, BARC-West, Beltsville, MD 20705, USA

Wang, G. L., J. M. Dong, et al. (1995). The distribution of *Gossypium hirsutum* chromatin in *G. barbadense* germ plasm: Molecular analysis of introgressive plant breeding. *Theoretical and Applied Genetics* 91(6-7): 1153-1161. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843, USA

Wang, X. F. and D. Yakir (1995). Temporal and spatial variations in the oxygen-18 content of leaf water in different plant species. *Plant Cell and Environment* 18(12): 1377-1385. {a} Dep. Environmental Sci. Energy Research, Weizmann Inst. Sci., Rehovot 76100, Israel

- Wang, Z., Y. Yin, et al. (1995). The effect of DPC (N,N-dimethyl piperidinium chloride) on the ^{14}C - ^{14}C assimilation and partitioning of ^{14}C assimilates within the cotton plants interplanted in a wheat stand. *Photosynthetica Prague* 31(2): 197-202. Dep. Agron., Shandong Agric. Univ., Taian, Shandong Province 271018, China
- Wang, Z. L. S. C. W., X. X. Zhang, et al. (1995). Peroxidase isozyme study in the interspecific advanced lines from *Gossypium*. *Acta Botanica Sinica* 37(6): 465-470. Inst. Genetics, Academia Sinica, Beijing 100101, China
- Warner, D. A., A. S. Holaday, et al. (1995). Response of carbon metabolism to night temperature in cotton. *Agronomy Journal* 87(6): 1193-1197. {a} USDA Plant Stress Water Conservation Res. Unit, Route 3, Box 215, Lubbock, TX 79401, USA
- Weathersbee, A. A., III, D. D. Hardee, et al. (1995). Differences in yield response to cotton aphids (Homoptera: Aphididae) between smooth-leaf and hairy-leaf isogenic cotton lines. *Journal of Economic Entomology* 88(3): 749-754. {a} Southern Insect Manage. Lab., USDA-ARS, Stoneville, MS 38776, USA
- Wendel, J. F., A. Schnabel, et al. (1995). An Unusual Ribosomal DNA Sequence from *Gossypium gossypioides* Reveals Ancient, Cryptic, Intergenomic Introgression. *Molecular Phylogenetics and Evolution* 4(3): 298-313. {a} Dep. Bot., Iowa State Univ., Ames, IA 50011, USA
- Williams, H. J., I. Sattler, et al. (1995). Diversity in cyclic sesquiterpene production by *Gossypium hirsutum*. *Phytochemistry Oxford* 40(6): 1633-1636. {a} Cent. Biological NMR, Dep. Chem., Texas A and M Univ., College Station, TX 77843, USA
- Wood, D. C., L. V. Vu, et al. (1995). Purification and characterization of neomycin phosphotransferase II from genetically modified cottonseed (*Gossypium hirsutum*). *Journal of Agricultural and Food Chemistry* 43(4): 1105-1109. {a} AA4I, Monsanto Co., 700 Chesterfield Parkway N., St. Louis, MO 63198, USA
- Yamauchi, A., H. M. Taylor, et al. (1995). Axial resistance to water flow of intact cotton taproots. *Agronomy Journal* 87(3): 439-445. {a} USDA-ARS, Cropping Systems Res. Lab., Route 3, Box 215, Lubbock, TX 79401, USA
- Yang, Y. and D. W. Gabriel (1995). Intragenic recombination of a single plant pathogen gene provides a mechanism for the evolution of new host specificities. *Journal of Bacteriology* 177(17): 4963-4968. {a} Plant Molecular and Cellular Biology Program and Plant Pathology Dep., Univ. of Florida, Gainesville, FL 32611, USA
- Zhao, X., R. A. Wing, et al. (1995). Cloning and characterization of the majority of repetitive DNA in cotton (*Gossypium L.*). *Genome* 38(6): 1177-1188. {a} Dep. Soil and Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA
- Zheng, S., D. Ji, et al. (1995). Evaluation of Fusarium wilt-resistance in primitive races of *Gossypium hirsutum L.* *Genetic Resources and Crop Evolution* 42(3): 257-261. Dep. Agronomy, Zhejiang Agric. Univ., Hangzhou 310029, China
- Abdullaev, A., K. A. Aliev, et al. (1994). Effect of high CO_2 concentration on the characteristics of C-3-plants photosynthetic metabolism. *Fiziologiya i Biokhimiya Kul'turnykh Rastenii* 26(2): 131-135. Inst. Plant Physiol. Biophys., Acad. Sci. Taj., Dushanbe, Tajikistan
- Ahmad, S., A. Hussain, et al. (1994). Heterosis studies on quality traits in intraspecific crosses of *Gossypium hirsutum L.* *Pakistan Journal of Scientific and Industrial Research* 37(4): 153-155. {a} Cotton Res. Inst., Multan, Pakistan
- Ansari, B. A. (1994). Heterotic performance of intraspecific hybrids of *Gossypium hirsutum L.* *Sarhad Journal of Agriculture* 10(6): 681-686. Dep. Plant Breeding, Genetics Fac. Crop Production, Sindh Agric. Univ., Tandojam, Pakistan

- Ashraf, M., M. M. Saeed, et al. (1994). Tolerance to high temperature in cotton (*Gossypium hirsutum* L.) at initial growth stages. *Environmental and Experimental Botany* 34(3): 275-283. {a} Inst. Pure Applied Biol., Bahauddin Zakariya Univ., Multan, Pakistan
- Aslam, M., R. M. S. Iqbal, et al. (1994). Pollen irradiation in cotton (*Gossypium hirsutum* L.). *Pakistan Journal of Botany* 26(2): 341-346. Nuclear Inst. Agric. and Biol., Faisalabad, Pakistan
- Averina, N. G., I. D. Shegai, et al. (1994). An analysis of the phytochrome content in etiolated cotyledon leaves of a chlorophyll-deficient cotton mutant. *Fiziologiya i Biokhimiya Kul'turnykh Rastenii* 26(1): 22-26. Inst. Photobiol., Acad. Sci. Belarus, Minsk, Belarus
- Avtar, R., B. S. Chhabra, et al. (1994). Hybrid-seed production in cytoplasmic male-sterile lines of upland cotton (*Gossypium hirsutum*) through insect pollination. *Indian Journal of Agricultural Sciences* 64(7): 459-461. Dep. Plant Breed., Chaudhary Charan Singh Haryana Agric. Univ., Hisar 125 004, India
- Avtar, R., R. Kumar, et al. (1994). Boll-setting potential in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 64(1): 62-63. Dep. Plant Breeding, Chaudhary Charan Singh Haryana Agric. Univ., Hisar 125 004, India
- Baker, W. H., J. S. McConnell, et al. (1994). Yield response of dryland cotton to supplemental foliar potassium nitrate. *Arkansas Agricultural Experiment Station Research Series*(436): 41-43. {a} Soil Test Lab., Marianna, AR, USA
- Baker, W. H., J. S. McConnell, et al. (1994). Petiole sap as a method for monitoring cotton nitrogen and potassium in the field. *Arkansas Agricultural Experiment Station Research Series*(436): 71-78. {a} Soil Test Lab., Marianna, AR, USA
- Baker, W. H., S. J. McConnell, et al. (1994). Yield response of cotton to supplemental foliar urea. *Arkansas Agricultural Experiment Station Research Series*(436): 79-82. {a} Soil Test Lab., Marianna, AR, USA
- Baker, W. H., W. E. Sabbe, et al. (1994). Assessment of cotton nitrogen status using a hand-held chlorophyll meter. *Arkansas Agricultural Experiment Station Research Series*(436): 67-70. {a} Soil Test Lab., Marianna, AR, USA
- Ball, R. A., D. M. Oosterhuis, et al. (1994). Growth dynamics of the cotton plant during water-deficit stress. *Agronomy Journal* 86(5): 788-795. {a} Dep. Agronomy, Altheimer Lab., Univ. Arkansas, 276 Altheimer Dr., Fayetteville, AR 72703, USA
- Bararpour, M. T., R. E. Talbert, et al. (1994). Spotted spurge (*Euphorbia maculata*) interference with cotton (*Gossypium hirsutum*). *Weed Science* 42(4): 553-555. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA
- Barkhatova, T. A. (1994). Problem of cotton leaf abscission: Actuality and consideration level (review). *Sel'skokhozyaistvennaya Biologiya*(1): 13-19. Sci. Prod. Assoc. "Biolog", Acad. Sci. Uzb., Tashkent, Uzbekistan
- Basallote, M. J., J. Bejarano, et al. (1994). Soil solarization: A strategy for the control of diseases caused by soilborne plant pathogens and for the reduction of crop rotations. *Investigacion Agraria Produccion y Proteccion Vegetales*(Suppl. 2): 207-220. {a} Depto. Proteccion Vegetal, CIDA Cordoba, Apartado 4240, 1480 Cordoba, Spain
- Bauske, E. M., K. R. Rodriguez, et al. (1994). Management of *Meloidogyne incognita* on cotton by use of botanical aromatic compounds. *Nematropica* 24(2): 143-150. {a} Dep. Hortic., Alabama Cooperative Extension Serv., Auburn Univ., Auburn, AL 36849, USA
- Bednarz, C. W., D. M. Oosterhuis, et al. (1994). Plant physiological aspects of potassium nutrition in cotton. *Arkansas Agricultural Experiment Station Research Series*(436): 54-57. {a} Agron. Dep., Fayetteville, AR, USA

Benedict, C. R., R. J. Kohel, et al. (1994). Crystalline cellulose and cotton fiber strength. *Crop Science* 34(1): 147-151. {a} Dep. Biochem. Biophysics, Texas A and M Univ., College Station, TX 77843, USA

Betsche, T. (1994). Atmospheric CO₂ enrichment: Kinetics of chlorophyll a fluorescence and photosynthetic CO₂ uptake in individual, attached cotton leaves. *Environmental and Experimental Botany* 34(1): 75-86. Dep. Physiol. Veg. et Ecosystemes, CEA, Centre Cadarache, F 13108 St. Paul lez Durance, France

Bhatade, S., V. G. Reddy, et al. (1994). Stability analysis of seed-cotton yield in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 64(12): 875-877. Cotton Research Station, Marathwada Agric. Univ., Nanded, Maharashtra 431 604, India

Bilbro, J. D. and D. W. Fryrear (1994). Wind erosion losses as related to plant silhouette and soil cover. *Agronomy Journal* 86(3): 550-553. {a} Cropping Systems Res. Lab., USDA-ARS, P.O. Box 909, Big Spring, TX 79721-0909, USA

Bilger, W. and O. Bjorkman (1994). Relationship among violaxanthin deepoxidation, thylakoid membrane conformation, and nonphotochemical chlorophyll fluorescence quenching in leaves of cotton (*Gossypium hirsutum* L.). *Planta Heidelberg* 193(2): 238-246. {a} Carnegie Inst. Wash., Dep. Plant Biol., 290 Panama St., Stanford, CA 94305, USA

Boman, R. K. and R. L. Westerman (1994). Nitrogen and mepiquat chloride effects on the production of nonrank, irrigated, short-season cotton. *Journal of Production Agriculture* 7(1): 70-75. {a} Dep. Agron., Okla. State Univ., Stillwater, OK 74078, USA

Bondada, B. R., D. M. Oosterhuis, et al. (1994). Cotton yield response to foliar nitrogen fertilization in relation to boll load, petiole and fertilizer nitrogen. *Arkansas Agricultural Experiment Station Research Series*(436): 88-91. {a} Agron. Dep., Fayetteville, AR, USA

Bondada, B. R., D. M. Oosterhuis, et al. (1994). Anatomical considerations related to photosynthesis in cotton (*Gossypium hirsutum* L.) leaves, bracts, and the capsule wall. *Journal of Experimental Botany* 45(270): 111-118. {a} Altheimer Lab., Dep. Agron., Univ. Arkansas, Fayetteville, AR 72703, USA

Boquet, D. J., E. B. Moser, et al. (1994). Boll weight and within-plant yield distribution in field-grown cotton given different levels of nitrogen. *Agronomy Journal* 86(1): 20-26. {a} Louisiana Agricultural Exp. Stn., Northeast Res. Stn., P. O. Box 438, St. Joseph, LA 71366, USA

Bordovsky, J. P., W. M. Lyle, et al. (1994). Crop rotation and tillage effects on soil water and cotton yield. *Agronomy Journal* 86(1): 1-6. {a} Texas A and M Agricultural Research and Ext. Center, Rt. 1, Box 219, Lubbock, TX 79401-9757, USA

Bowman, D. T. and D. P. Schmitt (1994). Screening cotton for tolerance to *Hoplolaimus columbus*. *Plant Disease* 78(7): 695-697. {a} Crop Sci. Dep., North Carolina State Univ., Raleigh, NC 27695, USA

Brock, P. M., J. R. B. Inwood, et al. (1994). Systemic induced resistance to *Alternaria macrospora* in cotton (*Gossypium hirsutum*). *Australasian Plant Pathology* 23(3): 81-85. Dep. Crop Sci., Univ. Sydney, Sydney, NSW 2006, Australia

Brouder, S. M. and K. G. Cassman (1994). Evaluation of a mechanistic model of potassium uptake by cotton in vermiculitic soil. *Soil Science Society of America Journal* 58(4): 1174-1183. Dep. Agron. Range Sci., Univ. Calif., Davis, CA 95616, USA

Brouder, S. M. and K. G. Cassman (1994). Cotton root and shoot response to localized supply of nitrate, phosphate and potassium: Split-pot studies with nutrient solution and vermiculitic soil. *Plant and Soil* 161(2): 179-193. {a} Dep. Agronomy Range Science, Univ. Calif.-Davis, Davis, CA 95616, USA

Brown, B. A., R. M. Hayes, et al. (1994). Effect of tillage and cover crop on fluometuron adsorption and degradation under controlled conditions. *Weed Science* 42(4): 629-634. {a} P.O. Box 1071, Knoxville, TN 37901, USA

Brubaker, C. L. and J. F. Wendel (1994). Reevaluating the origin of domesticated cotton (*Gossypium hirsutum*: Malvaceae) using nuclear restriction fragment length polymorphisms (RFLPs). *American Journal of Botany* 81(10): 1309-1326. {a} Dep. Bot., Iowa State Univ., Ames, IA 50011, USA

Bryson, C. T., J. E. Hanks, et al. (1994). Purple nutsedge (*Cyperus rotundus*) control in reduced-tillage cotton (*Gossypium hirsutum* L.) with low-volume technology. *Weed Technology* 8(1): 28-31. {a} USDA/ARS, South Weed Sci. Lab., Stoneville, MS 38776, USA

Carvalho, C. A. M. and A. C. Fraga (1994). The effects of temperature and imbibition time on cotton seeds (*Gossypium hirsutum* L.) during the exudate (phenolphthalein) pH test. *Ciencia e Pratica* 18(4): 365-369. {a} Cx. Postal 43, Pitangui, MG, CEP 35.650-000, Brazil

Carvalho, L. H., E. J. Chiavegato, et al. (1994). Effects of growth phytohormones and pruning on the cotton crop. *Bragantia* 53(2): 247-254. {a} Secao Algodao, Inst. Agronomico, Caixa Postal 28, 13001-970 Campinas, Brazil

Chu, C. C., D. H. Akey, et al. (1994). Response of cotton plants to aldicarb soil application. *Journal of the Agricultural Association of China*(167): 76-90. {a} USDA-ARS Western Cotton Res. Lab., Brawley, CA 92227, USA

Cohen, A. C. and D. L. Hendrix (1994). Demonstration and preliminary characterization of alpha-amylase in the sweetpotato whitefly, *Bemisia tabaci* (Aleyrodidae: Homoptera). *Comparative Biochemistry and Physiology B Comparative Biochemistry and Molecular Biology* 109(4): 593-601. {a} Western Cotton Res. Lab., USDA, ARS, 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

Corbin, B. R., Jr., M. McClelland, et al. (1994). Dissipation of fluometuron and trifluralin residues after long-term use. *Weed Science* 42(3): 438-445. {a} Valent U.S.A. Corp., P.O. Box 5008, Greenville, MS 38704, USA

Corpas, F. J., J. Bunkelmann, et al. (1994). Identification and immunochemical characterization of a family of peroxisome membrane proteins (PMPs) in oilseed glyoxysomes. *European Journal of Cell Biology* 65(2): 280-290. {a} Dep. Botany, Arizona State Univ., Tempe, AZ 85287-1601, USA

Cotty, P. J. (1994). Influence of field application of an atoxigenic strain of *Aspergillus flavus* on the populations of *A. flavus* infecting cotton bolls and on the aflatoxin content of cottonseed. *Phytopathology* 84(11): 1270-1277. Southern Reg. Res. Cent., Agric. Res. Serv., U.S. Dep. Agric., P.O. Box 19687, New Orleans, LA 70179, USA

Davidonis, G. and O. Hinojosa (1994). Influence of seed location on cotton fiber development in planta and in vitro. *Plant Science Limerick* 103(1): 107-113. {a} USDA, ARS, Southern Reg. Res. Cent., P.O. Box 19687, New Orleans, LA 70179-0687, USA

Dixon, D. C., R. W. Seagull, et al. (1994). Changes in the accumulation of alpha- and beta-tubulin isotypes during cotton fiber development. *Plant Physiology Rockville* 105(4): 1347-1353. {a} United States Dep. Agriculture, Agricultural Res. Serv., Southern Regional Res. Cent., P.O. Box 19687, New Orleans, LA 70179, USA

Du, Y. J., F. S. Yan, et al. (1994). Olfaction in host plant selection of the soybean aphid *Aphis glycines*. *Acta Entomologica Sinica* 37(4): 385-392. Inst. Zoology, Academia Sinica, Beijing 100080, China

Dubery, I. A. and F. Smit (1994). Phenylalanine ammonia-lyase from cotton (*Gossypium hirsutum*) hypocotyls: Properties of the enzyme induced by a *Verticillium dahliae* phytotoxin. *Biochimica et Biophysica Acta* 1207(1): 24-30. {a} Dep. Chem. Biochem., RAU-University, P.O. Box 524, Auckland Park 2006, South Africa

Dugas, W. A., M. L. Heuer, et al. (1994). Sap flow measurements of transpiration from cotton grown under ambient and enriched CO₂ concentrations. *Agricultural and Forest Meteorology* 70(1-4): 231-245. {a} Blackland Res. Cent., Texas Agric. Exp. Stn., 808 E. Blackland Rd., Temple, TX 76502, USA

Durant, J. A., M. E. Roof, et al. (1994). Early season incidence of thrips (Thysanoptera) on wheat, cotton, and three wild host plant species in South Carolina. *Journal of Agricultural Entomology* 11(1): 61-71. {a} Dep. Entomology, Clemson University, Pee Dee Research and Education Center, Florence, SC 29501-9603, USA

Duttamajumder, S. K. and J. P. Verma (1994). Population dynamics of virulence genes of *Xanthomonas campestris* pv. *malvacearum* on cotton. *Journal of Phytopathology Berlin* 140(2): 145-152. Indian Inst. Sugarcane Res., Lucknow U.P.-226 002, India

Echekwu, C. A. and S. O. Alabi (1994). A Diallel Analysis of Earliness in Interspecific Crosses of Cotton. *Discovery and Innovation* 6(3): 241-243. Inst. Agric. Res., Ahmadu Bello Univ., PMB 1044 Samaru, Zaria, Nigeria

Edmisten, K. L., C. W. Wood, et al. (1994). Effects of early-season foliar fertilization on cotton growth, yield and nutrient concentration. *Journal of Plant Nutrition* 17(4): 683-692. {a} Crop Sci. Dep., North Carolina State Univ., Raleigh, NC 27695-7620, USA

Ferguson, D. L. and J. J. Burke (1994). Methionyl sulfoxide content and protein-methionine-S-oxide reductase activity in response to water deficits or high temperature. *Physiologia Plantarum* 90(2): 253-258. {a} USDA, Cropping Systems Res. Lab., Route 3, Box 215, Lubbock, TX 79401, USA

Flint, H. M., F. D. Wilson, et al. (1994). The effect of plant water stress on beneficial and pest insects including the pink bollworm and the sweetpotato whitefly in two short-season cultivars of cotton. *Southwestern Entomologist* 19(1): 11-22. USDA-ARS, Western Cotton Res. Lab., 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

Foster, J. D., Jr., L. M. Verhalen, et al. (1994). Prometryn tolerance in glanded versus glandless isolines of cotton. *Crop Science* 34(1): 67-71. {a} Merck Sharp Dohme, 12700 W. 117th St., Overland Park, KS 66210, USA

Fuzatto, M. G., E. Cia, et al. (1994). Yield stability of cotton genotypes facing the occurrence of disease and nematodes. *Bragantia* 53(1): 47-52. Secao Algodao, Inst. Agron., C.P. 28, 13001-970 Campinas, SP, Brazil

Gazaway, W. S. and B. Armstrong (1994). First report of Columbia lance nematode (*Hoplolaimus columbus*) on cotton in Alabama. *Plant Disease* 78(6): 640. Dep. Plant Pathol., Auburn Univ., Auburn, AL 36849, USA

Gerik, T. J., B. S. Jackson, et al. (1994). Plant nitrogen status and boll load of cotton. *Agronomy Journal* 86(3): 514-518. {a} Blackland Res. Cent., Texas A and M Univ. System, 808 Blackland Rd., Temple, TX 76502, USA

Gilmore, A. M. and O. Bjorkman (1994). Adenine nucleotides and the xanthophyll cycle in leaves: I. Effects of CO₂- and temperature-limited photosynthesis on adenylate energy charge and violaxanthin de-epoxidation. *Planta Heidelberg* 192(4): 526-536. {a} Dep. Plant Biol., Carnegie Inst. Washington, 290 Panama St., Stanford, CA 94305, USA

Gilmore, A. M. and O. Bjorkman (1994). Adenine nucleotides and the xanthophyll cycle in leaves: II. Comparison of the effects of CO₂- and temperature-limited photosynthesis on photosystem II fluorescence quenching, the adenylate energy charge and violaxanthin de-epoxidation in cotton. *Planta Heidelberg* 192(4): 537-544. {a} Dep. Plant Biol., Carnegie Inst. Washington, 290 Panama St., Stanford, CA 94305, USA

Giri, A. N., D. G. Giri, et al. (1994). Performance of cotton (*Gossypium* species) genotypes under different levels of nitrogen. *Indian Journal of Agronomy* 39(3): 432-436. {a} All-India Co-ordinated Res. Project on Weed Control, Maharashtra 431 402, India

- Gossett, D. R., E. P. Millhollon, et al. (1994). Antioxidant response to NaCl stress in salt-tolerant and salt-sensitive cultivars of cotton. *Crop Science* 34(3): 706-714. {a} Dep. Biol. Sci., Louisiana State Univ., Shreveport, Shreveport, LA 71115, USA
- Gossett, D. R., E. P. Millhollon, et al. (1994). The effects of NaCl on antioxidant enzyme activities in callus tissue of salt-tolerant and salt-sensitive cotton cultivars (*Gossypium hirsutum* L.). *Plant Cell Reports* 13(9): 498-503. {a} Dep. Biol. Sci., La. State Univ., Shreveport, LA 71115, USA
- Goswami, C. L. (1994). Axial control of pigments accumulation in cotyledonary leaf of cotton. *Indian Journal of Plant Physiology* 37(2): 111-112. Dep. Bot., CCS Haryana Agric. Univ., Hisar 125 044, India
- Gouia, H., M. H. Ghorbal, et al. (1994). Effects of NaCl on flows of N and mineral ions and on NO-3-reduction rate within whole plants of salt-sensitive bean and salt-tolerant cotton. *Plant Physiology Rockville* 105(4): 1409-1418. {a} Physiologie Vegetale, Dep. Sciences Biologiques, Faculte Sciences Tunis, Campus Univ., Tunisia
- Goyal, A., S. Roy, et al. (1994). Micropropagation of *Gossypium hirsutum* variety L.H.900. *Journal of Phytological Research* 7(2): 143-146. Dep. Botany, Univ. Rajasthan, Jaipur 302004, India
- Gridi, P. I. L., E. Cia, et al. (1994). Cotton improvement for multi-resistance to diseases and pests in field conditions. *Bragantia* 53(1): 33-45. {a} Secao Algodao, Inst. Agronomico, C.P. 28, 13001-970 Campinas, SP, Brazil
- Guo, C., D. M. Oosterhuis, et al. (1994). Enhancing mineral nutrient uptake of cotton plants with plant growth regulators. *Arkansas Agricultural Experiment Station Research Series*(436): 83-87. Agron. Dep., Fayetteville, AR, USA
- Hadas, A., E. Rawitz, et al. (1994). Short-term variations of soil physical properties as a function of the amount and C/N ratio of decomposing cotton residues: I. Soil aggregation and aggregate tensile strength. *Soil and Tillage Research* 32(2-3): 183-198. {a} Inst. Soils and Water, Agric. Res. Organ., Volcani Cent., Bet Dagan 50250, Israel
- Hameed, S., S. Khalid, et al. (1994). Cotton leaf curl disease in Pakistan caused by a whitefly transmitted geminivirus. *Plant Disease* 78(5): 529. {a} Plant Virol. Lab., Natl. Agric. Res. Centre, Islamabad-45500, Pakistan
- Hansen, G., A. Das, et al. (1994). Constitutive expression of the virulence genes improves the efficiency of plant transformation by *Agrobacterium*. *Proceedings of the National Academy of Sciences of the United States of America* 91(16): 7603-7607. {a} Ciba-Geigy Corporation, PO Box 12257, Research Triangle Park, NC 27709, USA
- Hashmi, R. A., N. Bano, et al. (1994). Die lectric properties of leaves from some plant species. *Pakistan Journal of Botany* 26(2): 467-475. {a} Condensed Matter Res. Lab., Dep. Physics, Univ. Karachi, Karachi 75270, Pakistan
- He, J. and C. Sun (1994). A scheme for introgression of delayed gland morphogenesis gene from wild *Gossypium bickii* into cultivated upland cotton (*G. hirsutum*). *Acta Genetica Sinica* 21(1): 52-58. Inst. Genetics, Academia Sinica, Beijing 100101, China
- Hedin, P. A. and J. C. McCarty, Jr. (1994). Multiyear study of the effects of kinetin and other plant growth hormones on yield, agronomic traits, and allelochemicals of cotton. *Journal of Agricultural and Food Chemistry* 42(10): 2305-2307. {a} Crop Sci. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Mississippi State, MS 39762-5367, USA
- Hedin, P. A. and J. C. McCarty, Jr. (1994). Effects of several commercial plant growth regulator formulations on yield and allelochemicals of cotton (*Gossypium hirsutum* L.). *Journal of Agricultural and Food Chemistry* 42(6): 1355-1357. {a} Crop Sci. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Mississippi State, MS 39762-5367, USA

- Heitholt, J. J. (1994). Comparison of adjuvant effects on cotton leaf potassium concentration and lint yield. *Journal of Plant Nutrition* 17(12): 2221-2233. USDA-ARS, Cotton Physiology Genetics, P.O. Box 345, Stoneville, MS 38776, USA
- Heitholt, J. J. (1994). Canopy characteristics associated with deficient and excessive cotton plant population densities. *Crop Science* 34(5): 1291-1297. USDA-ARS, Cotton Physiol. Genet., P.O. Box 345, Stoneville, MS 38776, USA
- Heitholt, J. J. (1994). Supplemental boron, boll retention percentage, ovary carbohydrates, and lint yield in modern cotton genotypes. *Agronomy Journal* 86(3): 492-497. USDA-ARS, Cotton Physiol. and Genetics, P.O. Box 345, Stoneville, MS 38776, USA
- Heitholt, J. J. (1994). Effects of foliar urea- and triazone-nitrogen, with and without boron, on cotton. *Journal of Plant Nutrition* 17(1): 57-70. USDA-ARS, Cotton Physiology Genetics, P.O. Box 345, Stoneville, MS 38776, USA
- Heitholt, J. J. and J. H. Schmidt (1994). Receptacle and ovary assimilate concentrations and subsequent boll retention in cotton. *Crop Science* 34(1): 125-131. {a} USDA-ARS, Cotton Physiology Genetics, P.O. Box 345, Stoneville, MS 38776, USA
- Hileman, D. R., G. Huluka, et al. (1994). Canopy photosynthesis and transpiration of field-grown cotton exposed to free-air CO₂ enrichment (FACE) and differential irrigation. *Agricultural and Forest Meteorology* 70(1-4): 189-207. {a} Dep. Biol., Tuskegee Univ., Tuskegee, AL 36088, USA
- Hira, G. S. (1994). Effect of water-table depth on yield of cotton (*Gossypium* species). *Indian Journal of Agricultural Sciences* 64(8): 550-551. Dep. Soils, Punjab Agric. Univ., Ludhiana 141 004, India
- Hirimburegama, K. and N. Gamage (1994). In vitro callus and cell cultures of *Gossypium hirsutum* L. (Cotton). *Journal of the National Science Council of Sri Lanka* 22(4): 305-312. Dep. Bot., Univ. Colombo, PO Box 1490, Colombo, Sri Lanka
- Holt, S. J., J. M. Stewart, et al. (1994). Flower bud development in greenhouse-grown cotton. *Crop Science* 34(4): 973-976. {a} Dep. Agron., Univ. Ark., Fayetteville, AR 72701, USA
- Holtz, B. A., A. E. Karu, et al. (1994). Enzyme-linked immunosorbent assay for detection of *Thielaviopsis basicola*. *Phytopathology* 84(9): 977-983. {a} Dep. Plant Pathol., Univ. Calif. Davis, Kearney Agric. Cent., 9240 S. Riverbend Ave., Parlier, CA 93648, USA
- Holtz, B. A. and A. R. Weinhold (1994). *Thielaviopsis basicola* in San Joaquin Valley soils and the relationship between inoculum density and disease severity of cotton seedlings. *Plant Disease* 78(10): 986-990. {a} Dep. Plant Pathol., Univ. Calif. Davis, Kearney Agric. Cent., 9240 S. Riverbend Ave., Parlier, CA 93648, USA
- Huluka, G., D. R. Hileman, et al. (1994). Effects of elevated CO₂ water stress on mineral concentration of cotton. *Agricultural and Forest Meteorology* 70(1-4): 141-152. {a} Dep. Agric. Sci., Tuskegee Univ., Tuskegee, AL 36088, USA
- Hunsaker, D. J., G. R. Hendrey, et al. (1994). Cotton evapotranspiration under field conditions with CO₂ enrichment and variable soil moisture regimes. *Agricultural and Forest Meteorology* 70(1-4): 247-258. {a} USDA, ARS, US Water Conservation Lab., 4331 Broadway Rd., Phoenix, AZ 85040, USA
- Iqbal, R. M. S., M. B. Chaudhry, et al. (1994). Development of a high yielding cotton mutant, NIAB-92 through the use of induced mutations. *Pakistan Journal of Botany* 26(1): 99-104. Nuclear Inst. Agric. Biol., Faisalabad, Pakistan
- Jacobs, K. A., S. E. Eentz, et al. (1994). Screening three *Cercis* species for susceptibility to *Verticillium dahliae*. *Plant Disease* 78(9): 925. U.S. Natl. Arboretum, 3501 New York Ave. NE, Washington, DC 20002-1958, USA

Jacobson, B. D., D. S. Murray, et al. (1994). Soil-water extraction profiles of cotton (*Gossypium hirsutum*) and weed species. *Weed Technology* 8(2): 190-198. {a} Agron. Dep., Univ. Kentucky, Lexington, KY 40546, USA

Jafri, A. Z. and R. Ahmad (1994). Plant growth and ionic distribution in cotton (*Gossypium hirsutum* L.) under saline environment. *Pakistan Journal of Botany* 26(1): 105-114. {a} Dep. Bot., Islamia Degree Coll., Multan, Pakistan

Jakobsen, K. S., D. W. Hughes, et al. (1994). Simultaneous induction of postabscission and germination mRNAs in cultured dicotyledonous embryos. *Planta Heidelberg* 192(3): 384-394. {a} Dep. Botany, Univ. Georgia, Athens, GA 30602-7271, USA

Janes, L. D., D. M. Oosterhuis, et al. (1994). Foliar-applied potassium nitrate effects on cotton genotypes. *Arkansas Agricultural Experiment Station Research Series*(436): 51-53. {a} Agron. Dep., Fayetteville, AR, USA

Jenkins, J. N. and J. C. McCarty, Jr. (1994). Comparison of four cotton genotypes for resistance of *Heliothis virescens*. *Crop Science* 34(5): 1231-1233. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA

John, M. E. and M. W. Petersen (1994). Cotton (*Gossypium hirsutum* L.) pollen-specific polygalacturonase mRNA: Tissue and temporal specificity of its promoter in transgenic tobacco. *Plant Molecular Biology* 26(6): 1989-1993. {a} Agracetus Inc., 8520 University Green, Middleton, WI 53562, USA

Johnson, W. C., III, J. Cardina, et al. (1994). Dynamics of subeconomic threshold populations of sicklepod (*Cassia obtusifolia*) in a peanut-cotton-corn rotation. *Weed Science* 42(3): 364-368. {a} Res. Agron., USDA-ARS, For. Res. Agron., Agric. Res. Statistician, Coastal Plain Exp. Stn., Tifton, GA 31793-0748, USA

Kadas, I., G. Arvai, et al. (1994). A simple method for the preparation of various 1-oxo-hydrindene-2-acetic and -propionic acids: Valuable precursors of strigol and its analogues. *Tetrahedron* 50(9): 2895-2906. {a} Res. Group, Hung. Acad. Sci., Dep. Organic Chem. Technology, Technical Univ. Budapest, P.O.B. 91, H-1521 Budapest, Hungary

Karunyal, S., G. Renuga, et al. (1994). Effects of tannery effluent on seed germination, leaf area, biomass and mineral content of some plants. *Bioresource Technology* 47(3): 215-218. {a} Dep. Plant Sci., Sch. Biol. Sci., Madurai Kamaraj Univ., Madurai-625 021, India

Kasperbauer, M. J. (1994). Cotton plant size and fiber developmental responses to FR/R ratio reflected from the soil surface. *Physiologia Plantarum* 91(2): 317-321. Coastal Plains Res. Cent., Agric. Res. Serv., US Dep. Agric., Florence, SC 29502-3039, USA

Kaur, R. and O. S. Singh (1994). Effect of seedling hardening on early growth and water relations of cotton (*Gossypium hirsutum* (L.) American cotton). *Indian Journal of Agricultural Research* 28(4): 235-238. {a} Dep. Biotechnol., Univ. Hortic. For., Solan-173 230, India

Keese, R. J. and N. D. Camper (1994). Uptake and translocation of (14C)MSMA in cotton and MSMA-resistant and -susceptible cocklebur. *Pesticide Biochemistry and Physiology* 49(2): 138-145. Dep. Plant Pathol. Physiol., Clemson Univ., Box 340377, Clemson, SC 29634-0377, USA

Keisling, T. C. and C. W. Smith (1994). Research plot lengths for spindle harvested cotton. *Communications in Soil Science and Plant Analysis* 25(17-18): 2981-2988. {a} Univ. Arkansas, NEREC, P.O. Box 48, Keiser, AR 72351, USA

Khah, E. M. and H. C. Passam (1994). Sensitivity of seed of cotton cv. Zeta-2 to damage during acid-delinting. *Plant Varieties and Seeds* 7(1): 51-57. {a} Dep. Horticulture, Agric. Univ. Athens, 75 Iera Odos, Votanikos, Athens 118 55, Greece

- Khoshkhoo, N., P. A. Hedin, et al. (1994). Free sugars in root-knot nematode susceptible and resistant cotton plant roots and leaves. *Journal of Agricultural and Food Chemistry* 42(2): 366-368. {a} Crop Sci. Res. Lab., Agric. Res. Serv., US Dep. Agric., Mississippi State, MS 39762-5367, USA
- Khoshkhoo, N., P. A. Hedin, et al. (1994). Terpenoid aldehydes in root-knot nematode susceptible and resistant cotton plants. *Journal of Agricultural and Food Chemistry* 42(1): 204-208. {a} Crop Sci. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Mississippi State, MS 39762-5367, USA
- Kimball, B. A., R. L. Lamorte, et al. (1994). Effects of free-air CO₂ enrichment on energy balance and evapotranspiration of cotton. *Agricultural and Forest Meteorology* 70(1-4): 259-278. {a} USDA, ARS, US Water Conservation Lab., 4331 E. Broadway Rd., Phoenix, AZ 85040, USA
- Kinloch, R. A. and R. K. Sprenkel (1994). Plant-parasitic nematodes associated with cotton in Florida. *Journal of Nematology* 26(4 Suppl.): 749-752. Univ. Fla., Agric. Res. Educ. Cent., Jay, FL 32565, USA
- Kleifeld, Y., B. Rubin, et al. (1994). Soil-incorporated trifluralin protects cotton (*Gossypium hirsutum*) from root-absorbed herbicides. *Weed Research* 34(6): 461-469. {a} Dep. Weed Sci., A.R.O. Newe Ya'ar Exper. Stn., P.O. Haifa 31999, Israel
- Klingaman, T. E. and L. R. Oliver (1994). Influence of cotton (*Gossypium hirsutum*) and soybean (*Glycine max*) planting date on weed interference. *Weed Science* 42(1): 61-65. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA
- Korn, R. W. (1994). Pattern formation in plant epidermis through inhibition of immediately adjacent cells by pattern elements. *Protoplasma* 180(3-4): 145-152. Biol. Dep., Bellarmine Coll., Louisville, KY 40205, USA
- Ladygin, V. G., I. D. Shegai, et al. (1994). Spectral characteristics and photochemical activity of leaves in "xantha" cotton mutants. *Fiziologiya Rastanii Moscow* 41(1): 10-16. {a} Inst. Soil Sci. Photosynth., Russ. Acad. Sci., 142292 Pushchino, Russia
- Lakkineni, K. C., S. N. Bhardwaj, et al. (1994). Effect of temperature on early growth and seed-cotton yield in upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 64(9): 653-654. {a} Div. Plant Physiol., Indian Agric. Res. Inst., New Delhi 110 012, India
- Landolt, P. J., R. R. Heath, et al. (1994). Effects of host plant, *Gossypium hirsutum* L., on sexual attraction of cabbage looper moths, *Trichoplusia ni* (Hubner) (Lepidoptera: Noctuidae). *Journal of Chemical Ecology* 20(11): 2959-2974. {a} US Dep. Agric., Agric. Res. Serv., Insect Attractants Behavior Basic Biol. Res. Lab., P.O. Box 14565, 1700 S.W. 23rd Dr., Gainesville, FL 32604, USA
- Lascano, R. J., R. L. Baumhardt, et al. (1994). Soil and plant water evaporation from strip-tilled cotton: Measurement and simulation. *Agronomy Journal* 86(6): 987-994. {a} Texas Agric. Exp. Stn., Rt. 3, Box 219, Lubbock, TX 79401-9757, USA
- Liu, R. J. and X. S. Luo (1994). A new method to quantify the inoculum potential of arbuscular mycorrhizal fungi. *New Phytologist* 128(1): 89-92. {a} Dep. Horticulture, Lai-Yang Agric. Coll., Lai-Yang, Shandong 265200, China
- Loughrin, J. H., A. Manukian, et al. (1994). Diurnal cycle of emission of induced volatile terpenoids by herbivore-injured cotton plants. *Proceedings of the National Academy of Sciences of the United States of America* 91(25): 11836-11840. {a} Insect Attractants, Behavior Basic Biol. Research Lab., U.S. Dep. Agric., Agric. Research Serv., 1700 Southwest 23rd Drive, Gainesville, FL 32604, USA
- Mahmoud, S. M. M., M. M. Rabag, et al. (1994). Some aspects affecting preplanting management of cotton seedling disease caused by *Rhizoctonia solani*. *Egyptian Journal of Phytopathology* 22(1): 39-57. {a} Res. Inst., Agric. Res. Centre, Giza, Egypt

Manoharan, M., K. H. Doddayenkanna, et al. (1994). Histology of pollen degeneration in cytoplasmic male sterile (CMS) cotton (*Gossypium hirsutum* L.). Proceedings of the Indian National Science Academy Part B Biological Sciences 60(6): 565-570. {a} Dep. Botany, Karnatak Univ., Dharwad 580 003, India

Manoharan, T. and S. Uthamasamy (1994). Differential susceptibility of field population of gram podborer (*Helicoverpa armigera*) to insecticides in Tamil Nadu. Indian Journal of Agricultural Sciences 64(2): 126-131. Dep. Agric. Entomology, Tamil Nadu Agric. Univ., Coimbatore 641 003, India

Maredia, K. M., N. P. Tugwell, et al. (1994). Technique for screening cotton germplasm for resistance to tarnished plant bug, *Lygus lineolaris* (Palisot de beauvois). Southwestern Entomologist 19(1): 63-70. {a} Dep. Entomol., Mich. State Univ., East Lansing, MI 48824, USA

Martinez, V. and A. Lauchli (1994). Salt-induced inhibition of phosphate uptake in plants of cotton (*Gossypium hirsutum* L.). New Phytologist 126(4): 609-614. {a} Cent. Edafol. Biol. Aplicada del Segura, CSIC, Apdo 4195, 30080 Murcia, Spain

Marur, C. J., L. Sodek, et al. (1994). Free amino acids in leaves of cotton plants under water deficit. Revista Brasileira de Fisiologia Vegetal 6(2): 103-108. {a} Eng. Agric., Dr., IAPAR. CP. 1331, Londrina, PR 86001-970, Brazil

Mauney, J. R., B. A. Kimball, et al. (1994). Growth and yield of cotton in response to a free-air carbon dioxide enrichment (FACE) environment. Agricultural and Forest Meteorology 70(1-4): 49-67. {a} USDA ARS, Western Cotton Res. Lab., 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

May, O. L. and C. C. Green (1994). Genetic variation for fiber properties in elite Pee Dee cotton populations. Crop Science 34(3): 684-690. {a} USDA-ARS, Coastal Plains Soil, Water, and Plant Res. Cent., P.O. Box 3039, Florence, SC 29502-3039, USA

McCall, P. J., T. C. J. Turlings, et al. (1994). Herbivore-induced volatile emissions from cotton (*Gossypium hirsutum* L.) seedlings. Journal of Chemical Ecology 20(12): 3039-3050. {a} Insect Attractants, Behavior Basic Biology Res. Lab., Agricultural Res. Serv., U.S. Dep. Agriculture, P.O. Box 14565, Gainesville, FL 32604, USA

McCarty, J. C., Jr. and P. A. Hedin (1994). Effects of 1,1-dimethylpiperidinium chloride on the yields, agronomic traits, and allelochemicals of cotton (*Gossypium hirsutum* L.), a nine year study. Journal of Agricultural and Food Chemistry 42(10): 2302-2304. {a} Crop Sci. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Mississippi State, MS 39762-5367, USA

McConnell, J. S., W. H. Baker, et al. (1994). Response of cotton grown on compacted soils to phosphorus and potassium fertilization and deep tillage. Arkansas Agricultural Experiment Station Research Series(436): 44-50. {a} Southeast Res. and Extension Cent., Monticello, AR, USA

McCully, M. E. (1994). Accumulation of high levels of potassium in the developing xylem elements in roots of soybean and some other dicotyledons. Protoplasma 183(1-4): 116-125. Dep. Biol., Carleton Univ., 1125 Colonel By Drive, Ottawa, ON K1S 5B6, Canada

McDonald, M. B., L. J. Elliot, et al. (1994). DNA extraction from dry seeds for RAPD analyses in varietal identification studies. Seed Science and Technology 22(1): 171-176. Dep. Agron., The Ohio State Univ., Columbus, OH 43210, USA

McKinion, J. M. and R. A. Sequeira (1994). Mechanics of model building. Hortscience 29(12): 1413-1415. Crop Simulation Res. Unit, Agric. Res. Serv., Miss. State Univ., MS 39762-5367, USA

McMichael, B. L. and J. J. Burke (1994). Metabolic activity of cotton roots in response to temperature. Environmental and Experimental Botany 34(2): 201-206. USDA-ARS, Cropping Systems Res. Lab., Plant Stress and Water Conservation Res. Unit, Route 3, Box 215, Lubbock, TX 79401, USA

McSorley, R., D. W. Dickson, et al. (1994). Host status of selected tropical rotation crops to four populations of root-knot nematodes. *Nematropica* 24(1): 45-53. {a} Dep. Entomology Nematology, P.O. Box 110620, Univ. Fla., Gainesville, FL 32611, USA

McSorley, R., D. W. Dickson, et al. (1994). Effects of tropical rotation crops on *Meloidogyne arenaria* population densities and vegetable yields in microplots. *Journal of Nematology* 26(2): 175-181. {a} Entomol. and Nematol. Dep., P.O. Box 110620, Univ. Florida, Gainesville, FL 32611-0611, USA

Medlin, C. R. (1994). History, life cycle, and control of the cotton bollworm. *Journal of Natural Resources and Life Sciences Education* 23(1): 70-72. C/O Kevin J. Donnelly, Dep. Agronomy, Oklahoma State Univ., Stillwater, OK 74078, USA

Meena, R., K. Rathinavel, et al. (1994). Seed development and maturation in upland (*Gossypium hirsutum*) and tree (*G. arboreum*) cottons. *Indian Journal of Agricultural Sciences* 64(2): 111-113. {a} Regional Station, CICR, Sirsa, Haryana 125 055, India

Merrill, S. D. and D. R. Upchurch (1994). Converting root numbers observed at minirhizotrons to equivalent root length density. *Soil Science Society of America Journal* 58(4): 1061-1067. {a} USDA-ARS, P.O. Box 459, Mandan, ND 58554, USA

Meshram, L. D., R. A. Ghongade, et al. (1994). Development of male sterile systems from various sources in cotton (*Gossypium* spp.). *PKV Research Journal* 18(1): 83-86. {a} PKV, Akola, India

Meshram, L. D. and M. A. Tayyab (1994). Interspecific hybrid between *Gossypium hirsutum* L. x *Gossypium australe* F. Muell. *PKV Research Journal* 18(1): 124-126. Cotton Res. Unit, CRS, Punjabrao Krishi Vidyapeeth, Akola, India

Meshram, L. D. and M. A. Tayyab (1994). Cytomorphology of two interspecific hybrids of *Gossypium*. *PKV Research Journal* 18(1): 63-67. PKV, Akola, India

Miley, W. N., D. M. Oosterhuis, et al. (1994). Effects of foliar application of five potassium fertilizers on cotton yield and quality. *Arkansas Agricultural Experiment Station Research Series*(436): 62-66. {a} North Little Rock, AR, USA

Millar, A. A., M. R. Olive, et al. (1994). The expression and anaerobic induction of alcohol dehydrogenase in cotton. *Biochemical Genetics* 32(7-8): 279-300. {a} Dep. Botany, Univ. British Columbia, 3529-6270 University Blvd., Vancouver, BC V6T 1Z4, Canada

Mitchell, C. C., A. E. Hiltbold, et al. (1994). Evaluation of a liquid potassium bicarbonate/amino acid co-product as a source of potassium, nitrogen, and sulfur. *Journal of Plant Nutrition* 17(12): 2119-2134. {a} Dep. Agronomy Soils, Ala. Agricultural Experiment Station, Auburn University, Auburn, AL 36849, USA

Montandon, R., J. E. Slosser, et al. (1994). Late-Season Termination Effects on Cotton Fruiting, Yield, and Boll Weevil (Coleoptera: Curculionidae) Damage in Texas Dryland Cotton. *Journal of Economic Entomology* 87(6): 1647-1652. Texas Agric. Exp. Stn., PO Box 1658, Vernon, TX 76385, USA

Mullins, G. L., D. W. Reeves, et al. (1994). In-row subsoiling and potassium placement effects on root growth and potassium content of cotton. *Agronomy Journal* 86(1): 136-139. {a} Dep. Agronomy and Soils, Auburn University, Auburn, AL 36849-5412, USA

Naranjo, S. E. and H. M. Flint (1994). Spatial distribution of preimaginal *Bemisia tabaci* (Homoptera: Aleyrodidae) in cotton and development of fixed-precision sequential sampling plans. *Environmental Entomology* 23(2): 254-266. Western Cotton Res. Lab., USDA-ARS, 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

Novo, J. P. S. and D. Gabriel (1994). Occurrence of cotton pests in cotton gins in the region of campinas, Sao Paulo State. *Anais da Sociedade Entomologica do Brasil* 23(3): 449-453. {a} Inst. Agronomico Campinas, Secao Entomol., Caixa postal 28, 13001-970 Campinas, SP, Brazil

- Oosterhuis, D. M. and L. D. Janes (1994). Foliar fertilization of cotton with potassium in Arkansas 1989-1993. Arkansas Agricultural Experiment Station Research Series(436): 58-61. Agron. Dep., Fayetteville, AR, USA
- Oosterhuis, D. M. and D. Zhao (1994). Increased root length and branching in cotton by soil application of the plant growth regulator PGR-IV. *Plant and Soil* 167(1): 51-56. Dep. Agronomy, Univ. Arkansas, Fayetteville, AR 72703, USA
- Pachepsky, L. B., V. R. Reddy, et al. (1994). Cotton canopy photosynthesis model for predicting the effect of temperature and elevated carbon dioxide concentration. *Biotronics* 23: 35-46. {a} Duke Univ. Phytotron, Dep. Botany, Duke Univ., Science Drive, Durham, NC 27708, USA
- Page, S. L. J. and J. Bridge (1994). The African cotton-root nematode, *Meloidogyne acrona*; its pathogenicity and intra-generic infectivity within *Gossypium*. *Fundamental and Applied Nematology* 17(1): 67-73. International Inst. Parasitol., 395 A Hatfield Road, St. Albans AL4 0XU, UK
- Paramasivam, S. and G. A. Breitenbeck (1994). Distribution of nitrogen in soils of the southern Mississippi River alluvial plain. *Communications in Soil Science and Plant Analysis* 25(3-4): 247-267. Dep. Agron., Louisiana Agric. Exp. Station, LSU Agric. Cent., Baton Rouge, LA 70803-2110, USA
- Park, Y. H. and R. J. Kohel (1994). Effect of concentration of MgCl₂ on random-amplified DNA polymorphism. *Biotechniques* 16(4): 652-654, 656. {a} USDA-ARS, Southern Crops Res. Lab., College Station, TX 77845, USA
- Peeters, M. C., K. Willems, et al. (1994). Protoplast-to-plant regeneration in cotton (*Gossypium hirsutum* L. cv. Coker 312) using feeder layers. *Plant Cell Reports* 13(3-4): 208-211. {a} Loab. Tropical Crop Husbandry, Catholic Univ. Leuven, Kardinaal Mercierlaan 92, B-3001 Heverlee, Belgium
- Pettigrew, W. T. (1994). Source-to-sink manipulation effects on cotton lint yield and yield components. *Agronomy Journal* 86(4): 731-735. USDA-ARS, Cotton Physiol. and Genet. Res., P.O. Box 345, Stoneville, MS 38776, USA
- Pettigrew, W. T. and W. R. Meredith, Jr. (1994). Leaf gas exchange parameters vary among cotton genotypes. *Crop Science* 34(3): 700-705. {a} Cotton Physiol. and Genetics Res., P.O. Box 345, Stoneville, MS 38776, USA
- Pinter, P. J., Jr., B. A. Kimball, et al. (1994). Effects of free-air carbon dioxide enrichment on PAR absorption and conversion efficiency by cotton. *Agricultural and Forest Meteorology* 70(1-4): 209-230. {a} US Water Conservation Lab., 4331 E. Broadway Rd., Phoenix, AZ 85040, USA
- Porteous, L. A., J. L. Armstonrg, et al. (1994). An effective method to extract DNA from environmental samples for polymerase chain reaction amplification and DNA fingerprint analysis. *Current Microbiology* 29(5): 301-307. {a} Biotechnology Program, United States Environmental Protection Agency, Environmental Res. Lab., 200 S.W. 35 Street, Corvallis, OR 97333, USA
- Poswal, M. A. T. (1994). Variation for resistance to bacterial blight among different hybrid pools of cotton in Nigeria. *Acta Phytopathologica et Entomologica Hungarica* 29(1-2): 81-88. Dep. Crop Protection, Inst. Agric. Res., Ahmadu Bello Univ., Samaru-Zaria, Nigeria
- Prasad, M. and R. Prasad (1994). Response of upland cotton (*Gossypium hirsutum*) to micro-nutrients and sulphur. *Indian Journal of Agronomy* 39(4): 707-708. Div. Agron., Indian Agric. Res. Inst., New Delhi 110 012, India
- Prasad, M. and R. Prasad (1994). Response of upland cotton (*Gossypium hirsutum*) to biofertilizer and nitrogen fertilization. *Indian Journal of Agronomy* 39(2): 334-336. Div. Agronomy, Indiana Agric. Res. Inst., New Delhi 110 012, India

- Prior, S. A., H. H. Rogers, et al. (1994). Free-air CO₂ enrichment of cotton: Vertical and lateral root distribution patterns. *Plant and Soil* 165(1): 33-44. {a} Natl. Soil Dynamics Lab., Agric. Res. Serv., U.S. Dep. Agric., P.O. Box 3439, Auburn, AL 36831-3439, USA
- Prior, S. A., H. H. Rogers, et al. (1994). Effects of free-air CO₂ enrichment on cotton root growth. *Agricultural and Forest Meteorology* 70(1-4): 69-86. {a} Natl. Soil Dynamics Lab., Agric. Res. Serv., US Dep. Agric., PO Box 3439, Auburn, AL 36831-3439, USA
- Puri, S., S. Singh, et al. (1994). Growth and productivity of crops in association with an *Acacia nilotica* tree belt. *Journal of Arid Environments* 27(1): 37-48. Dep. Forestry, Haryana Agric. Univ., Hisar 125 004, India
- Quisenberry, J. E., L. D. McDonald, et al. (1994). Responses of photosynthetic rates to genotypic differences in sink-to-source ratios in upland cotton (*Gossypium hirsutum* L.). *Environmental and Experimental Botany* 34(3): 245-252. {a} Plant Stress and Water Conservation Unit, Route 3, Box 215, Lubbock, TX 79401, USA
- Rasooly, R., B. Raccah, et al. (1994). Laboratory transmission of the citrus stubborn disease agent by a leafhopper from the *Circulifer tenellus* complex from the Jordan Valley. *Phytoparasitica* 22(3): 209-218. {a} Dep. Virology, ARO, Volcani Cent., Bet Dagan 50250, Israel
- Rawitz, E., A. Hadas, et al. (1994). Short-term variations of soil physical properties as a function of the amounts and C/N ratio of decomposing cotton residues: II. Soil compressibility, water retention and hydraulic conductivity. *Soil and Tillage Research* 32(2-3): 199-212. {a} Seagram Cent. Soil and Water Sci., Fac. Agric., Hebrew Univ., Rehovot 76100, Israel
- Reddy, V. R. (1994). Modeling cotton growth and phenology in response to temperature. *Computers and Electronics in Agriculture* 10(1): 63-73. USDA:ARS:BA:NRI:SRL, Systems Research Lab., USDA Agricultural Research Service, Build. 007, Room 008, BARC-West, Beltsville, MD 20705, USA
- Reddy, V. R., L. B. Pachepsky, et al. (1994). Response of crop photosynthesis to carbon dioxide, temperature, and light: Experimentation and modeling. *Hortscience* 29(12): 1415-1422. U.S. Dep. Agric., Agric. Res. Serv., Natural Resources Inst., Systems Res. Lab., Build. 007, Room 008, Beltsville Agric. Res. Cent.-West, Beltsville, MD 20705, USA
- Reddy, V. R., K. R. Reddy, et al. (1994). Carbon dioxide and temperature effects on cotton leaf initiation and development. *Biotronics* 23: 59-74. {a} Systems Res. Lab., USDA-ARS, Build. 007, Room 008, BARC-West, Beltsville, MD 20705, USA
- Reddy, V. R., K. R. Reddy, et al. (1994). Carbon dioxide enrichment and temperature effects on root growth in cotton. *Biotronics* 23: 47-57. {a} USDA, ARS, BA, NRI, Systems Res. Lab., Build 007, Room 008, BARC-West, Beltsville, MD 20705, USA
- Reinisch, A. J., J. M. Dong, et al. (1994). A detailed RFLP map of cotton, *Gossypium hirsutum* X *Gossypium barbadense*: Chromosome organization and evolution in a disomic polyploid genome. *Genetics* 138(3): 829-847. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA
- Resende, M. L. V., J. Flood, et al. (1994). Host specialization of *Verticillium dahliae*, with emphasis on isolates from cocoa (*Theobroma cacao*). *Plant Pathology Oxford* 43(1): 104-111. {a} Cocoa Res. Cent., CEPLAC, Cx. Postal 07, 45600-Itabuna-BA, Brazil
- Rodriguez, K. R., B. N. Kokalis, et al. (1994). Rotations of coastal bermudagrass, cotton, and bahiagrass for management of *Meloidogyne arenaria* and southern blight in peanut. *Journal of Nematology* 26(4 Suppl.): 665-668. {a} Dep. Plant Pathol., Auburn Univ., Auburn, AL 36849, USA
- Rufty, T. W., Jr., R. B. Thomas, et al. (1994). Growth response of cotton to CO₂ enrichment in differing light environments. *Physiologia Plantarum* 91(3): 503-509. {a} USDA ARS, Oxford, NC 27565, USA

Runion, G. B., E. A. Curl, et al. (1994). Effects of free-air CO₂ enrichment on microbial populations in the rhizosphere and phyllosphere of cotton. *Agricultural and Forest Meteorology* 70(1-4): 117-130. {a} Sch. Forestry, 108 M. White Smith Hall, Auburn Univ., Auburn, AL 36849, USA

Sabino, N. P., N. M. D. Silva, et al. (1994). Effects of urea and ammonium sulfate applications on the characteristics and technological properties of cotton fiber. *Bragantia* 53(1): 75-82. {a} Secao Tecnologia Fibras, Inst. Agronomico, C.P. 28, 13001-970 Campinas, SP, Brazil

Saha, S. and D. M. Stelly (1994). Chromosomal location of Phosphoglucomutase 7 locus in *Gossypium hirsutum*. *Journal of Heredity* 85(1): 35-39. Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA

Samora, P. J., D. M. Stelly, et al. (1994). Localization and mapping of the Le-1 and GI-2 loci of cotton (*Gossypium hirsutum* L.). *Journal of Heredity* 85(2): 152-157. {a} Dep. Soil Crop Sci., Texas Agric. Experiment Station Texas A M Univ., College Station, TX 77843-2474, USA

Sankaralingam, A. and E. C. McGawley (1994). Interrelationships of *Rotylenchulus reniformis* with *Rhizoctonia solani* on Cotton. *Journal of Nematology* 26(4): 475-485. {a} Dep. Plant Pathol., Tamil Nadu Agric. Univ., Coimbatore 641003, India

Sankaralingam, A. and E. C. McGawley (1994). Influence of *Rhizoctonia solani* on Egg Hatching and Infectivity of *Rotylenchulus reniformis*. *Journal of Nematology* 26(4): 486-491. {a} Dep. Plant Pathol., Tamil Nadu Agric. Univ., Coimbatore 641003, India

Schmutz, A., T. Jenny, et al. (1994). A caffeoyl-fatty acid-glycerol ester from wax associated with green cotton fibre suberin. *Phytochemistry Oxford* 36(6): 1343-1346. {a} Institu Botanische Biologie, Universitat Freiburg, A. Gockelstrasse 3, CH-1700 Freiburg, Switzerland

Schouest, L. P., Jr. and T. A. Miller (1994). Automated pheromone traps show male pink bollworm (Lepidoptera: Gelechiidae) mating response is dependent on weather conditions. *Journal of Economic Entomology* 87(4): 965-974. Dep. Entomol., Univ. Calif., Riverside, CA 92521, USA

Schroer, J. and P. Welzel (1994). Asymmetric synthesis of a key synthetic precursor for (+)-strigol and sorgolactone. *Tetrahedron* 50(23): 6839-6858. {a} Fakultat fuer Chemie, Ruhr-Universitaet, D-44780 Bochum, Germany

Schuster, G. L. and R. S. Halliwell (1994). Six new hosts of tomato spotted wilt virus in Texas. *Plant Disease* 78(1): 100. Dep. Plant Pathology Microbiology, Tex. Agricultural Experiment Station, College Station, TX 77843, USA

Scott, W. P., J. E. Mulrooney, et al. (1994). Mortality of susceptible and resistant *Heliothis virescens* larvae exposed to various petroleum oils and insecticide rates. *Journal of Entomological Science* 29(4): 466-470. {a} USDA-ARS, Southern Insect Management Lab., P.O. Box 346, Stoneville, MS 38776, USA

Sharma, R. K. and R. S. S. Tomar (1994). Response of upland cotton (*Gossypium hirsutum*) to plant spacing and nitrogen under rainfed conditions. *Indian Journal of Agronomy* 39(2): 274-276. {a} ECF Scheme, Zonal Agric. Res. Station, JNKVV, Moena, Madhya Pradesh 476 001, India

Shripathi, V. and G. S. Swamy (1994). Effects of triacontanol on the lipid composition of cotton (*Gossypium hirsutum* L.) leaves and its interaction with indole-3-acetic acid and benzyladenine. *Plant Growth Regulation* 14(1): 45-50. {a} Dep. Botany, Karnatak Univ., Dharwad 580 003, India

Singh, D. (1994). Effect of foliar application of lime on rainfed upland cotton (*Gossypium hirsutum*) and tree cotton (*G. arboreum*). *Indian Journal of Agricultural Sciences* 64(10): 697-700. Project Directorate Cropping Systems Res., Modipuram, Uttar Pradesh 250 110, India

Singh, T., J. S. Brar, et al. (1994). Effect of Wheat Residues, Nitrogen Levels and its Time of Application on Yield of Cotton (*Gossypium hirsutum* L.). *Annals of Biology Ludhiana* 10(2): 280-281. PAU Regional Research Station, Bathinda-151 001, India

Skinner, R. H. and A. C. Cohen (1994). Phosphorus nutrition and leaf age effects on sweetpotato whitefly (Homoptera: Aleyrodidae) host selection. *Environmental Entomology* 23(3): 693-698. West. Cotton Res. Lab., USDA-ARS, Phoenix, AZ 85040, USA

Skinner, R. H. and J. W. Radin (1994). The effect of phosphorus nutrition on water flow through the apoplastic bypass in cotton roots. *Journal of Experimental Botany* 45(273): 423-428. {a} USDA-ARS, Western Cotton Research Lab., 4135 E. Broadway Road, Phoenix, AZ 85040, USA

Smith, J. E. and D. J. Longstreth (1994). Leaf expansion and carbon assimilation in cotton leaves grown at two photosynthetic photon flux densities. *American Journal of Botany* 81(6): 711-717. {a} Dep. Botany, Louisiana State Univ., Baton Rouge, LA 70803-1705, USA

Snipes, C. E. and C. C. Baskin (1994). Influence of early defoliation on cotton yield, seed quality, and fiber properties. *Field Crops Research* 37(2): 137-143. {a} Miss. Agricultural Forestry Experiment Station, Stoneville, MS 38776, USA

Snipes, C. E. and J. D. Byrd, Jr. (1994). The influence of fluometuron and MSMA on cotton yield and fruiting characteristics. *Weed Science* 42(2): 210-215. Miss. Agric. Forest Exp. Station, P.O. Box 197, Stoneville, MS 38766, USA

Snipes, C. E. and G. D. Wills (1994). Influence of temperature and adjuvants on thidiazuron activity in cotton leaves. *Weed Science* 42(1): 13-17. Mississippi Agric For. Exp. Stn., Delta Branch, Stoneville, MS 38776, USA

Snyder, C. S., G. M. Lorenz, et al. (1994). Late-season foliar fertilization of cotton with potassium nitrate in Jefferson County. *Arkansas Agricultural Experiment Station Research Series*(436): 36-40. {a} State Extension Office, Little Rock, AR, USA

Solanki, S. and S. V. S. Chauhan (1994). Effect of some pesticides on pollen fertility in some crops. *Journal of Phytological Research* 7(1): 83-84. Dep. Bot., R.B.S. Coll., Agra-282 002, India

Souza, J. G. D. and J. V. D. Silva (1994). Phenology and physiology of the perennial cotton after the selection for root starch storage. *Revista Brasileira de Fisiologia Vegetal* 6(2): 145-148. {a} Eng. Agric., M.Sc., Pesquisador do CNPA, CP 174, 58107-720 Campina Grande, PB, Brazil

Steinkraus, D. C. and P. H. Slaymaker (1994). Effect of temperature and humidity on formation, germination, and infectivity of conidia of *Neozygites fresenii* (Zygomycetes: Neozygitaceae) from *Aphis gossypii* (Homoptera: Aphididae). *Journal of Invertebrate Pathology* 64(2): 130-137. Dep. Entomol., Univ. Arkansas, Fayetteville, AR 72701, USA

Stewart, S. D. and M. J. Gaylor (1994). Effects of host switching on oviposition by the tarnished plant bug (Heteroptera: Miridae). *Journal of Entomological Science* 29(2): 231-238. Dep. Entomology, Ala. Agricultural Experiment Station, Auburn Univ., Auburn, AL, USA

Stipanovic, R. D., H. L. Kim, et al. (1994). Raimondalone, a sesquiterpene from a cotton interspecific hybrid. *Phytochemistry Oxford* 36(4): 953-956. {a} USDA, ARS, Southern Crops Res. Lab., College Station, TX 77845, USA

Taher, D. (1994). Expert consultation on cotton pest problems and their control in the Near East (Izmir, Turkey, September 5-7, 1994). *FAO Food and Agriculture Organization of the United Nations Plant Protection Bulletin* 42(3): 139-148. Regional Plant Prot. Off., FAO Regional Off. Near East, PO Box 2223 Dokkii, Cairo, Egypt

- Taware, S. P. and V. P. Patil (1994). Genetic analysis of pink bollworm resistance and other quantitative characters in cotton. *Indian Journal of Genetics and Plant Breeding* 54(2): 137-141. Dep. Genetics Plant Breeding, Agharkar Res. Inst., Gopal Ganesh Agarkar Rd., Pune 411004, India
- Thullen, R. J. and P. E. Keeley (1994). Control of Ivyleaf Morningglory (*Ipomoea hederacea*) in Cotton (*Gossypium hirsutum*) by Combinations of Methods. *Weed Technology* 8(4): 772-776. Agric. Res. Serv., Cotton Res. Stn., 17053 Shafter Ave., Shafter, CA 93262, USA
- Tomar, R. S. S., R. K. Sharma, et al. (1994). Performance of American cotton (*Gossypium hirsutum*) in relation to planting pattern and intercropping with legumes. *Indian Journal of Agronomy* 39(3): 397-402. Coll. Agric., Jawaharlal Nehru Krishi Vishwa Vidyalaya, Indore, Madhya Pradesh 452 001, India
- Tomar, S. K. and O. S. Rana (1994). Incidence of jassid in relation to variety and time of sowing in cotton (*Gossypium* species). *Indian Journal of Agricultural Sciences* 64(1): 70-71. Cotton Res. Stn., Govind Ballabh Pant Univ. Agric. Technol., Bulandshahr, Uttar Pradesh 203 001, India
- Torbert, H. A. and D. W. Reeves (1994). Fertilizer nitrogen requirements for cotton production as affected by tillage and traffic. *Soil Science Society of America Journal* 58(5): 1416-1423. {a} USDA-ARS Grassland Soil Water Res. Lab., 808 East Blackland Rd., Temple, TX 76502, USA
- Trelease, R. N., S. M. Choe, et al. (1994). Conservative amino acid substitutions of the c-terminal tripeptide (Ala-Arg-Met) on cottonseed isocitrate lyase preserve import in vivo into mammalian cell peroxisomes. *European Journal of Cell Biology* 65(2): 269-279. {a} Dep. Botany, Arizona State Univ., Tempe, AZ 85287-1601, USA
- Turley, R. B., D. L. Ferguson, et al. (1994). Isolation and characterization of a cDNA encoding ribosomal protein S16 from cotton (*Gossypium hirsutum* L.). *Plant Physiology Rockville* 106(3): 1219-1220. {a} U.S. Dep. Agric., Agric. Res. Service, Cotton Physiol. Genetics Res., P.O. Box 345, Stoneville, MS 38776, USA
- Turley, R. B., D. L. Ferguson, et al. (1994). Isolation and characterization of a cDNA encoding ribosomal protein L41 from cotton (*Gossypium hirsutum* L.). *Plant Physiology Rockville* 105(4): 1449-1450. {a} United States Dep. Agriculture, Agricultural Res. Serv., Cotton Physiology Genetics Res., P.O. Box 345, Stoneville, MS 38776, USA
- Tyagi, A. P. (1994). Correlation coefficients and selection indices in upland cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research* 28(3): 189-196. Dep. Biol., Sch. Pure Applied Sciences, University South Pacific, PO Box 1168, Suva, Fiji, India
- Tyagi, A. P. (1994). Association of yield components and fibre traits and their functions in upland cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research* 28(3): 159-165. Dep. Biol., Sch. Pure Applied Sciences, Univ. South Pacific, PO Box 1168, Suva, Fiji, India
- Uma, M. S., S. G. Patil, et al. (1994). Performance of field crops under different levels of natural salinity of Tungabhadra project command area. *Indian Journal of Agricultural Sciences* 64(8): 555-557. Agric. Res. Stn., Univ. Agric. Sci., Gangavati, Karnataka 583 227, India
- Uthamasamy, S. and T. Manoharan (1994). Occurrence of leaf perforator (*Bucculatrix loxoptila*) on upland (*Gossypium hirsutum*) and egyptian (*G. barbadense*) cotton. *Indian Journal of Agricultural Sciences* 64(3): 202-204. Dep. Agric. Entomol., Cent. Plant Prot. Studies, Tamil Nadu Agric. Univ., Coimbatore 641 003, India
- Vaissiere, B. E. and S. B. Vinson (1994). Pollen morphology and its effect on pollen collection by honey bees, *Apis mellifera* L. (Hymenoptera: Apidae), with special reference to upland cotton, *Gossypium hirsutum* L. (Malvaceae). *Grana* 33(3): 128-138. {a} Station de Zoologie Apidologie INRA, BP 91, F84143 Montfavet Cedex, France
- Venkatakrishnan, A. S. (1994). Effect of agrochemicals on flower production, and bud and boll shedding of upland cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 39(3): 504-505. Dep. Agron., Tamil Nadu Agric. Univ., Coimbatore 641 003, India

- Wafler, U. and H. Meier (1994). Enzyme activities in developing cotton fibres. *Plant Physiology and Biochemistry Montrouge* 32(5): 697-702. {a} Institut de Biologie Vegetale, Universite de Fribourg, 3 rue Albert Gockel, 1700 Fribourg, Switzerland
- Wall, G. W., J. S. Amthor, et al. (1994). COTCO2: A cotton growth simulation model for global change. *Agricultural and Forest Meteorology* 70(1-4): 289-342. {a} US Water Conservation Lab., 4331 E. Broadway Rd., Phoenix, AZ 85040, USA
- Wan, C. Y. and T. A. Wilkins (1994). A modified hot borate method significantly enhances the yield of high-quality RNA from cotton (*Gossypium hirsutum* L.). *Analytical Biochemistry* 223(1): 7-12. Dep. Agron. Range Sci., Univ. Calif., Davis, CA 95616-8515, USA
- Wang, C. Y. and T. A. Wilkins (1994). Isolation of multiple cDNAs encoding the vacuolar H⁺-ATPase subunit B from developing cotton (*Gossypium hirsutum* L.) ovules. *Plant Physiology Rockville* 106(1): 393-394. {a} Dep. Agronomy Range Sci., Univ. Calif., Davis, CA 95616-8515, USA
- Wang, Y. (1994). Light and electron microscopy study on the endoplasmic reticulum in the microspore of *Gossypium hirsutum*. *Chinese Journal of Botany* 6(1): 23-28. Inst. Vegetables Flowers, Chinese Acad. Agric. Sci., Beijing 100081, China
- Wang, Y., C. H. Lou, et al. (1994). Cytoplasmic connections between tapetal cells of *Gossypium hirsutum*. *Acta Phytophysiological Sinica* 20(3): 308-312. Sch. Biol., Beijing Agric. Univ., Beijing 100094, China
- Wang, Y., S. J. Yang, et al. (1994). Nuclear invagination and nuclear vacuole formation in several plants. *Acta Botanica Sinica* 36(12): 963-966. {a} Inst. Flower Vegetable, Chinese Acad. Agric. Sci., Beijing 100081, China
- Wang, Z. Z., K. Q. Li, et al. (1994). Variation and high frequency somatic embryogenesis of embryogenic callus in *Gossypium hirsutum*. *Acta Botanica Sinica* 36(5): 331-338. Dep. Biol., Shanxi Normal Univ., Xi'an 710062, China
- Weathersbee, A. A., III, D. D. Hardee, et al. (1994). Effects of cotton genotype on seasonal abundance of cotton aphid (Homoptera: Aphididae). *Journal of Agricultural Entomology* 11(1): 29-37. Southern Insect Management Laboratory, U. S. Department Agriculture, Agricultural Research Service, Stoneville, MS 38776, USA
- Wendel, J. F., R. Powley, et al. (1994). Genetic diversity in and phylogenetic relationships of the Brazilian endemic cotton, *Gossypium mustelinum* (Malvaceae). *Plant Systematics and Evolution* 192(1-2): 49-59. Dep. Botany, Iowa State Univ., Ames, IA 50011, USA
- Westerman, R. B. and D. S. Murray (1994). Silverleaf Nightshade (*Solanum elaeagnifolium*) Control in Cotton (*Gossypium hirsutum*) with Glyphosate. *Weed Technology* 8(4): 720-727. Dep. Agron., Oklahoma State Univ., Stillwater, OK 74078, USA
- Wiegand, C. L., J. D. Rhoades, et al. (1994). Photographic and videographic observations for determining and mapping the response of cotton to soil salinity. *Remote Sensing of Environment* 49(3): 212-223. {a} USDA/ARS, Remote Sensing Res. Unit, 2413 E. Hwy. 83, Weslaco, TX 78596-8344, USA
- Wiese, A. F., W. L. Harman, et al. (1994). Economic evaluation of conservation tillage systems for dryland and irrigated cotton (*Gossypium hirsutum*) in the southern Great Plains. *Weed Science* 42(2): 316-321. {a} Texas A and M Univ. Res. and Ext. Cent., 6500 Amarillo Blvd. West, Amarillo, TX 79106, USA
- Wilkins, T. A., C. Y. Wan, et al. (1994). Ancient origin of the vacuolar H⁺-ATPase 69-kilodalton catalytic subunit superfamily. *Theoretical and Applied Genetics* 89(4): 514-524. {a} Dep. Agron. Range Sci., Univ. Calif., Davis, CA 95616-8515, USA

- Willis, G. H., L. L. McDowell, et al. (1994). Permethrin and sulprofos washoff from cotton plants as a function of time between application and initial rainfall. *Journal of Environmental Quality* 23(1): 96-100. {a} USDA-ARS, Soil and Water Res. Unit, 4115 Gourrier Ave., Baton Rouge, LA 70808, USA
- Wilson, F. D., H. M. Flint, et al. (1994). Yield, yield components, and fiber properties of insect-resistant cotton lines containing a *Bacillus thuringiensis* toxin gene. *Crop Science* 34(1): 38-41. {a} USDA-ARS, Western Cotton Res. Lab., 4135 E. Broadway, Phoenix, AZ 85040, USA
- Wilson, L. J. (1994). Resistance of Okra-Leaf Cotton Genotypes to Twospotted Spider Mites (Acari: Tetranychidae). *Journal of Economic Entomology* 87(6): 1726-1735. Commonwealth Sci. Ind. Res. Organ., Div. Plant Ind., Cotton Res. Unit, PO Box 59, Narrabri, NSW 2390, Australia
- Womac, A. R., J. E. Mulrooney, et al. (1994). Influence of oil droplet size on the toxicity of bifenthrin on cotton to tobacco budworm (*Heliothis virescens*). *Pesticide Science* 40(1): 77-83. {a} Agric. Eng. Dep., Univ. Tenn., PO Box 1071, Knoxville, TN 37901, USA
- Wood, C. W., H. A. Torbet, et al. (1994). Free-air CO₂ enrichment effects on soil carbon and nitrogen. *Agricultural and Forest Meteorology* 70(1-4): 103-116. {a} Dep. Agronomy Soils, 202 Funchess Hall, Auburn Univ., Auburn, AL 36849-5412, USA
- Yang, Y. H., J. C. Zhou, et al. (1994). Interactions between *Agrobacterium tumefaciens* and cotton (*Gossypium hirsutum*) cells. *Chinese Journal of Botany* 6(1): 64-68. {a} Dep. Agron., Huazhong Agric. Univ., Wuhan 430070, China
- York, A. C., D. L. Jordan, et al. (1994). Peanut Control in Rotational Crops. *Peanut Science* 21(1): 40-43. {a} Dep. Crop Sci., Box 7620, North Carolina State Univ., Raleigh, NC 27695-7620, USA
- Zhang, J. S., C. P. He, et al. (1994). Structure of micropyle in *Gossypium hirsutum* and the pathway of pollen tube growth. *Acta Botanica Sinica* 36(9): 727-729. Lab. Plant Reproductive Biology, Coll. Life Sciences, Wuhan Univ., Wuhan 430072, China
- Zhong, H. and A. Lauchli (1994). Spatial distribution of solutes, K, Na, Ca and their deposition rates in the growth zone of primary cotton roots: Effects of NaCl and CaCl₂. *Planta Heidelberg* 194(1): 34-41. {a} Dep. Integrative Biol., Univ. Calif., Berkeley, CA 94720, USA
- Zhu, J. and B. S. Weir (1994). Analysis of cytoplasmic and maternal effects I. A genetic model for diploid plant seeds and animals. *Theoretical and Applied Genetics* 89(2-3): 153-159. {a} Program Stat. Genet., Dep. Stat., N.C. State Univ., Raleigh, NC 27695-8203, USA
- Zodape, S. T. and V. D. Chauhan (1994). Stimulation of lipase in cotton seed by pulsed concentrated solar radiation. *Indian Journal of Plant Physiology* 37(4): 289-290. Cent. Salt Marine Chem. Res. Inst., Bhavnagar 364 002, India
- Aamlid, T. S. (1993). Autumn treatment in smooth meadowgrass (*Poa pratensis* L.) grown for seed. *Norsk Landbruksforskning* 7(2): 117-138. Landvik Res. Stn., N-4899 Grimstad, Norway
- Abzalov, M. F. and G. N. Fatkhullaeva (1993). Studies of the anthocyanin pigmentation genetics and its relationship with the leaves shape in cotton *Gossypium hirsutum* L. *Genetika* 29(8): 1356-1365. {a} Sci. Prod. Assoc. "Biolog", Acad. Sci. Uzb., Tashkent, Uzbekistan
- Ansari, B. A., M. M. Rajper, et al. (1993). Hybrid vigour in intervarietal crosses of upland cotton. *Pakistan Journal of Botany* 25(1): 7-12. Dep. Plant Breed. Genet., Sindh Agric. Univ., Tandojam, Pakistan
- Appel, A. G., E. P. Benson, et al. (1993). Laboratory and field evaluations of an entomogenous nematode (Nematoda: Steinernematidae) for German cockroach (Dictyoptera: Blattellidae) control. *Journal of Economic Entomology* 86(3): 777-784. {a} Dep. Entomol., Ala. Agric. Exp. Stn., Auburn University, AL 36849-5413, USA

- Arshad, M., M. Hanif, et al. (1993). Correlation studies on some commercial cotton varieties of *Gossypium hirsutum* L. *Sarhad Journal of Agriculture* 9(1): 49-53. Central Cotton Res. Inst., Multan
- Baker, R. S. (1993). Response of cotton (*Gossypium hirsutum*) to preplant-applied hormone-type herbicides. *Weed Technology* 7(1): 150-153. Plant Physiol., Delta Res. Ext. Cent., P.O. Box 197, Stoneville, MS 38776, USA
- Baloch, M. J., A. R. Lakho, et al. (1993). Heterosis in interspecific cotton hybrids. *Pakistan Journal of Botany* 25(1): 13-20. Cotton Res. Inst., Sakrand, Sindh, Pakistan
- Basra, A. S., G. R. Dhillon, et al. (1993). Influence of ABA and calcium on dark fixation of carbon dioxide in cotton fibres. *Phyton Horn* 33(1): 1-6. {a} Dep. Botany, Punjab Agric. Univ., Ludhiana-141004, India
- Baudracco, S., P. J. Grima, et al. (1993). Quantitative cytochemical localization of cinnamyl alcohol dehydrogenase activity in plant tissues. *Phytochemical Analysis* 4(5): 205-209. {a} Plant Cell Mol. Sci. Group, Div. Life Sci., King's College London, Campden Hill Rd., London W8 7AH, UK
- Bauer, P. J., J. J. Camberato, et al. (1993). Cotton yield and fiber quality response to green manures and nitrogen. *Agronomy Journal* 85(5): 1019-1023. {a} USDA-ARS Coastal Plains Soil, Water and Plant Res. Cent., P.O. Box 3039, Florence, SC 29502, USA
- Baumhardt, R. L., J. W. Keeling, et al. (1993). Tillage and residue effects on infiltration into soils cropped to cotton. *Agronomy Journal* 85(2): 379-383. {a} Texas Agric. Exp. Stn., Route 3, Box 219, Lubbock, TX 79401
- Baumhardt, R. L., C. W. Wendt, et al. (1993). Tillage and furrow diking effects on water balance and yields of sorghum and cotton. *Soil Science Society of America Journal* 57(4): 1077-1083. {a} Texas Agric. Exp. Stn., Route 3, Box 219, Lubbock, TX 79401, USA
- Begum, S., A. Rahman, et al. (1993). Studies on the hemicellulose from Dhabdhabey (S-718) jute. *Dhaka University Studies Part B Science* 41(2): 107-112. Dep. Chemistry, Dhaka Univ., Dhaka-1000, Bangladesh
- Bhaskar, K. S., S. T. Gaikwad, et al. (1993). Response of upland cotton (*Gossypium hirsutum*) to the levels of fertilizers in shallow soils of Sanogi watershed near Nagpur. *Indian Journal of Agronomy* 38(1): 89-92. National Bureau Soil Survey Land-Use Planning, Nagpur, Maharashtra 440 010, India
- Boquet, D. J., E. B. Moser, et al. (1993). Nitrogen effects on boll production of field-grown cotton. *Agronomy Journal* 85(1): 34-39. {a} Louisiana Agric. Exp. Stn., Northeast Res. Stn., P.O. Box 438, St. Joseph, LA 71366
- Bourland, F. M., C. E. Ortiz, et al. (1993). Registration of Miscot 8001, Miscot 8004 and Miscot 8006 germplasm lines of cotton. *Crop Science* 33(5): 1106. {a} Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA
- Bradow, J. M. (1993). Inhibitions of cotton seedling growth by volatile ketones emitted by cover crop residues. *Journal of Chemical Ecology* 19(6): 1085-1108. USDA, ARS, Southern Regional Res. Cent., P.O. Box 19687, New Orleans, LA 70179, USA
- Brendecke, J. W., R. D. Axelson, et al. (1993). Soil microbial activity as an indicator of soil fertility: Long-term effects of municipal sewage sludge on an arid soil. *Soil Biology and Biochemistry* 25(6): 751-758. {a} Dep. Soil. Water Sci., Stat. Support Unit, Coll. Agric., Univ. Ariz., Tucson, AZ 85721, USA
- Brubaker, C. L., J. A. Koontz, et al. (1993). Bidirectional cytoplasmic and nuclear introgression in the New World cottons, *Gossypium barbadense* and *G. hirsutum* (Malvaceae). *American Journal of Botany* 80(10): 1203-1208. {a} Dep. Bot., Iowa State Univ., Ames, IA 50011, USA
- Brubaker, C. L. and J. R. Wendel (1993). On the specific status of *Gossypium lanceolatum* Todaro. *Genetic Resources and Crop Evolution* 40(3): 165-170. Dep. Botany, Iowa State Univ., Ames, IA 50011, USA

- Carmi, A., Z. Plaut, et al. (1993). Cotton root growth as affected by changes in soil water distribution and their impact on plant tolerance to drought. *Irrigation Science* 13(4): 177-182. {a} Inst. Soils and Water, Agric. Res. Organization, Volcani Cent., P.O. Box 6, Bet Dagan 50250, Israel
- Chapman, K. D. and T. S. Moore, Jr. (1993). Catalytic properties of a newly discovered acyltransferase that synthesizes N-acylphosphatidylethanolamine in cottonseed (*Gossypium hirsutum* L.) microsomes. *Plant Physiology Rockville* 102(3): 761-769. {a} Botany Dep., La. State Univ., Baton Rouge, LA 70803, USA
- Close, T. J., R. D. Fenton, et al. (1993). A view of plant dehydrins using antibodies specific to the carboxy terminal peptide. *Plant Molecular Biology* 23(2): 279-286. Dep. Botany Plant Sci., Univ. California, Riverside, CA 92521, USA
- Colyer, P. D. and P. R. Vernon (1993). Effect of tillage on cotton plant populations and seedling diseases. *Journal of Production Agriculture* 6(1): 108-111. {a} Red River Res. Station, P.O. Box 8550, La. State Univ. Agric. Center, Bossier City, LA 71113-8550
- Conkerton, E. J., D. C. Chapital, et al. (1993). Fruiting sites in cotton: Seed quality. *Journal of Agricultural and Food Chemistry* 41(6): 882-885. {a} Southern Regional Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., P.O. Box 19687, New Orleans, LA 70179-0687, USA
- Cook, C. G. and Z. K. M. El (1993). Fruiting and lint yield of cotton cultivars under irrigated and nonirrigated conditions. *Field Crops Research* 33(4): 411-421. {a} USDA-Agricultural Res. Serv., Subtropical Agric. Res. Lab., Weslaco, TX 78596, USA
- Cox, J. S., C. E. Shamu, et al. (1993). Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. *Cell* 73(6): 1197-1206. Dep. Biochem. Biophysics, Univ. Calif. Med. Sch., San Francisco, CA 94143-0448, USA
- Crane, C. F., H. J. Price, et al. (1993). Identification of a homeologous chromosome pair by in situ DNA hybridization to ribosomal RNA loci in meiotic chromosomes of cotton (*Gossypium hirsutum*). *Genome* 36(6): 1015-1022. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843, USA
- Croise, L. and F. Lieutier (1993). Effects of drought on the induced defence reaction of Scots pine to bark beetle-associated fungi. *Annales des Sciences Forestieres Paris* 50(1): 91-97. INRA, Station de Zool. Forestiere, Ardon, 45160 Olivet, France
- Dabo, S. M., E. D. Mitchell, Jr., et al. (1993). A method for the isolation of nuclear DNA from cotton (*Gossypium*) leaves. *Analytical Biochemistry* 210(1): 34-38. {a} Biochem. Molecular Biol. Dep., Oklahoma State Univ., Stillwater, Oklahoma 74078-0454, USA
- Dani, R. G., O. Y. Vesmanovo, et al. (1993). Studies on clonal micropropagation of cotton through meristem culture. *Advances in Plant Sciences* 6(2): 260-264. {a} Div. Crop. Improvement, Cent. Inst. Cotton Res., Post Bag No. 225, GPO Panjiri-Wardha Rd., Nagpur-440-001, India
- Davidonis, G. (1993). Cotton fiber growth and development in vitro: Effects of tunicamycin and monensin. *Plant Science Limerick* 88(2): 229-236. U.S. Dep. Agriculture, ARS, Southern Regional Res. Cent., P.O. Box 19687, New Orleans, LA 70119, USA
- Davis, C. J. G. and D. C. Sheppard (1993). Redistribution of metals and nutrients by fire ants in a flue dust contaminated pasture. *Soil Biology and Biochemistry* 25(6): 747-750. {a} Agron. Dep., Coastal Plain Exp. Station, Univ. Georgia, Tifton, GA 31793-0748, USA
- Davis, D., J. Merida, et al. (1993). Independent elicitation of the oxidative burst and phytoalexin formation in cultured plant cells. *Phytochemistry Oxford* 32(3): 607-611. {a} Dep. Med. Chemistry Pharmacognosy, Purdue Univ., West Lafayette, IN 47907-1333, USA

Delwiche, C. F. and T. D. Sharkey (1993). Rapid appearance of carbon-13 in biogenic isoprene when carbon-13 carbon dioxide is fed to intact leaves. *Plant Cell and Environment* 16(5): 587-591. {a} Dep. Bot., Univ. Wis., Madison, WI 53706-1381, USA

Dhingra, H. R., C. L. Goswami, et al. (1993). Studies on the reproductive functions of cotton. *Indian Journal of Plant Physiology* 36(2): 118-120. {a} Dep. Botany, Haryana Agric. Univ., Hisar-125 004, India

Donald, W. W. (1993). Models and sampling for studying weed seed survival with wild mustard (*Sinapis arvensis*) as a case study. *Canadian Journal of Plant Science* 73(2): 637-645. Cropping Systems and Water Quality Res. Unit, Agric. Res. Serv., U.S. Dep. Agric., 244 Agric. Eng. Build., Univ. Mo., Columbia, MO 65211, USA

Duarsa, M. A. P., M. J. Hill, et al. (1993). Soil moisture and temperature affect condensed tannin concentrations and growth in *Lotus corniculatus* and *Lotus pedunculatus*. *Australian Journal of Agricultural Research* 44(7): 1667-1681. {a} Div. Anim. Prod., CSIRO, Pastoral Res. Lab., Private Mail Bag, Post Office, Armidale, N.S.W. 2350

Dudienas, C. and H. Kimati (1993). Auxanographic characterization of *Colletotrichum gossypii* and *Colletotrichum gossypii* var. *cephalosporioides*. *Summa Phytopathologica* 19(3-4): 152-156. {a} Secao Fitopatologia, Inst. Agronomico, Caixa Postal 28, 13.020 Campinas, SP, Brazil

Ekukole, G. (1993). Host plants of *Aphis gossypii* Glover (Homoptera: Aphididae) in the Maroua are North Cameroon. *Journal of African Zoology* 107(2): 169-174. Inst. Agric. Res., P.O. Box 33 Maroua, Cameroon

Ellis, J. R., R. D. Lacewell, et al. (1993). Preferred irrigation strategies in light of declining government support. *Journal of Production Agriculture* 6(1): 112-118. {a} Dep. Agric. Economic, Washington State Univ., Pullman, WA 99164-6210

Ephrath, J. E., A. Marani, et al. (1993). Photosynthetic rate, stomatal resistance and leaf water potential in cotton (*Gossypium hirsutum* L.) as affected by soil moisture and irradiance. *Photosynthetica Prague* 29(1): 63-71. {a} Centre Desert Agrobiologie, J. Blaustein Inst. Desert Res., Ben Gurion Univ. Negev, Sede Boqer 84993, Israel

Ferguson, W. and J. Yee (1993). Evaluation of professional scouting programs in cotton production. *Journal of Production Agriculture* 6(1): 100-103. {a} USDA-Economic Res. Serv., 1301 New York Ave. NW, Washington, DC 20005-4788

Firoozabady, E. and D. L. Deboer (1993). Plant regeneration via somatic embryogenesis in many cultivars of cotton (*Gossypium hirsutum* L.). *In Vitro Cellular and Developmental Biology Plant* 29P(4): 166-173. {a} DNA Plant Technol. Corporation, 6701 San Pablo Ave., Oakland, CA 94608, USA

Franklyn, K. M. and J. R. Warmington (1993). Cloning and nucleotide sequence analysis of the *Candida albicans* enolase gene. *FEMS Federation of European Microbiological Societies Microbiology Letters* 111(1): 101-107. {a} Sch. Biomedical Sci., Curtin Univ. Technol., GPO Box U1987, Perth, Western Australia 6001, Australia

Frenay, J. R., D. L. Chen, et al. (1993). Use of nitrification inhibitors to increase fertilizer nitrogen recovery and lint yield in irrigated cotton. *Fertilizer Research* 34(1): 37-44. {a} Div. Plant Industry, CSIRO, G.P.O. Bo 1600, Canberra, A.C.T. 2601, Australia

Furbeck, S. M., F. M. Bourland, et al. (1993). Relationships of seed and germination measurements with resistance to seed weathering in cotton. *Seed Science and Technology* 21(3): 505-512. Dep. Agron., Miss. State Univ., P.O. Box 5248, Mississippi State, MS 39762, USA

Furbeck, S. M., F. M. Bourland, et al. (1993). Inheritance of resistance to seed deterioration in cotton. *Euphytica* 69(3): 203-209. {a} Dep. Crop Sci., North Carolina State Univ., Raleigh, NC 27607, USA

Gomez, M., J. S. Johnston, et al. (1993). Nuclear 2C DNA content of *Gossypium hirsutum* L. accessions determined by flow cytometry. *Biologisches Zentralblatt* 112(4): 351-357. {a} Dep. Soil and Crop Sci., Tex. A and M Univ., College Station, TX 77843, USA

Goswami, C. L. (1993). Effect of nitrogen on diurnal variations in adaxial and abaxial stomatal response of cotton (*Gossypium hirsutum* L. CV. H-777). *Indian Journal of Plant Physiology* 36(2): 137-138. Dep. Botany, Haryana Agric. Univ., Hisar-125 004, India

Grontoft, M. (1993). A rapid screening method for testing the resistance of cotyledons to downy mildew in *Brassica napus* and *Brassica campestris*. *Plant Breeding* 110(3): 207-211. Plant Protection Centre, Box 44, S-230 53 Alnarp, Sweden

Guimaraez, R. M., A. C. Fraga, et al. (1993). Effects of the method of osmotic conditions over germination and cotton seed (*Gossypium hirsutum* L.) vigour in ideal conditions and hydric and thermic stress. *Ciencia e Pratica* 17(3): 215-223. Engenheiro Agronomo, M.Sc., Faepe, Lavras, MG, Brazil

Guinn, G. and D. L. Brummett (1993). Leaf age, decline in photosynthesis, and changes in abscisic acid, indole-3-acetic acid, and cytokinin in cotton leaves. *Field Crops Research* 32(3-4): 269-275. US Dep. Agric., Agric. Res. Serv., Western Cotton Res. Lab., 4135 E. Broadway, Phoenix, Ariz. 85040, USA

Hamilton, J. G. and M. P. Zalucki (1993). Interactions between a specialist herbivore, *Crociosema plebejana*, and its host plants *Malva parviflora* and cotton, *Gossypium hirsutum*: Oviposition preference. *Entomologia Experimentalis et Applicata* 66(3): 207-212. {a} Australian Plague Locust Commission, DPIE, GPO Box 858, Canberra ACT 2601, Australia

Hamilton, J. G. and M. P. Zalucki (1993). Interactions between a specialist herbivore, *Crociosema plebejana*, and its host plants *Malva parviflora* and cotton, *Gossypium hirsutum*: Larval performance. *Entomologia Experimentalis et Applicata* 66(3): 199-205. {a} Australian Plague Locust Commission, DPIE, GPO Box 858, Canberra ACT 2601, Australia

Hassan, S., R. D. Ahmad, et al. (1993). Genetic analysis of yield and yield components in various crosses of American upland cotton (*Gossypium hirsutum* L.). *Sarhad Journal of Agriculture* 9(4): 327-335. {a} Fac. Agric., Gomal Univ., D. I. Khan, Pakistan

Heitholt, J. J. (1993). Cotton boll retention and its relationship to lint yield. *Crop Science* 33(3): 486-490. Cotton Physiol. and Genetics, USDA-ARS, P.O. Box 345, Stoneville, MS 38776, USA

Heitholt, J. J., W. T. Pettigrew, et al. (1993). Growth, boll opening rate, and fiber properties of narrow-row cotton. *Agronomy Journal* 85(3): 590-594. {a} USDA-ARS, Cotton Physiol. Genetics, P.O. Box 345, Stoneville, MS 38776, USA

Hendrix, D. L. (1993). Rapid extraction and analysis of nonstructural carbohydrates in plant tissues. *Crop Science* 33(6): 1306-1311. West. Cotton Res. Lab., USDA-ARS, 4135 E. Broadway Rd., Phoenix, AZ 85040, USA

Hesler, L. S. and G. R. Sutter (1993). Effect of trap color, volatile attractants, and type to toxic bait dispenser or captures of adult corn rootworm beetles (Coleoptera: Chrysomelidae). *Environmental Entomology* 22(4): 743-750. Northern Grain Insects Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., Brookings, SD 57006, USA

Heussen, S. C. and E. B. Dowdle (1993). Erythrina protease inhibitor: Kinetics of interaction with tissue plasminogen activator. *South African Journal of Science* 89(3): 131-135. {a} Dep. Clin. Sci. Immunol., Univ. Cape Town Med. Sch., Observatory, 7925 South, Central African Republic

Hong, S. S. and W. Y. Soh (1993). Early ontogeny of vascular cambium in cotyledonary node of seedlings in *Pinus koraiensis* S. et Z. *Korean Journal of Botany* 36(1): 29-34. {a} Dep. Biol. Educ., Chonnam Natl. Univ., Kwangju, North Korea

- Hu, S. Y., Y. Wang, et al. (1993). The regional aggregation of plastids and mitochondria in microsporocyte and microspore of *Gossypium hirsutum*. *Chinese Journal of Botany* 5(1): 11-17. Biol. Dep., Peking Univ., Beijing 100871, China
- Hu, S. Y., Z. f. Yuan, et al. (1993). The giant stacked rough endoplasmic reticulum and its activity during cotton microspore development. *Acta Botanica Sinica* 35(8): 573-579. Dept. Biology, Peking University, Beijing 100871, China
- Huberman, M., E. Pressman, et al. (1993). Pith autolysis in plants: IV. The activity of polygalacturonase and cellulase during drought stress induced pith autolysis. *Plant and Cell Physiology* 34(6): 795-801. {a} Biol. Dep., Wake Forest Univ., Winston-Salem, NC 27109, USA
- Ibrahim, G. and H. I. Nirenberg (1993). Response of some Sudanese cotton cultivars to race 1 and 5 of *Fusarium oxysporum* f. sp. *vasinfectum*. *Zeitschrift fuer Pflanzenkrankheiten und Pflanzenschutz* 100(6): 645-651. {a} Botany and Plant Pathol. Sect., Agric. Res. Corp., Wad Medani, Sudan
- Islam, Z., M. S. Islam, et al. (1993). Optimum fertilization for upland cotton (*Gossypium hirsutum*) in red-brown terrace soil of Bangladesh. *Indian Journal of Agricultural Sciences* 63(4): 200-203. Div. Soil Sci., Bangladesh Agric. Res. Isnt., Gazipur, Bangladesh 1701
- Jacks, T. J., T. P. Hensarling, et al. (1993). Permanent gases inside healthy and microbially infected cotton fruit during development. *Biochemical and Biophysical Research Communications* 191(3): 1284-1287. {a} Southern Regional Res. Center, ARS, USDA, PO Box 19687, New Orleans, LA 70179, USA
- Jacks, T. J. and O. Hinojosa (1993). Superoxide radicals in intact tissues and in dimethyl sulphoxide-based extracts. *Phytochemistry Oxford* 33(3): 563-568. Southern Regional Res. Cent., U.S. Dep. Agric., P.O. Box 19867, New Orleans, LA 70179, USA
- Jeffers, D. P. and P. A. Roberts (1993). Effect of planting date and host genotype on the root-knot nematode-fusarium wilt disease complex of cotton. *Phytopathology* 83(6): 645-654. {a} Dep. Nematology, Univ. Calif., Riverside, CA 92521, USA
- Jenkins, J. N., W. L. Parrott, et al. (1993). Growth and survival of *Heliothis virescens* (Lepidoptera: Noctuidae) on transgenic cotton containing a truncated form of the delta endotoxin gene from *Bacillus thuringiensis*. *Journal of Economic Entomology* 86(1): 181-185. {a} Crop Sci., Res. Lab., USDA-ARS, P.O. Box 5367, Mississippi State, MS 39762
- Johnson, D. H., D. L. Jordan, et al. (1993). Nicosulfuron, primisulfuron, imazethapyr, and DPX-PE350 injury to succeeding crops. *Weed Technology* 7(3): 641-644. {a} American Cyanamid Co., P.O. Box 400, Princeton, NJ 08543-0400, USA
- Jones, R. W., J. R. Cate, et al. (1993). Pollen feeding and survival of the boll weevil (Coleoptera: Curculionidae) on selected plant species in Northeastern Mexico. *Environmental Entomology* 22(1): 99-108. {a} Dep. Entomol., Texas A and M University, College Station, TX 77843, USA
- Jordan, D. L., R. E. Frans, et al. (1993). Total postemergence herbicide programs in cotton (*Gossypium hirsutum*) with sethoxydim and DPX-PE350. *Weed Technology* 7(1): 196-201. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72701, USA
- Jordan, D. L., R. E. Frans, et al. (1993). Cotton (*Gossypium hirsutum*) response to DPX-PE350 applied postemergence. *Weed Technology* 7(1): 159-162. Dep. Agron., Univ. Ark., Fayetteville, AR 72701, USA
- Jordan, D. L., R. E. Frans, et al. (1993). DPX-PE350 does not interact with early-season insecticides in cotton (*Gossypium hirsutum*). *Weed Technology* 7(1): 92-96. Dep. Agron., Univ. Arkansas, Fayetteville, AR 72703, USA

- Jordan, D. L., D. H. Johnson, et al. (1993). Carryover of DPX-PE350 to grain sorghum (*Sorghum bicolor*) and soybean (*Glycine max*) on two Arkansas soils. *Weed Technology* 7(3): 645-649. {a} Northeast. Res. Stn., La. State Univ., P.O. Box 438, St. Joseph, LA 71366, USA
- Jordan, D. L., M. R. McClelland, et al. (1993). Effect of MSMA on entireleaf morningglory (*Ipomoea hederacea* var. *integriuscula*) control with postemergence herbicides. *Weed Technology* 7(1): 36-41. {a} Dep. Agron., Univ. Arkansas, Fayetteville, AR 72703, USA
- Jordan, D. L., A. C. York, et al. (1993). Clomazone as a component in cotton (*Gossypium hirsutum*) herbicide programs. *Weed Technology* 7(1): 202-211. {a} Dep. Crop Sci., N.C. State Univ., Raleigh, NC 27695-7620, USA
- Kabilan, V. and K. Murugesan (1993). Studies on the phytotoxicity of a few pesticides. *Geobios Jodhpur* 20(3): 143-147. CAS Botany, Univ. Madras, Madras-600 025, India
- Kalabokis, V., M. M. Santoro, et al. (1993). Effect of sodium and nucleotide on the stability of solubilized calcium free calcium-ATPase from scallop sarcoplasmic reticulum. *Biochemistry* 32(16): 4389-4396. {a} Dep. Med. Biochemistry, Southern Ill. Univ., Carbondale, IL 62901-4409, USA
- Karimov, K. K. and S. V. Dontsova (1993). Evaluating the viability of cotton seeds on the basis of their electrical conductance. *Izvestiya Akademii Nauk Respubliki Tadjikistan Otdelenie Biologicheskikh Nauk*(3): 24-26. Inst. Plant Physiol. Biophys., Taj. Acad. Sci., Dushanbe, Tajikistan
- Keeling, J. W., C. G. Henniger, et al. (1993). Effects of DPX PE350 on cotton (*Gossypium hirsutum*) growth, yield, and fiber quality. *Weed Technology* 7(4): 930-933. Texas Agric. Exp. Stn., Lubbock, TX 79401, USA
- Kerns, D. L. and M. J. Gaylor (1993). Induction of cotton aphid outbreaks by insecticides in cotton. *Crop Protection* 12(5): 387-393. {a} Sandoz Agro Inc., PO Box 5189, Greenville, MS 38704, USA
- Khoshkhoo, N., P. A. Hedin, et al. (1993). Effects of bioregulators on the terpenoid aldehydes in root-knot nematode infected cotton plants. *Journal of Agricultural and Food Chemistry* 41(12): 2442-2446. {a} Crop Sci. Res. Lab., Agric. Res. Service, Dep. Agric., Mississippi State, MS 39762-5367, USA
- Kimball, B. A. and J. R. Mauney (1993). Response of cotton to varying carbon dioxide, irrigation, and nitrogen: Yield and growth. *Agronomy Journal* 85(3): 706-712. {a} USDA-ARS, U.S. Water Conserv. Western Cotton Res. Lab., Phoenix, AZ 85040, USA
- Kloth, R. H. (1993). New evidence relating the pilose allele and micronaire reading in cotton. *Crop Science* 33(4): 683-687. USDA-ARS, Cotton Physiol. and Genetics Res. Unit, P.O. Box 345, Stoneville, MS 38776, USA
- Koeniger, M. and K. Winter (1993). Growth and photosynthesis of *Gossypium hirsutum* L at high photon flux densities: Effects of soil temperatures and nocturnal air temperatures. *Agronomie Paris* 13(5): 423-431. {a} Smithsonian Tropical Res. Inst., P.O. Box 2072, Balboa, Panama
- Kohel, R. J., C. R. Benedict, et al. (1993). Incorporation of (¹⁴C)glucose into crystalline cellulose in aberrant fibers of a cotton mutant. *Crop Science* 33(5): 1036-1040. {a} Crop Germplasm Res. Unit, Southern Crops Res. Lab., USDA-ARS, Route 5 Box 805, College Station, TX 77845, USA
- Koniger, M. and K. Winter (1993). Reduction of photosynthesis in sun leaves of *Gossypium hirsutum* L. under conditions of high light intensities and suboptimal leaf temperatures. *Agronomie Paris* 13(7): 659-668. Smithsonian Tropical Res. Inst., PO Box 2072, Balboa, Panama
- Koraddi, V. R., S. B. Modak, et al. (1993). Studies on efficient utilization of rain water and soil moisture in rainfed cotton. *Journal of Maharashtra Agricultural Universities* 18(1): 27-29. Agric. Res. Station, Dharwad-580 007, India

- Kuznetsov, V. V., V. Y. Rakitin, et al. (1993). Why does heat shock increase salt resistance in cotton plants? *Plant Physiology and Biochemistry Montrouge* 31(2): 181-188. {a} Lab. Molecular Mechanisms of Adaptation, Inst. Plant Physiology, Academy Sci. Russia, Botanicheskaya 35, 127276 Moscow, Russia
- Landolt, P. (1993). Suitability of six species of Commelinaceae as larval hosts of *Mouralia tinctorides* (Lepidoptera: Noctuidae, Plusiinae). *Florida Entomologist* 76(4): 572-576. United States Dep. Agric., Agric. Res. Serv., P.O. Box 1565, Gainesville, FL 32604, USA
- Landolt, P. J. (1993). Effects of host plant leaf damage on cabbage looper moth attraction and oviposition. *Entomologia Experimentalis et Applicata* 67(1): 79-85. U.S. Dep. Agric., Agric. Res. Serv., P.O. Box 14565, Gainesville, FL 32604, USA
- Lege, K. E., C. W. Smith, et al. (1993). Planting date and plant density effects on condensed tannin concentration of cotton. *Crop Science* 33(2): 320-324. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, TX 77843-2474, USA
- Leidi, E. O. and J. C. Gutierrez (1993). Note about variability in root growth of cotton at seedling stage: Effect of genotype and osmotic stress. *Investigacion Agraria Produccion y Proteccion Vegetales* 8(2): 165-169. {a} Dpto. Biol. Vegetal, IRNAS, Apdo. 1052, 41080 Sevilla, Spain
- Leidi, E. O., J. M. Lopez, et al. (1993). Searching for tolerance to water stress in cotton genotypes: Photosynthesis, stomatal conductance and transpiration. *Photosynthetica Prague* 28(3): 383-390. {a} Dpto. Biologia Vegetal, I.R.N.A.S., Aptdo. 1052, E-41080 Sevilla, Spain
- Li, L. and R. M. Brown, Jr. (1993). Beta-Glucan synthesis in the cotton fiber: II. Regulation and kinetic properties of beta-glucan synthases. *Plant Physiology Rockville* 101(4): 1143-1148. {a} Dep. Botany, Univ. Texas Austin, Austin, TX 78713-7640, USA
- Li, L., R. R. Drake, Jr., et al. (1993). Beta-Glucan synthesis in the cotton fiber: III. Identification of UDP-glucose-binding subunits of beta-glucan synthases by photoaffinity labeling with (beta-phosphorus-32)-5'-N-3-UDP-glucose. *Plant Physiology Rockville* 101(4): 1149-1156. {a} Dep. Botany, Univ. Texas Austin, Austin, TX 78713-7640, USA
- Liebig, M. A., A. J. Jones, et al. (1993). Controlled wheel traffic effects on soil properties in ridge tillage. *Soil Science Society of America Journal* 57(4): 1061-1066. {a} Dep. Agronomy, USDA-ARS, Univ. Nebraska, Lincoln, NE 68583, USA
- Lopez, J. M., E. O. Leidi, et al. (1993). Photosynthesis, stomatal conductance, water use efficiency and leaf temperature of cotton cultivars under water stress. *Investigacion Agraria Produccion y Proteccion Vegetales* 8(1): 17-27. {a} Dpto. de Algodon, CIDA, Las Torres-Tomejil, Aptdo. Oficial, 41200 Alcala del Rio
- Mace, M. E., R. D. Stipanovic, et al. (1993). Toxicity of cotton phytoalexins to zoopathogenic fungi. *Natural Toxins* 1(5): 294-295. {a} Cotton Pathology Res. Unit, Southern Crops Res. Lab., USDA, ARS, Route 5, Box 805, College Station, TX 77845, USA
- Mahaffee, W. F. and P. A. Backman (1993). Effects of seed factors on spermosphere and rhizosphere colonization of cotton by *Bacillus subtilis* GB03. *Phytopathology* 83(10): 1120-1125. Dep. Plant Pathol., Ala. Agric. Exp. Stn., Auburn Univ., Auburn, AL 36849, USA
- Marienfild, S. and R. Stelzer (1993). X-ray microanalyses in roots of aluminum-treated *Avena sativa* plants. *Journal of Plant Physiology* 141(5): 569-573. Botanisches Inst., Tierarztlichen Hochschule Hannover, Buenteweg 17 d, D-3000 Hannover 71, Germany
- Martsinkovskaya, A. I., V. L. Motin, et al. (1993). Cotton DNA fingerprinting for *Gossypium*: The use of M13 phage DNA as a hybridization probe. *Biopolimery i Kletka* 9(1): 51-53. Inst. Bioorg. Chem., Acad. Sci. Uzb., Tashkent, Uzbekistan

- Mascagni, H. J., Jr., T. C. Keisling, et al. (1993). Response of fast fruiting cotton cultivars to nitrogen on a clayey soil. *Journal of Production Agriculture* 6(1): 104-108. {a} Soil Test. Res. Lab., Hwy 1 S, Marianna, AR 72360
- McCall, P. J., T. C. J. Turlings, et al. (1993). Role of plant volatiles in host location by the specialist parasitoid *Microplitis croceipes* Cresson (Braconidae: Hymenoptera). *Journal of Insect Behavior* 6(5): 625-639. Insect. Attractants, Behavior Basis Biol. Res. Lab., Agric. Res. Serv., U.S. Dep. Agric., P.O. Box 14565, Gainesville, FL 32604, USA
- McConnell, J. S., W. H. Baker, et al. (1993). Nitrogen fertilization of cotton cultivars of differing maturity. *Agronomy Journal* 85(6): 1151-1156. {a} Univ. Arkansas, Southeast Res. Ext. Center, Monticello, AR, USA
- McWhorter, C. G. (1993). A 16-yr survey on levels of Johnsongrass (*Sorghum halepense*) in Arkansas, Louisiana, and Mississippi. *Weed Science* 41(4): 669-677. Plant Physiol., App. Tech. Res. Unit, Agric. Res. Serv., U.S. Dep. Agric., Stoneville, MS 38776, USA
- Mert, H. H. (1993). Investigation on the endogenous ABA levels of seeds of cotton cultivars under salt stress conditions. *Turkish Journal of Botany* 17(4): 201-205. Biol. Dep., Fac. Education, Dokuz Eylul Univ., 35150 Buca, Izmir, Turkey
- Mohod, V. K. and S. B. Parde (1993). Assessment of isolines in some of the varieties of upland cotton for ginning percentage. *Journal of Maharashtra Agricultural Universities* 18(2): 258-260. Punjabrao Krishi Vidyapeeth, Akola-444 104, India
- Monks, C. D. and P. A. Banks (1993). Effect of straw, ash, tillage on dissipation of imazaquin and imazethapyr. *Weed Science* 41(1): 133-137. {a} Div. Plant Soil Sci., P.O. Box 6108, 1090 Agric Sci., West Va. Univ., Morgantown, WV 26506-6108, USA
- Moody, D. S., D. G. Bottrell, et al. (1993). Late season migration of the boll weevil in the Texas rolling plains. *Southwestern Entomologist* 18(1): 1-10. {a} Route No. 1, Box 409, Kingsville, TX, USA
- Morris, D. A. (1993). The role of auxin in the apical regulation of leaf abscission in cotton (*Gossypium hirsutum* L.). *Journal of Experimental Botany* 44(261): 807-814. Dep. Biol., Sch. Biol. Sci., Univ. Southampton, Bassett Crescent East, Southampton SO9 3TU, UK
- Mullins, G. L. (1993). Cotton root growth as affected by phosphorus fertilizer placement. *Fertilizer Research* 34(1): 23-26. Auburn Univ., Dep. Agron. Soils, Auburn, AL 36849-5412, USA
- Mullins, G. L. and C. H. Burmester (1993). Uptake of sulfur by four cotton cultivars grown under field conditions. *Journal of Plant Nutrition* 16(6): 1071-1081. {a} Auburn Univ., Dep. Agron. and Soils, Auburn, AL 36849-5412, USA
- Mullins, G. L. and C. H. Burmester (1993). Accumulation of copper, iron, manganese and zinc by four cotton cultivars. *Field Crops Research* 32(1-2): 129-140. {a} Auburn Univ., Dep. Agronomy Soils, 202 Funches Hall, Auburn, AL 36849-5412, USA
- Nandeshwar, S. B. and A. A. Saoji (1993). Evaluation of hybrid progenies from stored pollen in cotton. *Advances in Plant Sciences* 6(2): 213-218. {a} Crop Improvement Div., Central Inst. Cotton Res., Post Bag No. 225, Nagpur-440 001, M.S., India
- Naranjo, S. E. and J. M. Martin (1993). Comparative development, reproduction, and oviposition of pink bollworm (Lepidoptera: Gelechiidae) on a resistant okra-leaf cotton and commercial upland an pima cultivars. *Journal of Economic Entomology* 86(4): 1094-1103. Western Cotton Res. Lab., USDA-ARS, 4135 E. Broadway Road, Phoenix, AZ 85040, USA
- Navasero, R. C. and S. B. Ramaswamy (1993). Influence of plant age, water stress, larval damage, and presence of conspecific eggs on oviposition by *Heliothis virescens* (F.) on cotton. *Journal of Applied Entomology* 115(1): 97-106. {a} Dep. Entomology, Mississippi State Univ., Mississippi State, MS 39762

Ockerby, S. E., D. J. Lyons, et al. (1993). Irrigation frequency and nitrogen fertilizers modify cotton yield at Emerald, Central Queensland. *Australian Journal of Agricultural Research* 44(6): 1389-1402. {a} Queensland Dep. Primary Industries, P.O. Box 591, Ayr, Qld. 4807, Australia

Ogle, H. J., A. M. Stirling, et al. (1993). Pathogenicity of fungi associated with seedling disease of cotton. *Australian Journal of Experimental Agriculture* 33(7): 923-929. Dep. Agric., Univ. Queensland, Brisbane, Qld 4072, Australia

Ojala, A. (1993). Effects of temperature and irradiance on the growth of two freshwater photosynthetic cryptophytes. *Journal of Phycology* 29(3): 278-284. Inst. Freshwater Ecol., Ferry House, Ambleside, Cumbria LA22 0LP, UK

Okamura, H., A. Mimura, et al. (1993). Antioxidant activity of tannins and flavonoids in *Eucalyptus rostrata*. *Phytochemistry Oxford* 33(3): 557-561. Biotechnol. Res. Lab., Kobe Steel Ltd., 25-14, Kanondai 1-chome, Tsukuba, Ibaraki 304, Japan

Okuda, K., L. Li, et al. (1993). Beta-Glucan synthesis in the cotton fiber: I. Identification of beta-1,4- and beta-1,3-glucans synthesized in vitro. *Plant Physiology Rockville* 101(4): 1131-1142. {a} Dep. Botany, Univ. Texas Austin, Austin, TX 78713-7640, USA

Olszyk, D., A. Bytnerowicz, et al. (1993). Cotton yield losses and ambient ozone concentrations in California's San Joaquin Valley. *Journal of Environmental Quality* 22(3): 602-611. {a} USEPA/Environ. Res. Lab., Corvallis, OR 97333, USA

Oosterhaven, K., K. J. Hartmans, et al. (1993). Inhibition of potato (*Solanum tuberosum*) sprout growth by the monoterpene S-carvone: Reduction of 3-hydroxy-3-methylglutaryl-coenzyme A reductase activity without effect on its mRNA level. *Journal of Plant Physiology* 141(4): 463-469. Agrotechnol. Res. Inst., P.O. Box 17, 6700 AA Wageningen, Netherlands Antilles

Padhi, A. K., B. K. Sahoo, et al. (1993). Production potential, economics and energetics of upland cotton (*Gossypium hirsutum*) based intercropping systems under upland, rainfed situation. *Indian Journal of Agricultural Sciences* 63(3): 160-165. {a} National Agric. Res. Project, Kalimela, Orissa 764 047, India

Palit, P. (1993). Radiation and carbon use efficiency of field grown jute (*Corchorus* spp.) in relation to potential primary production. *Photosynthetica Prague* 28(3): 369-375. *Platn Physiol. Sect.*, Central Res. Inst. Jute and Allied Fibre, Barrackporre 743 101, India

Panizzi, A. R. and R. M. L. Alves (1993). Performance of nymphs and adults of the southern green stink bug (Heteroptera: Pentatomidae) exposed to soybean pods at different phenological stages of development. *Journal of Economic Entomology* 86(4): 1088-1093. {a} Centro Nacional Pesquisa Soja, Empresa Brasileira Pesquisa Agropecuaria, Caixa Postal 1061, Londrina, Parana 86001-970, Brasil

Patel, J. K. (1993). Response of rainfed upland cotton (*Gossypium hirsutum*) to triaccontanol and naphthalene acetic acid sprays. *Indian Journal of Agronomy* 38(1): 97-101. Water Management Project, Regional Agric. Res. Station, Indira Gandhi Krishi Vishwa Vidyalaya, Sarkanda, Bilaspur, Madhya Pradesh 495 001, India

Pettigrew, W. T., J. J. Heitholt, et al. (1993). Early season ethephon application effects on cotton photosynthesis. *Agronomy Journal* 85(4): 821-825. {a} USDA-ARS, Cotton Physiol. Genetics, P.O. Box 345, Stoneville, MS 38776, USA

Pettigrew, W. T., J. J. Heitholt, et al. (1993). Gas exchange differences and comparative anatomy among cotton leaf-type isolines. *Crop Science* 33(6): 1295-1299. {a} USDA-ARS, Cotton Physiol. Genet. Res., P.O. Box 345, Stoneville, MS 38776, USA

Pierce, M. L., M. Essenberg, et al. (1993). A comparison of the quantities of exopolysaccharide produced by *Xanthomonas campestris* pv. *malvacearum* in susceptible and resistant cotton cotyledons during early stages of infection. *Phytopathology* 83(3): 344-349. Dep. Biochem. Molecular Biology, Oklahoma State University, Stillwater, OK 74078-0454

- Polak, J., J. Chod, et al. (1993). Purification of hop mosaic virus and preparation of antiserum suitable for ELISA. *Ochrana Rostlin* 29(3): 161-167. {a} Res. Inst. Crop Production, Div. Phytomed., 161 06 Praha 6-Ruzyne, Czechoslovakia
- Pollak, P. E., T. Vogt, et al. (1993). Chalcone synthase and flavonol accumulation in stigmas and anthers of *Petunia hybrida*. *Plant Physiology Rockville* 102(3): 925-932. {a} Dep. Hortic., Prog. Plant Physiol. Wash. State Univ., Pullman, WA 99164-6414, USA
- Popham, P. L., S. M. Pike, et al. (1993). Water relation alterations observed during hypersensitive reaction induced by bacteria. *Plant Physiology Rockville* 103(4): 1243-1247. {a} Dep. Plant Pathol., Univ. Missouri, Columbia, MO 65211, USA
- Poswal, M. A. T. (1993). Evaluation of local and introduced cotton (*Gossypium hirsutum* L.) cultivars in the Nigerian savanna. *Tropical Agriculture* 70(3): 208-213. Dep. Agron., Univ. Fort Hare, Alice 5700, South Africa
- Prasad, M., B. L. Meena, et al. (1993). Effect of intercropping in upland cotton (*Gossypium hirsutum*) on growth and yield of component crops. *Indian Journal of Agronomy* 38(2): 342-344. Div. Agron., Indian Agric. Res. Inst., New Delhi 110 012, India
- Prasad, M. and R. Prasad (1993). Productivity and upland cotton (*Gossypium hirsutum*) genotypes under different levels of nitrogen and spacing. *Indian Journal of Agronomy* 38(4): 606-608. Div. Agron., Indian Agric. Res. Inst., New Delhi 110 012, India
- Puste, A. M. (1993). Response of tossa jute (*Corchorus olitorius*) to levels of nitrogen and inter-row spacing. *Indian Journal of Agricultural Sciences* 63(5): 268-271. Dep. Agron., Bidhan Chandra Krishi Viswavidyalaya, Mohanpur, West Bengal 741 235, India
- Rao, N. R. and J. D. Ownby (1993). Development of an ELISA for estimation of the copper nutritional status of wheat and cotton. *Plant and Soil* 155-156: 453-456. Dep. Botany, Oklahoma State Univ., Stillwater, OK 74078, USA
- Rasulov, B. K. (1993). Regulation of photosynthetic carbon dioxide exchange in leaves during development of water deficit in the radical area of cotton plants. *Fiziologiya Rastanii Moscow* 40(4): 668-675. Dep. Genet., Acad. Sci. Tadzh., Dushanbe, Tajikistan
- Ray, N. and V. K. Khaddar (1993). Effect of water stagnation at various growth stages of cotton plants grown under salt affected soil conditions. *Advances in Plant Sciences* 6(1 Suppl.): 125-136. J. N. Agricultural University, Indore Campus, Indore-452 001, India
- Ray, N. and S. C. Pandey (1993). The influence of restricted boll number on certain yield attributes and yield of cotton. *Proceedings of the National Academy of Sciences India Section B Biological Sciences* 63(2): 243-247. Jawaharlal Nehru Agric. Univ., Indore Campus, Indore-452 001, India
- Reddy, K. R., H. F. Hodges, et al. (1993). A temperature model for cotton phenology. *Biotronics* 22: 47-59. {a} Dep. Agron., Miss. State Univ., PO Box 9649, Mississippi State, MS 39762, USA
- Reddy, K. R., H. F. Hodges, et al. (1993). Temperature effects on pima cotton leaf growth. *Agronomy Journal* 85(3): 681-686. {a} Dep. Agron., Mississippi State Univ., Mississippi State, MS 39762, USA
- Reddy, V. R. (1993). Modeling mepiquat chloride-temperature interactions in cotton: The model. *Computers and Electronics in Agriculture* 8(3): 227-236. NRI Systems Res. Lab., Bld. 011A, Rm 165B, BARC-West, Beltsville, MD 20705-2350, USA
- Rikin, A., J. W. Dillwith, et al. (1993). Correlation between the circadian rhythm of resistance to extreme temperatures and changes in fatty acid composition in cotton seedlings. *Plant Physiology Rockville* 101(1): 31-36. {a} Dep. Bot., Okla. State Univ., Stillwater, OK 74078, USA

Ring, D. R. and J. H. Benedict (1993). Comparison of insect injury-cotton yield response functions and economic injury levels for *Helicoverpa zea* and *Heliothis virescens* (Lepidoptera: Noctuidae) in the lower gulf coast of Texas. *Journal of Economic Entomology* 86(4): 1228-1235. Texas Agricultural Res., Extension Cent., Route 2, Box 589, Corpus Christi, TX 78406, USA

Ring, D. R., J. H. Benedict, et al. (1993). Economic injury levels and development and application of response surfaces relating insect injury, normalized yield, and plant physiological age. *Environmental Entomology* 22(2): 273-282. Texas Agric. Res. and Extension Cent., Route 2, Box 589, Corpus Christi, TX 78406, USA

Ring, D. R., J. H. Benedict, et al. (1993). Cotton yield response to cotton fleahopper (Hemiptera: Miridae) infestations on the lower gulf coast of Texas. *Journal of Economic Entomology* 86(6): 1811-1819. {a} Texas Agric. Res. Extension Cent., Route 2, Box 589, Corpus Christi, TX 78406, USA

Rodriguez, K. R., D. G. Robertson, et al. (1993). Bahiagrass-cotton rotations and the management of *Meloidogyne incognita* and *Hoplolaimus galeatus*. *Nematropica* 23(2): 141-147. Dep. Plant Pathology, Auburn University, Auburn, AL 36849, USA

Rogers, G. S., L. Payne, et al. (1993). Nitrogen and phosphorus requirements of cotton and wheat under changing atmospheric CO₂ concentrations. *Plant and Soil* 155-156: 231-234. {a} Univ. Western Sydney, Hawkesbury, Bourke Street, Richmond, NSW 2753, Australia

Salisbury, C. D. and J. M. Chandler (1993). Interaction of cotton (*Gossypium hirsutum*) and velvetleaf (*Abutilon theophrasti*) plants for water is affected by their interaction for light. *Weed Science* 41(1): 69-74. {a} Texas Agric. Exp. Stn., Amrillo, TX 79106, USA

Santos, G. R. D., L. Zambolim, et al. (1993). Transmission of *Colletotrichum gossypii* var. *cephalosporioides* by cotton seeds produced on plants inoculated in different ages. *Summa Phytopathologica* 19(3-4): 177-180. Dep. Fitopatologia, Univ. Federal Vicosa, 36570-000-Vicosa, MG, Brazil

Savoy, B. R., D. L. Holshouser, et al. (1993). A system for morningglory (*Ipomoea* spp.) control in cotton (*Gossypium hirsutum* L.). *Weed Technology* 7(3): 615-618. {a} Sandoz Agro, Inc., P.O. Box KK, Gilroy, CA 95021-2348, USA

Schalk, J. M., B. M. Shepard, et al. (1993). Response of caterpillar pests and the parasite *Diadegma insulare* to collard cultivars and a pyrethrin insecticide. *Hortscience* 28(4): 308-310. {a} U.S. Dep. Agric., Agric. Res. Serv., U.S. Vegetable Lab., 2875 Savannah Highway, Charleston, SC 29414-5334, USA

Schmutz, A., T. Jenny, et al. (1993). Caffeic acid and glycerol are constituents of the suberin layers in green cotton fibres. *Planta Heidelberg* 189(3): 453-460. {a} Institut Botanische Biologie, Univ Freiburg, A. Gockelstrasse 3, CH-1700 Freiburg, Switzerland

Shapovalova, R. A., V. P. Kratenko, et al. (1993). Potato blight of cotton in Central Asia. *Doklady Rossiiskoi Akademii Sel'skokhozyaistvennykh Nauk*(3): 24-28. {a} Uzb. Inst. Genet., Tashkent, Uzbekistan

Sharpley, A. N. (1993). An innovative approach to estimate bioavailable phosphorus in agricultural runoff using iron oxide-impregnated paper. *Journal of Environmental Quality* 22(3): 597-601. USDA-ARS, Natl. Agric. Water Quality Lab., P.O. Box 1430, Durant, OK 74702-1430, USA

Sharpley, A. N., S. J. Smith, et al. (1993). Nitrogen and phosphorus fate from long-term poultry litter applications to Oklahoma soils. *Soil Science Society of America Journal* 57(4): 1131-1137. {a} USDA-ARS, ANatl. Agric. Water Quality Lab., P.O. Box 1430, Durant, OK 74702-1430, USA

She, J., J. Wu, et al. (1993). High frequency of plating efficiency and plant regeneration from protoplasts of cotton (*Gossypium hirsutum* L.). *Acta Genetica Sinica* 20(6): 536-540. Inst. Agrobiol. Genetics Physiol., Jiangsu Acad. Agric. Sci., Nanjing 210014, China

Sheck, A. L. and F. Gould (1993). The genetic basis of host range in *Heliothis virescens*: Larval survival and growth. *Entomologia Experimentalis et Applicata* 69(2): 157-172. Dep. Entomol., N.C. State Univ., Raleigh, NC 27642, USA

Shi, J., W. C. Mueller, et al. (1993). The inhibition of fungal growth in resistant cotton plants infected by *Fusarium oxysporum* f. sp. *vasinfectum*. *Journal of Phytopathology Berlin* 139(3): 253-260. Dep. Plant Sci., Univ. Rhode Island, Kingston, RI 02881, USA

Siddiqui, M. A. (1993). Heterobeltiosis for seed cotton yield and related character in *Hirsutum* cotton hybrids. *Journal of Maharashtra Agricultural Universities* 18(3): 403-405. Marathwada Agric. Univ., Parbhani 431 402, India

Simwat, G. S. and A. K. Dhawan (1993). Phytotoxic effect of spraying mixtures of systemic and contact insecticides on upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 63(6): 390-392. Dep. Entomol., Punjab Agric. Univ., Ludhiana 141 004, India

Singh, I. and G. S. Chouhan (1993). Effect of sowing time, cycocel spray and nitrogen fertilization on production potential of upland cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 38(1): 93-96. {a} Agric. Res. Station, Rajasthan Agric. Univ., Sumerpur 306 902, India

Singh, J., J. S. Gill, et al. (1993). Ovicidal effect of insecticides on eggs of gram podborer (*Helicoverpa armigera*) and spotted bollworm (*Earias vitella*). *Indian Journal of Agricultural Sciences* 63(12): 853-855. {a} Dep. Entomol., Punjab Agric. Univ., Ludhiana 141 004, India

Singh, M., N. B. Patil, et al. (1993). Improvement in fibre-bundle strength in medium-staple varieties of upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 63(11): 719-722. {a} Div. Genetics, IARI, New Delhi 110 012, India

Singh, M. and V. P. Singh (1993). Development of upland cotton (*Gossypium hirsutum*) for spring-summer cultivation. *Indian Journal of Agricultural Sciences* 63(10): 660-661. Div. Genetics, Indian Agric. Res. Inst., New Delhi 110 012, India

Singh, R., C. S. Praharaj, et al. (1993). Effect of irrigation and nitrogen regime on yield and yield-contributing characters of upland cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 38(4): 609-612. {a} Punjab Agric. Univ., Ludhiana 141 004, India

Soares, J. J. and A. A. D. De (1993). Influence of planting data and boll weevil *Anthonomus grandis* Boh. (Coleoptera: Curculionidae) attack on square and boll shedding in cotton. *Anais da Sociedade Entomologica do Brasil* 22(2): 253-258. {a} EMBRAPA, Centro Nacional Pesquisa Algodao, Caixa Postal 174, 58107-720 Campina Grande, Brazil

Soares, J. J. and P. T. Yamamoto (1993). Ovipositional behavior of *Anthonomus grandis* Boh. (Coleoptera: Curculionidae) under different levels of natural infestation. *Anais da Sociedade Entomologica do Brasil* 22(2): 333-339. {a} EMBRAPA, Centro Nacional Pesquisa Algodao, Caixa Postal 174, 58107-720 Campina Grande, Brazil

Sohu, R. S., G. S. Chahal, et al. (1993). A note on measurement of bloom rate in cotton. *Crop Improvement* 20(2): 219-221. Dep. Plant Breeding, Punjab Agric. Univ., Ludhiana, India

Stapelton, G. S., E. C. Murdock, et al. (1993). Postemergence control of cowpea (*Vigna unguiculata*) in cotton (*Gossypium hirsutum*). *Weed Technology* 7(2): 478-482. {a} Dep. Agron., Clemson Univ., Clemson, SC 29634, USA

Starr, J. L., C. M. Heald, et al. (1993). *Meloidogyne incognita* and *Rotylenchulus reniformis* and associated soil textures from some cotton production areas of Texas. *Journal of Nematology* 25(4 Suppl.): 895-899. {a} Tex. Agric. Exp. Stn., College Station, TX 77843, USA

- Steinitz, B., A. Navon, et al. (1993). Expression of insect resistance in in vitro-derived callus tissue infested with lepidopteran larvae. *Journal of Plant Physiology* 142(4): 480-484. {a} Dep. Plant Genetics, Agric. Res. Organization, Volcani Cent., Bet Dagan 50250, Israel
- Stoilova, A. (1993). Variability in reciprocal hybrids of *Gossypium hirsutum* L. and *G. barbadense* L. in BC-1F-1 and BC-1F-2 of F-3 backcrosses. *Genetika i Seleksiya* 26(5-6): 362-369. Inst. Cotton Durum Wheat, Chirpan 6200, Bulgaria
- Stoilova, A. and T. Lidanski (1993). Qualitative evaluation of reaction norm in cotton lines of interspecific origin (*Gossypium hirsutum* L. X *G. barbadense* L.). *Genetika i Seleksiya* 26(5-6): 391-395. {a} Inst. Cotton Durum Wheat, Chirpan 6200, Bulgaria
- Streeter, J. G. and S. O. Salminen (1993). Alterations in apoplastic and total solute concentrations in soybean nodules resulting from treatments known to affect gas diffusion. *Journal of Experimental Botany* 44(261): 821-828. {a} Dep. Agronomy, Ohio State Univ., Ohio Agric. Res. and Dev. Cent., Wooster, OH 44691, USA
- Suttle, J. C. and S. R. Abrams (1993). Abscission-promoting activities of abscisic acid and five abscisic acid analogs in cotton seedlings and explants. *Plant Growth Regulation* 12(1-2): 111-117. {a} USDA-ARS Northern Crop Sci. Lab., P.O. Box 5677, State Univ. Stn., Fargo, ND 58105-5677, USA
- Suttle, J. C. and J. F. Hulstrand (1993). Involvement of abscisic acid in ethylene-induced cotyledon abscission in cotton seedlings. *Plant Physiology Rockville* 101(2): 641-646. {a} United States Dep. Agric., Northern Crop Science Lab., P.O. Box 5677, State University Station, Fargo, ND 58105, USA
- Tamci, M. (1993). Phosphorus and zinc interaction in shoots of cotton plants. *Angewandte Botanik* 67(3-4): 138-140. Inst. Nuclear Sci., Univ. Ege, Bornova, Izmir, Turkey
- Tang, B., J. N. Jenkins, et al. (1993). F-2 hybrids of host plant germplasm and cotton cultivars: I. Heterosis and combining ability for lint yield and yield components. *Crop Science* 33(4): 700-705. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA
- Tang, B., J. N. Jenkins, et al. (1993). F-2 hybrids of host plant germplasm and cotton cultivars: II. Heterosis and combining ability for fiber properties. *Crop Science* 33(4): 706-710. {a} USDA-ARS, Crop Sci. Res. Lab., P.O. Box 5367, Mississippi State, MS 39762, USA
- Thomas, R. B., C. D. Reid, et al. (1993). Growth and maintenance components of leaf respiration of cotton grown in elevated carbon dioxide partial pressure. *Plant Cell and Environment* 16(5): 539-546. {a} Bot. Dep., Duke Univ., Durham, NC 27706, USA
- Timpa, J. D. and B. A. Triplett (1993). Analysis of cell-wall polymers during cotton fiber development. *Planta Heidelberg* 189(1): 101-108. {a} USDA, ARS, Southern Regional Res. Cent., P.O. Box 19687, New Orleans, LA 70179, USA
- Tiwari, R. J., K. Dwivedi, et al. (1993). Effect of gypsum on leaf-water potential of cottons (*Gossypium hirsutum*, *G. herbaceum* and *G. arboreum*) varieties grown in salt-affected vertisol of Madhya Pradesh. *Indian Journal of Agricultural Sciences* 63(11): 734-736. {a} Watershed Management Project, Coll. Agric., Sehore, Madhya Pradesh 466 001, India
- Tzen, J. T. C., Y. Z. Cao, et al. (1993). Lipids, proteins, and structure of seed oil bodies from diverse species. *Plant Physiology Rockville* 101(1): 267-276. {a} Dep. Bot. and Plant Sci., Univ. Calif., Riverside, CA 92521-0124, USA
- Ullrich, W. R., G. Kunz, et al. (1993). Role of ammonium accumulation in bacteria-induced hypersensitive and compatible reactions of tobacco and cotton plants. *Physiologia Plantarum* 89(3): 644-652. {a} Inst. Botanik, Technische Hochschule, Schnittpahnstrasse 3-5, D-64287 Darmstadt, Germany

- Vaidya, K. R. and M. M. Young (1993). Ethyl methanesulfonate induced variation in qualitative and quantitative characters of roselle (*Hibiscus sabdariffa* L.) (Malvaceae). *Revista Brasileira de Genetica* 16(2): 381-391. Dep. Botany, Univ. West Indies, Mona, Kingston 7, Jamaica
- Vencill, W. K., L. J. Giraud, et al. (1993). Soil moisture relations and critical period of *Cynodon dactylon* (L.) Pers. (coastal bermudagrass) competition in conservation-tillage cotton (*Gossypium hirsutum* L.). *Weed Research* 33(1): 89-96. {a} Dep. Agronomy, Univ. Georgia, Athens, GA, USA
- Walsh, J. D., M. S. Defelice, et al. (1993). Soybean (*Glycine max*) herbicide carryover to grain and fiber crops. *Weed Technology* 7(3): 625-632. {a} American Cyanamid Co., 1545 West Stonehenge, Sycamore, IL 61078, USA
- Walsh, J. D., M. S. Defelice, et al. (1993). Influence of tillage on soybean (*Glycine max*) herbicide carryover to grass and legume forage crops in Missouri. *Weed Science* 41(1): 144-149. {a} American Cyanamide Co., 1545 West Stonehenge, Sycamore, IL 60178, USA
- Wang, Y. and S. Y. Hu (1993). Cytoplasmic ultrastructural changes during microsporogenesis of *Gossypium hirsutum*: With emphasis on "cytoplasm reorganization". *Acta Botanica Sinica* 35(4): 255-260. Dep. Biol., Peking Univ., Beijing 100871, China
- Wankhade, S. T., S. G. Kharche, et al. (1993). Weed management in rainfed upland cotton (*Gossypium hirsutum*). *Indian Journal of Agronomy* 38(1): 102-104. Cotton Res. Unit, Punjabrao Krishi Vidyapeeth, Akola, Maharashtra 444 104, India
- Warner, D. A. and J. J. Burke (1993). Cool night temperatures alter leaf starch and photosystem II chlorophyll fluorescence in cotton. *Agronomy Journal* 85(4): 836-840. {a} USDA Plant Stress Water Conservation Res. Unit, Route 3, Box 215, Lubbock, TX 79401, USA
- Westerman, R. B., D. S. Murray, et al. (1993). Hogpotato (*Hoffmanseggia glauca*) control with herbicides and rotational crop response. *Weed Technology* 7(3): 650-656. {a} Dep. Agron., Okla. State Univ., Stillwater, OK 74078, USA
- Xie, W., N. L. Trolinder, et al. (1993). Cool temperature effects on cotton fiber initiation and elongation clarified using in vitro cultures. *Crop Science* 33(6): 1258-1264. {a} Dep. Biol. Sci., Box 43131, Tex. Tech Univ. Lubbock, TX 79409, USA
- Yoon, I. S. and B. G. Kang (1993). Sensitivity changes of auxin transport system in maize coleoptile segments. *Korean Journal of Botany* 36(1): 59-66. {a} Dep. Biol., Seoul Natl. Univ., Seoul, North Korea
- York, A. C. (1993). Peanut response to fluometuron applied to a preceding cotton crop. *Peanut Science* 20(2): 111-114. Dep. Crop Sci., N.C. State Univ., Raleigh, NC 27695-7620, USA
- York, A. C. and J. W. Wilcut (1993). Insecticides do not affect cotton (*Gossypium hirsutum*) response to imazaquin and imazethapyr. *Weed Science* 41(2): 269-280. {a} Dep. Crop Sci., N.C. State Univ., Raleigh, NC 27695-7620, USA
- Zekri, S. and C. Romero (1993). Public and private compromises in agricultural water management. *Journal of Environmental Management* 37(4): 281-290. {a} Dep. Agric. Econ. Manage. Stat., Cordoba Univ., Spain
- Zeng, Y. S., B. C. Ye, et al. (1993). Evidence for gene-for-gene complementarity: A novel specific elicitor from *Fusarium oxysporum* f. sp. *vasinfectum*. *Chinese Journal of Botany* 5(2): 97-101. Shanghai Inst. Biochem., Academia Sinica, Shanghai 20031, China
- Zhang, J., M. E. Mace, et al. (1993). Production and fungitoxicity of the terpenoid phytoalexins in cotton: Inoculated with *Fusarium oxysporum* f. sp. *vasinfectum*. *Journal of Phytopathology Berlin* 139(3): 247-252. USDA, ARS, Southern Crops Res. Lab., Cotton Pathol. Res. Unit, Route 5, Box 805, College Station, TX 77845, USA

- Zhang, J. F., J. Z. Sun, et al. (1993). Effect of frego bract gene on economic properties of cotton. *Acta Agronomica Sinica* 19(4): 315-320. Huazhong Agric. Univ., Wuhan 430070, Japan
- Zhong, H. and A. Lauchli (1993). Changes of cell wall composition and polymer size in primary roots of cotton seedlings under high salinity. *Journal of Experimental Botany* 44(261): 773-778. {a} Dep. Integrative Biol., Univ. California, Berkeley, CA 94720, USA
- Zhong, H. and A. Lauchli (1993). Spatial and temporal aspects of growth in the primary root of cotton seedlings: Effects of sodium chloride and calcium chloride. *Journal of Experimental Botany* 44(261): 763-771. {a} Dep. Integrative Biol., Univ. California, Berkeley, CA 94720, USA
- Abdullaev, A. A., B. B. Dzhumaev, et al. (1992). Combined action of environmental factors of photosynthetic carbon metabolism in cotton leaves. *Fiziologiya Rastanii Moscow* 39(2): 231-238. {a} Inst. Physiol. Biophys. Plant, Acad. Sci. Tadjh., Dushanbe, Tadjh
- Adeoti, A. A. and A. M. Emechebe (1992). The alternative hosts of *Coniella musaiaensis*. *Nigerian Journal of Botany* 5: 173-176. Dep. Crop Protection, Inst. Agric. Res., Ahmadu Bello University, Samaru, Zaria, Nigeria
- Ali, A. and M. J. Gaylor (1992). Effects of temperature and larval diet on development of the beet armyworm (Lepidoptera: Noctuidae). *Environmental Entomology* 21(4): 780-786.
- Allen, S. J., A. S. Hodgson, et al. (1992). The effects of various agronomic practices on the incidence of bacterial blight of cotton. *Plant Protection Quarterly* 7(2): 55-58.
- Almeida, I. M. G., L. O. S. Beriam, et al. (1992). Isolation and characterization of *Xanthomonas campestris* pv. *cyamopsidis* in Brazil and reaction of guar genotypes to the bacterium. *Summa Phytopathologica* 18(3-4): 255-261. {a} Instituto Biologico, Caixa Postal 70, 13001-Campinas, SP, Brazil
- Artiola, J. F. and I. L. Pepper (1992). Denitrification activity in the root zone of a sludge-amended desert soil. *Biology And Fertility Of Soils* 13(4): 200-205.
- Augustinussen, E. (1992). Effect of seed rate, nitrogen fertilization, growth regulation and harvest method on yield of fiber flax. *Tidsskrift for Planteavl* 96(5): 433-440. Statens Planteavlsfors., Landbrugscent., Afdeling Industriplanter og Froavl, 4000 Roskilde, Denmark
- Auld, B. A., H. E. Talbot, et al. (1992). Host range of three isolates of *Alternaria zinniae*, a potential biocontrol agent for *Xanthium* spp. *Plant Protection Quarterly* 7(3): 114-116. NSW Agric., Agric. Res. and Vet. Cent., Forest Road, Orange 2800, Australia
- Averina, N. G., I. D. Shegai, et al. (1992). Biosynthesis of chlorophyll in the cotyledonary leaves of a chlorophyll-deficient cotton mutant. *Fiziologiya Rastanii Moscow* 39(1): 85-92. {a} Inst. Photobiol., Acad. Sci. B., Minsk, Beloruss
- Babadzhanov, M. A. and Y. S. Nasyrov (1992). A multienzyme complex of key photosynthetic enzymes. *Fiziologiya Rastanii Moscow* 39(4): 753-759. {a} Tadjh. State Univ., Dushanbe, Tadjh
- Basra, A. S., R. S. Sarlach, et al. (1992). Hormonal effects on partitioning of carbon-14 sucrose in cotton fibres. *Acta Physiologiae Plantarum* 14(3): 137-142. Dep. Botany, Punjab Agricultural Univ., Ludhiana 141 004, India
- Bayley, C., N. Trolinder, et al. (1992). Engineering 2,4-D resistance into cotton. *Theoretical And Applied Genetics* 83(5): 645-649.
- Bevers, S. J. and J. E. Slosser (1992). Assessing cost effectiveness of planting dates and insecticide chemicals in Texas dryland cotton production. *Journal Of Production Agriculture* 5(3): 374-377.

- Bolger, T. P., D. R. Upchurch, et al. (1992). Temperature effects on cotton root hydraulic conductance. *Environmental And Experimental Botany* 32(1): 49-54.
- Boora, K. S., G. P. Lodhi, et al. (1992). Genetical architecture of yield and its components characters in alfalfa (*Medicago sativa* L.). *Annals of Biology Ludhiana* 8(2): 174-177. {a} CCS Haryana Agric. Univ., Hisar-125 004, India
- Bourland, F. M., D. M. Oosterhuis, et al. (1992). Concept for monitoring the growth and development of cotton plants using main-stem node counts. *Journal of Production Agriculture* 5(4): 532-538. {a} Dep. Agronomy, Univ. Arkansas, Fayetteville, Ariz. 72701
- Bradford, S. and J. Letey (1992). Cyclic and blending strategies for using nonsaline and saline waters for irrigation. *Irrigation Science* 13(3): 123-128. {a} Dep. Soil Environ. Sci., Univ. Calif., Riverside, Calif. 92521, USA
- Bradford, S. and J. Letey (1992). Simulated effects of water table and irrigation scheduling as factors in cotton production. *Irrigation Science* 13(3): 101-107. {a} Dep. Soil Environ. Sci., Univ. Calif., Riverside, Calif. 92521, USA
- Brugnoli, E. and O. Bjorkman (1992). Growth of cotton under continuous salinity stress: Influence on allocation pattern, stomatal and non-stomatal components of photosynthesis and dissipation of excess light energy. *Planta* 187(3): 335-347.
- Bruin, J., M. Dicke, et al. (1992). Plants are better protected against spider-mites after exposure to volatiles from infested conspecifics. *Experientia* 48(5): 525-529.
- Bryant, K. J., S. M. Masud, et al. (1992). Profit, cost, and soil erosion implications of dryland crop production systems: Texas High Plains. *Journal Of Production Agriculture* 5(3): 307-312.
- Byzova, M. V., A. S. Kraev, et al. (1992). Application of reverse transcriptase-polymerase chain reaction (RT-PCR) for identification of CHS gene from two cotton species specifically expressed in the petals. *Molekulyarnaya Biologiya Moscow* 26(4): 911-915. Cent. "Bioeng.", Acad. Sci. Russ., Moscow 117984, Russia
- Byzova, M. V., A. S. Kraev, et al. (1992). Molecular characterization of chalcone synthase gene families from two cotton species, using polymerase chain reaction. *Molekulyarnaya Biologiya* 26(2): 432-440.
- Carmi, A., Z. Plaut, et al. (1992). Establishment of shallow and restricted systems in cotton and its impact on plant response to irrigation. *Irrigation Science* 13(2): 87-91.
- Cassman, K. G., B. A. Roberts, et al. (1992). Cotton response to residual fertilizer potassium on vermiculitic soil: Organic matter and sodium effects. *Soil Science Society Of America Journal* 56(3): 823-830.
- Chapman, K. D. and R. N. Trelease (1992). Characterization of membrane proteins in enlarging cottonseed glyoxysomes. *Plant Physiology And Biochemistry* 30(1): 1-10.
- Choi, B. H., J. I. Lee, et al. (1992). Direct and indirect effects and determination coefficients of agronomic traits on seed cotton yield for Upland cotton culture after naked barley harvest. *Research Reports of the Rural Development Administration Suweon* 34(2 Upland and Indus. Crops): 59-67. Crop Exp. Stn., RDA, Suwon, North Korea
- Chu, C. C., T. J. Henneberry, et al. (1992). Effect of cotton defoliant on leaf abscission, immature bolls, and lint yields in a short-season production system. *Journal Of Production Agriculture* 5(2): 268-272.
- Constable, G. A., I. J. Rochester, et al. (1992). Cotton yield and nitrogen requirement is modified by crop rotation and tillage method. *Soil & Tillage Research* 23(1-2): 41-59.

- Cook, C. G. and Z. K. M. El (1992). Cotton seedling and first-bloom plant characteristics: Relationships with drought-influenced boll abscission and lint yield. *Crop Science* 32(6): 1464-1467. {a} USDA-ARS, Subtropical Agric. Res. Lab., Weslaco, TX 78596
- Culp, T. W. and C. C. Green (1992). Performance of obsolete and current cultivars and pee dee germplasm lines of cotton. *Crop Science* 32(1): 35-41.
- Dejoode, D. R. and J. F. Wendel (1992). Genetic diversity and origin of the Hawaiian Islands cotton, *Gossypium tomentosum*. *American Journal of Botany* 79(11): 1311-1319. {a} Dep. Botany, Iowa State Univ., Ames, Iowa 50011
- Deshmukh, V. N., Y. S. Guhe, et al. (1992). Response of wheat (*Triticum aestivum*) to various levels of irrigation with saline water under different crop sequences in Vertisol. *Indian Journal of Agricultural Sciences* 62(11): 737-740. {a} Dep. Agricultural Chemistry and Soil Sci., Punjabrao Krishi Vidyapeeth, Akola, Maharashtra 444 104, India
- Dever, J. K. and J. R. Gannaway (1992). Relative fiber uniformity between parent and F-1 and F-2 generations in cotton. *Crop Science* 32(6): 1402-1408. {a} Texas A and M Univ. Agric. Res. Ext. Cent., Rt. 3, Box 219, Lubbock, Texas 79401
- Dhawan, A. K., G. S. Simwat, et al. (1992). Field evaluation of asyemethrin (Chinmix) for bollworm control on cotton. *Indian Journal Of Plant Protection* 20(1): 24-26.
- Ellsworth, P. C. and J. R. J. Bradley (1992). Comparative damage potential and feeding dynamics of the European corn borer (Lepidoptera: Pyralidae) and cotton bollworm (Lepidoptera: Noctuidae) on cotton bolls. *Journal Of Economic Entomology* 85(2): 402-410.
- Endrizzzi, J. E. and D. T. Ray (1992). Mapping of the *chl-1*, *R-1*, *yg-1*, and *Dw* loci in the long arm of chromosome 16 of cotton. *Journal Of Heredity* 83(1): 1-5.
- Essenberg, M., M. L. Pierce, et al. (1992). A method for determining phytoalexin concentrations in fluorescent, hypersensitively necrotic cells in cotton leaves. *Physiological and Molecular Plant Pathology* 41(2): 101-109. {a} Dep. Biochem., Oklahoma State Univ., Stillwater, OK 74078, USA
- Essenberg, M., M. L. Pierce, et al. (1992). Development of fluorescent, hypersensitively necrotic cells containing phytoalexins adjacent to colonies of *Xanthomonas campestris* pv. *malvacearum* in cotton leaves. *Physiological and Molecular Plant Pathology* 41(2): 85-99. {a} Dep. Biochem., Oklahoma State Univ., Stillwater, OK 74078
- Estesen, B. J., N. A. Buck, et al. (1992). Residual life and toxicity to honey bees (Hymenoptera: Apidae) of selected insecticides applied to cotton in Arizona. *Journal Of Economic Entomology* 85(3): 700-709.
- Fargues, J. and N. K. Maniania (1992). Variation in susceptibility of *Spodoptera littoralis* (Noctuidae) to *Nomuraea rileyi* (Hyphomycetes). *Entomophaga* 37(4): 545-554. {a} I.N.R.A., Stn. Recherches Lutte Biol., La Miniere, 78285 Guyancourt Cedex, France
- Fernandez, C. J., J. T. Cothren, et al. (1992). Carbon and water economies of well-watered and water-deficient cotton plants treated with mepiquat chloride. *Crop Science* 32(1): 175-180.
- Gal, p. M. I. and L. G. Portenko (1992). Study of the pathogenicity of *Thielaviopsis basicola* to *Gossypium hirsutum* and *Gossypium barbadense*. *Izvestiya Akademii Nauk Respubliki Tadjhikistan Otdelenie Biologicheskikh Nauk*(3): 26-29. Dep. Gen. Genet. Cotton, Acad. Sci. Tadjh., Dushanbe, Tajikistan
- Giyasov, T. D., A. I. Klyuikova, et al. (1992). Nitrate reductase in germinating cotton seeds: Effects of light and substrate availability. *Fiziologiya Rastenii Moscow* 39(4): 807-814. {a} Div. Biochem. Biol. Fac., Tadz. State Univ., Dushanbe, Tadz

- Gossett, B. J., E. C. Murdock, et al. (1992). Resistance of Palmer amaranth (*Amaranthus palmeri*) to the dinitroaniline herbicides. *Weed Technology* 6(3): 587-591. {a} Dep. Agron, Clemson Univ., Clemson, S.C. 29634-0359
- Goulart, A. C. P. (1992). Effect of fungicides on the control of pathogens in cotton (*Gossypium hirsutum*) seeds. *Summa Phytopathologica* 18(2): 173-177. EMBRAPA-UEPAE Dourados, Caixa Postal 661, 79800-Dourados, MS
- Guinn, G. and D. L. Brummett (1992). Influence of defruiting on the abscisic acid and indole-3-acetic acid contents of cotton leaves. *Field Crops Research* 28(3): 257-262.
- Gupta, M. P. (1992). Amylase activity in cotton. *Indian Journal Of Agricultural Research* 25(3): 141-148.
- Gururajan, K. N., S. Henry, et al. (1992). Yield and seed-quality characteristics of varieties of glandless cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 62(5): 316-318.
- Harley, P. C., R. B. Thomas, et al. (1992). Modelling photosynthesis of cotton grown in elevated carbon dioxide. *Plant Cell And Environment* 15(3): 271-282.
- Hartung, W., E. W. Weiler, et al. (1992). Auxin and cytokinins in the apoplastic solution of dehydrated cotton leaves. *Journal of Plant Physiology* 140(3): 324-327. {a} Lehrstuhl Bot. I, Julius von Sachs Inst. Biowissenschaften Mittlerer Dallenbergweg 64, D-8700 Wuerzburg, Germany
- Hauck, C., S. Mueller, et al. (1992). A germination stimulant for parasitic flowering plants from *Sorghum bicolor*, a genuine host plant. *Journal Of Plant Physiology* 139(4): 474-478.
- Hedin, P. A., W. L. Parrott, et al. (1992). Relationships of glands, cotton square terpenoid aldehydes, and other allelochemicals to larval growth of *Heliothis virescens* (Lepidoptera; Noctuidae). *Journal Of Economic Entomology* 85(2): 359-364.
- Heilman, J. L., K. J. McInnes, et al. (1992). Evaporation from ridge-tilled soil covered with herbicide-killed winter wheat. *Soil Science Society of America Journal* 56(4): 1278-1286. {a} Dep. Soil and Crop Sci., Tex. A and M Univ., College Station, Tex. 77843
- Heitholt, J. J., W. T. Pettigrew, et al. (1992). Light interception and lint yield of narrow-row cotton. *Crop Science* 32(3): 728-733.
- Iida, Y., K. I. Watabe, et al. (1992). Effects of abscisic acid on the induction of desiccation tolerance in carrot somatic embryos. *Journal of Plant Physiology* 140(3): 356-360. {a} Gene Exp. Cent., Univ. Tsukuba, Tsukuba, Ibaraki 305, Japan
- Iqbal, M. M., S. K. Hussain, et al. (1992). Response of cotton to applied phosphorus in two soil series. *Sarhad Journal of Agriculture* 8(6): 689-696. {a} Nuclear Inst. Food Agric., Peshawar
- Ishag, H. M. (1992). Effects of foliar micronutrient fertilizers on the yield of irrigated cotton on the vertisols of the Sudan Gezira. *Experimental Agriculture* 28(3): 265-271.
- Itoh, J., T. Shioya, et al. (1992). Improvement of soil properties by cultivation of tropical leguminous crops in heavy clayey paddy fields. *Bulletin of the Hokuriku National Agricultural Experiment Station*(34): 27-41. {a} Joetsu Office, Cisso-Asahi Fertilizer Co. Ltd., Minami-honcho, Joetsu, Niigata 943, Japan
- John, M. E. and L. J. Crow (1992). Gene expression in cotton (*Gossypium hirsutum* L.) fiber: Cloning of the mRNAs. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 89(13): 5769-5773.
- Kaas, J. P., G. W. Elzen, et al. (1992). Flight responses in a wind tunnel of the parasitoid *Microplitis croceipes* to three spring and three summer host plants of *Heliothis* spp. *Entomologia Experimentalis Et Applicata* 63(3): 207-212.

Kalra, V. K. and P. D. Sharma (1992). Standardization of sampling technique and sample size for recording bollworm incidence in green bolls of cotton: A preliminary study. *Crop Research Hisar* 5(3): 541-544. Dep. Entomol., CCS Haryana Agric. Univ., Hisar-125 004, India

Karpilova, I. M. S. and S. M. Gaziyants (1992). Isolation of active ribulose-1,5-biphosphate carboxylase from the cotton species *Gossypium hirsutum* L. and *Gossypium barbadense* L. *Biopolimery i Kletka* 8(4): 30-38. Sci.-Eng. Cent. Autom. Biotech. Syst. "Sonar", V.M. Glushkov Inst. Cybern., Acad. Sci. Ukr., Kiev, Ukraine

Kasperbauer, M. J. and P. G. Hunt (1992). Cotton seedling morphogenic responses to FR/R ratio reflected from different colored soils and soil covers. *Photochemistry and Photobiology* 56(5): 579-584. {a} USDA, ARS, Coastal Plains Res. Center, Florence, SC 29504-3039

Kishor, P. B. K., J. D. Rao, et al. (1992). Activity of wall-bound enzymes in callus cultures of *Gossypium hirsutum* L. during growth. *Annals Of Botany* 69(2): 145-149.

Kleifeld, Y., T. Blumenfeld, et al. (1992). Control of purple nutsedge (*Cyperus rotundus*) in cotton with benfuresate. *Phytoparasitica* 20(1): 37-46.

Kloth, R. H. (1992). Variability of malate dehydrogenase among cotton cultivars with differing fiber traits. *Crop Science* 32(3): 617-621.

Kohel, R. J., E. V. Narbuth, et al. (1992). Fiber development of ligan lintless-2 mutant of cotton. *Crop Science* 32(3): 733-735.

Kolganova, T. V., D. K. Shrivastava, et al. (1992). Callogenesis and regeneration of the cotton *Gossypium hirsutum* L., 108-F. *Fiziologiya Rastenii Moscow* 39(2): 365-370. Inst. Mol. Genet., Acad. Sci. Russ., Moscow, Russia

Koraddi, V. R., P. Basavanna, et al. (1992). Response of rainfed cotton to sowing dates, spacings and fertilizer levels. *Journal of Maharashtra Agricultural Universities* 17(1): 69-72. {a} Agric. Research Station, Dharwad-580 007, India

Koroleva, N. S. and O. K. Yuldashev (1992). Cotton tissue culture: Communication II: Optimization of the phytohormonal composition of a medium for cotton callus formation. *Izvestiya Akademii Nauk Respubliki Tadjikistan Otdelenie Biologicheskikh Nauk*(1): 39-43. Inst. Physiol. Biophys. Plant, Acad. Sci. Tadjh., Dushanbe, Tajikistan

Koroleva, N. S. and O. K. Yuldashev (1992). Cotton tissue culture: Communication 4. Isolation, cultivation and protoplast fusion in *Gossypium hirsutum* and *Gossypium arboreum*. *Izvestiya Akademii Nauk Respubliki Tadjikistan Otdelenie Biologicheskikh Nauk*(2): 41-45. Inst. Physiol. Biophys. Plants, Acad. Sci. Tadjh., Dushanbe, Tajikistan

Kotova, V. V. (1992). Cotton disease caused by *Pythium ultimum* Trow var. *ultimum*. *Mikologiya i Fitopatologiya* 26(4): 284-288. All-Union Inst. Plant Prot., St. Petersburg, Russia

Lackey, B. A., B. A. Jaffee, et al. (1992). Sporulation of the nematophagous fungus *Hirsutiella rhossiliensis* from hyphae produced in vitro and added to soil. *Phytopathology* 82(11): 1326-1330. {a} Dep. Nematol., Univ. Calif., Davis, Calif. 95616-8668

Lagoke, S. T. O., K. O. Adejonwo, et al. (1992). Herbicide evaluation in the hoe-farming system cotton production in Northern Nigeria. *Tropical Pest Management* 38(3): 286-294.

Lesch, S. M., J. D. Rhoades, et al. (1992). Mapping soil salinity using calibrated electromagnetic measurements. *Soil Science Society Of America Journal* 56(2): 540-548.

Lewis, J. A., R. D. Lumsden, et al. (1992). Suppression of damping-off of peas and cotton in the field with composted sewage sludge. *Crop Protection* 11(3): 260-266.

- Liang, Z. L., R. Q. Jiang, et al. (1992). Studies on chromosome behaviour of F-1 and fertility restoration in hybrid of *Gossypium hirsutum* times *Gossypium bickii*. *Acta Botanica Sinica* 34(12): 931-936. Inst. Genet., Acad. Sinica, Beijing 100101, China
- Liu, C., J. Shun, et al. (1992). In vitro interspecific fertilization, embryo development and formation of hybrid seedlings between *Gossypium hirsutum* and *Gossypium arboreum*. *Euphytica* 60(2): 79-88.
- Ljubescic, N. and D. Matijevic (1992). The effect of amitrole on the pigment composition and ultrastructure of chromoplasts of tulip tree flowers. *Acta Botanica Croatica* 51: 13-19. Inst. Ruder Boskovic, P.O.B. 1016, 41001 Zagreb, Hrvatska, Croatia
- Maia, V. G. (1992). Effects of different rates and sources of nitrogen on the fecundity of *Tetranychus urticae* (Koch, 1836) in IAC 20 cotton plants (*Gossypium hirsutum* L.). *Anais da Sociedade Entomologica do Brasil* 21(3): 347-356. {a} Ex-Bolsista da FAPESP
- Majumdar, D. K. and S. K. Roy (1992). Response of summer sesame (*Sesamum indicum*) to irrigation, row spacing and plant population. *Indian Journal of Agronomy* 37(4): 758-762. {a} Dep. Agron., Palli Siksha Bhavana Visva-Bharati, Sriniketan, West Bengal 731 236
- Makhbubov, M. and P. D. Usmanov (1992). New wilt-resistant *Gossypium hirsutum* cultivar 1541. *Izvestiya Akademii Nauk Respubliki Tadjikistan Otdelenie Biologicheskikh Nauk*(2): 36-40. Dep. Genet., Acad. Sci. Tadjh., Dushanbe, Tajikistan
- Masood, S., M. Arshad, et al. (1992). Effect of transplanting and dibbling on the yield and yield components of cotton (*Gossypium hirsutum* L.). *Pakistan Journal Of Scientific And Industrial Research* 35(4): 142-144.
- McConnell, J. S., W. H. Baker, et al. (1992). Response of cotton to nitrogen fertilization and early multiple applications of mepiquat chloride. *Journal Of Plant Nutrition* 15(4): 457-468.
- Meiri, A., H. Frenkel, et al. (1992). Cotton response to water and salinity under sprinkler and drip irrigation. *Agronomy Journal* 84(1): 44-50.
- Meisner, J., M. Klein, et al. (1992). Effect of Margosan-O on the development of the leafhopper *Asymmetrasca decedens*. *Phytoparasitica* 20(1): 15-24.
- Meshram, M. K. and S. Raj (1992). Effect of bacterial blight infection at different stages of crop growth on intensity and seed cotton yield under rainfed conditions. *Indian Journal Of Plant Protection* 20(1): 54-57.
- Micinski, S., P. D. Colyer, et al. (1992). Cotton white flower counts and yield with and without early-season pest control. *Journal Of Production Agriculture* 5(1): 126-130.
- Mohan, P., M. G. Bhat, et al. (1992). Variability for biomass and harvest index in Asiatic (*Gossypium arboreum* L.) and American (*Gossypium hirsutum* L.) cottons. *Advances in Plant Sciences* 5(1): 100-105. Central Inst. Cotton Res., Nagpur-440 001, Maharashtra
- Mohan, P., P. Singh, et al. (1992). Variability for gossypol glands in upland cotton (*Gossypium hirsutum* L.). *Advances In Plant Sciences* 4(1): 165-170.
- Mohod, V. K., S. B. Parde, et al. (1992). Assessment of cotton isolines under various background of micronaire values. *Asian Journal of Plant Science* 4(2): 23-26. Dep. Botany, Punjabrao Krishi Vidyapeeth, Akola-444 005, India
- Morgan, P. W., C. J. He, et al. (1992). Intact leaves exhibit a climacteric-like rise in ethylene production before abscission. *Plant Physiology Rockville* 100(3): 1587-1590. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, Texas 77843

Mueller, T. C., R. E. Jones, et al. (1992). Comparison of PRZM and GLEAMS computer model predictions with field data for alachlor, metribuzin and norflurazon leaching. *Environmental Toxicology And Chemistry* 11(3): 427-436.

Mullins, G. L. and C. H. Burmester (1992). Uptake of calcium and magnesium by cotton grown under dryland conditions. *Agronomy Journal* 84(4): 564-569.

Mulrooney, J. E., W. L. Parrott, et al. (1992). Response of cotton to damage by tobacco budworm (Lepidoptera: Noctuidae): Changes in fruit distribution. *Journal Of Economic Entomology* 85(3): 981-986.

Naik, R. M., A. R. Dhage, et al. (1992). Differential carbon monoxide sensitivity of cytochrome C oxidase in the leaves of C-3 and C-4 plants. *Plant Physiology* 98(3): 984-987.

Narasimharao, P. and Y. Narasimharao (1992). Quality of effluent water discharged from paperboard industry and its effect on alluvial soil and crops. *Indian Journal Of Agricultural Sciences* 62(1): 9-12.

Nazarenko, L. V. (1992). Effect of sodium chloride on ribulosebisphosphate carboxylase of Euglena cells. *Fiziologiya Rastanii Moscow* 39(4): 748-752. K.A. Timiryazev Inst. Plant Physiol., Acad. Sci. Russ., Moscow, Russia

Ogunwolu, E. O. (1992). Field infestation and damage to soybean and cowpea by pod-sucking bugs in Benue State, Nigeria. *Insect Science and its Application* 13(6): 801-805. Dep. Crop Production, Univ. Agric., P.M.B. 2373, Makurdi, Nigeria

Ohana, P., D. P. Delmer, et al. (1992). Beta-Furfuryl-beta-glucoside: An endogenous activator of higher plant UDP-glucose:(1-3)-beta-glucan synthase: Biological activity, distribution and in vitro synthesis. *Plant Physiology* 98(2): 708-715.

Oliveira, S. H. F., B. C. Barros, et al. (1992). Effect of fungicides on the control of bean foliar diseases and seed quality. *Summa Phytopathologica* 18(2): 178-184. {a} Inst. Biol., Caixa Postal 70, 13001-Campinas, SP

Orgaz, F., L. Mateos, et al. (1992). Season length and cultivar determine the optimum evapotranspiration deficit in cotton. *Agronomy Journal* 84(4): 700-706.

Palmer, R. L. and W. M. Dugger (1992). Recovery of glucose from glucolipids in cell wall glucans in cotton fibres. *Phytochemistry* 31(8): 2631-2633.

Pandurangi, R. B., S. G. Wankhade, et al. (1992). Response of cotton (*Gossypium hirsutum* L.) to nitrogen and phosphorus grown under rainfed conditions. *Crop Research* 5(1): 54-58.

Paplomatas, E. J., D. M. Bassett, et al. (1992). Incidence of verticillium wilt and yield losses of cotton cultivars (*Gossypium hirsutum*) based on soil inoculum density of *Verticillium dahliae*. *Phytopathology* 82(12): 1417-1420. {a} Benaki Phytopathological Inst., Kifissia-Athens, Greece

Paris, Q. (1992). The return of von Liebig's "law of the minimum". *Agronomy Journal* 84(6): 1040-1046. Dep. Agric. Economics, Univ. Calif., Davis, Calif. 95616

Patil, F. B., Y. M. Shinde, et al. (1992). Heterosis in multiple environments for yield components and its relation with genetic divergence in cotton. *Indian Journal Of Genetics & Plant Breeding* 51(1): 118-124.

Patil, J. R., B. S. Janagoudar, et al. (1992). Effect of 2, 4-D on growth of cotton (*Gossypium hirsutum* L.) and possible recovery by antidotes. *Indian Journal of Plant Physiology* 35(4): 351-356. Dep. Crop Physiology, University Agric. Sci., Dharwad-580005

Percy, R. G. and E. L. Turcotte (1992). Interspecific hybrid fiber characteristics of cotton altered by unconventional *Gossypium barbadense* L. fiber genotypes. *Crop Science* 32(6): 1437-1441. {a} USDA-ARS, Maricopa Agric. Cent., 38760 W. Smith-Enke Road, Maricopa, Ariz. 85239

- Petersen, K. L., M. Fuchs, et al. (1992). Computing transpiration of sunlit and shaded cotton foliage under variable water stress. *Agronomy Journal* 84(1): 91-97.
- Pettigrew, W. T., J. J. Heitholt, et al. (1992). Early season floral bud removal and cotton growth, yield, and fiber quality. *Agronomy Journal* 84(2): 209-214.
- Polley, H. W., H. B. Johnson, et al. (1992). Determination of root biomasses of three species grown in a mixture using stable isotopes of carbon and nitrogen. *Plant And Soil* 142(1): 97-106.
- Powell, D. A. and T. S. J. Bellows (1992). Development and reproduction of two populations of *Eretmocerus* spp. (Hymenoptera: Aphelinidae) on *Bemisia tabaci* (Homoptera: Aleyrodidae). *Environmental Entomology* 21(3): 651-658.
- Prisco, J. T., C. R. B. Haddad, et al. (1992). Hydration-dehydration seed pre-treatment and its effects on seed germination under water stress conditions. *Revista Brasileira De Botanica* 15(1): 31-35.
- Pundarikakshudu, R., N. J. Mudholkar, et al. (1992). Effect of comparative efficiency of planting methods and soil depth on rainfed cotton (*Gossypium hirsutum*) in vertisols. *Indian Journal Of Agricultural Sciences* 62(4): 263-267.
- Qureshi, M. B., M. S. Awan, et al. (1992). Population dynamics, dispersal pattern and host records of dusky cotton bug, *Oxycarenus laetus* Kirby (Lygaeidae: Heteroptera) in Sindh. *Pakistan Journal of Zoology* 24(3): 223-225. Dep. Zoology, Univ. Sindh, Jamshoro
- Radin, J. W., L. L. Reaves, et al. (1992). Yield enhancement in cotton by frequent irrigations during fruiting. *Agronomy Journal* 84(4): 551-557.
- Raju, A. R. and S. G. Kharche (1992). Effects of sowing time and plant populations on fibre quality of cotton hybrids/varieties. *Agricultural Science Digest* 12(4): 189-192. {a} Cent. Inst. Cotton Res., Post Bag No. 225, G.P.O. Nagpur-440 001, India
- Rana, N. S. and H. B. Singh (1992). Uptake of nitrogen, phosphorus and potash by potato (*Solanum tuberosum*), mustard (*Brassica juncea*) and their intercropping system. *Indian Journal of Agronomy* 37(4): 666-668. Amar Singh College, Lakhaoti, Meerut Univ., Uttar Pradesh
- Rasulov, S. (1992). Ripening rate of cotton bolls as a function of the location of fruit elements and effect of synthetic growth regulators. *Izvestiya Akademii Nauk Respubliki Tadjikistan Otdelenie Biologicheskikh Nauk*(1): 46-50. K.A. Timiryazev Inst. Plant Physiol., Acad. Sci. Russ., Moscow, Russia
- Reddy, K. R., H. F. Hodges, et al. (1992). Temperature effects on cotton fruit retention. *Agronomy Journal* 84(1): 26-30.
- Reddy, K. R., V. R. Reddy, et al. (1992). Temperature effects on early season cotton growth and development. *Agronomy Journal* 84(2): 229-237.
- Reddy, V. R., A. Trent, et al. (1992). Mepiquat chloride and irrigation versus cotton growth and development. *Agronomy Journal* 84(6): 930-933. {a} Dep. Plant, Soil Entomol. Sci., Coll. Agric., Univ. Idaho, Moscow, Idaho 83843
- Reinbothe, S., A. Machmudova, et al. (1992). Jasmonate-induced proteins in cotton: Immunological relationship to the respective barley proteins and homology of transcripts to late embryogenesis abundant (Lea) mRNAs. *Journal Of Plant Growth Regulation* 11(1): 7-14.
- Rigi, D. G., S. V. Raikhelkar, et al. (1992). Seed-cotton yield and monetary economics of upland cotton (*Gossypium hirsutum*) as influenced by weed management. *Indian Journal of Agronomy* 37(4): 778-780. Marathwada Agric. Univ., Parbhani, Maharashtra 431 402

- Rikin, A. (1992). Circadian rhythm of heat resistance in cotton seedlings: Synthesis of heat-shock proteins. *European Journal of Cell Biology* 59(1): 160-165. Dep. Botany, Okla. State Univ., Stillwater, Okla. 74078-0293, USA
- Rikin, A. (1992). Temporal organization of chilling resistance in cotton seedlings: Effects of low temperature and relative humidity. *Planta* 187(4): 517-522.
- Roberts, E. M., N. R. Rao, et al. (1992). Effects of cycling temperatures on fiber metabolism in cultured cotton ovules. *Plant Physiology Rockville* 100(2): 979-986. {a} Dep. Biological Sciences, Texas Tech Univ., Lubbock, Tex. 79409
- Roberts, R. K. and B. C. English (1992). Cotton acreage response and fertilizer use. *Journal Of Production Agriculture* 5(1): 158-162.
- Rodge, R. P., V. D. Sondge, et al. (1992). Cropping-system planning for semi-arid Marathwada region under limited availability of irrigation water. *Indian Journal Of Agricultural Sciences* 62(7): 436-441.
- Ross, W. (1992). Confocal pH topography in plant cells: Acidic layers in the peripheral cytoplasm and the apoplast. *Botanica Acta* 105(4): 253-259. Martin-Luther-Univ. Halle, FB Pharmazie, Abt. Pharmazeutische Biotechnologie Zellphysiologie, Weinbergweg 16a, D-4050 Halle, PSF 8, Germany
- Ryser, U. (1992). Ultrastructure of the epidermis of developing cotton (*Gossypium*) seeds: Suberin, pits, plasmodesmata, and their implication for assimilate transport into cotton fibers. *American Journal Of Botany* 79(1): 14-22.
- Sanam, y. M. F. and D. A. Musaev (1992). Production and cytogenetic study of the homozygous translocation stocks of cotton. *Genetika* 28(11): 75-82. V.I. Lenin Tashk. State Univ., Tashkent, Uzbekistan
- Sanyasi, I. P. S. (1992). Genetic analysis of yield and number of bolls in upland cotton *Gossypium hirsutum* L.). *Indian Journal Of Agricultural Research* 25(3): 167-172.
- Selley, R. (1992). Relative efficiency of production alternatives: Irrigation in cotton. *Journal Of Production Agriculture* 5(3): 317-323.
- Sethi, H. N., G. M. Bharad, et al. (1992). Biomass production of cotton (*Gossypium hirsutum*) varieties as influenced by intercrops. *Indian Journal of Agronomy* 37(3): 451-455. Dep. Agronomy, Punjabrao Krishi Vidyapeeth, Akola, Maharashtra 444 104
- Shabashvili, E. Z. and V. I. Chikov (1992). Structural and functional response of cotton leaf chloroplasts to changes of sink-source interactions in plants. *Fiziologiya Rastenii Moscow* 39(3): 480-487. Kazan Inst. Biol., Kazan Sci. Cent., Acad. Sci. Russ., Kazan, Russia
- Shahina, F. and M. A. Maqbool (1992). Description of *Deladenus pakistanensis*, new species (Nematoda: Tylenchida) from Pakistan. *Pakistan Journal Of Nematology* 10(1): 1-6.
- Shanmugham, K. (1992). Seed soaking and foliar application of growth-regulants and anti-transpirant chemicals for increasing drought resistance in rainfed upland cotton (*Gossypium hirsutum*). *Indian Journal of Agricultural Sciences* 62(11): 744-750. Regional Station, Central Institute Cotton Research, Coimbatore, Tamil Nadu 641 003, India
- Sharma, J. K., K. N. Namdeo, et al. (1992). Effect of sowing dates, plant spacings and varieties on yield of *hirsutum* cotton. *Crop Research Hisar* 5(3): 440-444. All India Coordinated Cotton Improvement Project, B. M. Coll. Agric., Khandwa-450 001, India
- Sharma, J. P. and R. A. Agarwal (1992). Effect of insecticidal application on the fibre quality, seed-cotton and carryover of *Pectinophora gossypiella* (Saund.). *Journal of Insect Science* 5(2): 153-156. {a} HRS, Seobagh, Kullu 175 101, India

- Shu, T. T. and E. Y. Chang (1992). Studies of fertilization rates and application of sucker control agents on yield and quality of tobacco variety TT8. Bulletin of Taiwan Tobacco Research Institute Taiwan Tobacco and Wine Monopoly Bureau(37): 1-7. Dep. Agron., Tobacco Res. Inst., TTWMB, Taichung, Taiwan
- Siddiqui, M. A. and R. A. Patil (1992). Combining ability for yield and yield contributing traits and earliness in upland cotton under irrigated conditions. Journal of Maharashtra Agricultural Universities 17(3): 383-386. Agric. Res. Stn., Badnapur, India
- Silvertooth, J. C., J. E. Watson, et al. (1992). Bromide and nitrate movement in an irrigated cotton production system. Soil Science Society Of America Journal 56(2): 548-561.
- Simwat, G. S., A. K. Dhawan, et al. (1992). Criteria for initiating insecticide spray to control bollworms (*Pectinophora gossypiella*, *Earias vittella* and *Earias insulana*) of cotton (*Gossypium* spp.). Indian Journal Of Agricultural Sciences 62(1): 85-87.
- Singh, D., M. S. Brar, et al. (1992). Critical concentrations of potassium in cotton (*Gossypium hirsutum*). Journal Of Agricultural Science 118(1): 71-76.
- Singh, D. and H. Singh (1992). Calibration of a tractor-mounted sprayer for optimum control of bollworms (*Pectinophora gossypiella*, *Earias insulana* and *Earias vitella*) in upland cotton (*Gossypium hirsutum*). Indian Journal Of Agricultural Sciences 62(9): 637-642.
- Singh, K., N. Singh, et al. (1992). Nitrogen requirement of sorghum fodder grown in different cropping systems. International Journal of Tropical Agriculture 10(3): 165-167. Dep. Agron., CCS Haryana Agric. Univ., Hisar-125 004, India
- Singh, M., V. P. Singh, et al. (1992). Breeding suitable plant type for high productivity in upland cotton (*Gossypium hirsutum*). Indian Journal Of Agricultural Sciences 62(5): 319-324.
- Singh, M., V. P. Singh, et al. (1992). Selection for greater fibre strength in medium and superior staple upland cotton (*Gossypium hirsutum*). Indian Journal Of Agricultural Sciences 62(5): 325-328.
- Singh, S. B. (1992). Variation in leaf anatomy of upland cotton (*Gossypium hirsutum*), tree cotton (*Gossypium arboreum*) and asiatic cotton (*Gossypium herbaceum*). Indian Journal of Agricultural Sciences 62(12): 819-821. Div. Crop Improvement, Central Inst. Cotton Res., Nagpur, Maharashtra 440 001, India
- Smith, C. W., J. C. McCarty, Jr., et al. (1992). Condensed tannins in cotton and bollworm-budworm (*Lepidoptera: Noctuidae*) resistance. Journal of Economic Entomology 85(6): 2211-2217. {a} Dep. Soil Crop Sci., Texas A and M Univ., College Station, Texas 77843
- Snipes, C. E., D. L. Colvin, et al. (1992). Cotton (*Gossypium hirsutum*) yield response to cultivation timing and frequency. Weed Technology 6(1): 31-35.
- Snipes, C. E. and T. C. Mueller (1992). Cotton (*Gossypium hirsutum*) yield response to mechanical and chemical weed control systems. Weed Science 40(2): 249-254.
- Snipes, C. E., J. E. Street, et al. (1992). Cotton (*Gossypium hirsutum*) injury from simulated quinclorac drift. Weed Science 40(1): 106-109.
- Srinivasaperumal, S., P. Samuthiravelu, et al. (1992). Life tables and energetics of *Earias vittella* (Fab.) (*Noctuidae: Lepidoptera*) reared on three different hosts. Insect Science and its Application 13(6): 749-754. Dep. Anim. Physiol., Sch. Biol. Sci., Madurai Kamaraj Univ., Madurai 625 021, India
- Steiner, J. J. and T. A. Jacobsen (1992). Time of planting and diurnal soil temperature effects on cotton seedling field emergence and rate of development. Crop Science 32(1): 238-244.

Stevens, W. E., J. R. Johnson, et al. (1992). Tillage and winter cover management effects of fruiting and yield of cotton. *Journal of Production Agriculture* 5(4): 570-575. {a} GOSSYM -COMAX Information Unit, 102 Technol. Center, Starkville, Miss. 39759

Stipanovic, R. D., M. E. Mace, et al. (1992). The role of free radicals in the decomposition of the phytoalexin desoxyhemigossypol. *Journal of the Chemical Society Perkin Transactions I*(23): 3189-3192. US Dep. Agric., Agric. Res. Serv., South. Crops Food Anim. Protection Res. Lab., Route 5, Box 805, College Station, TX 77845, USA

Stoilova, A. (1992). Type of hybrid development in *Gossypium hirsutum* L. x *Gossypium barbadense* L. backcrosses with both parental species: II. Characteristics of BC-1 in backcrosses of F-2. *Genetika I Seleksiya* 25(2): 153-159.

Sukhadia, N. M. and M. V. Dhoble (1992). Productivity and water-use efficiency of rainy-season crops under different dates of sowing. *Indian Journal of Agronomy* 37(4): 669-675. {a} Dep. Agron., G.A.U. Junagadh 360 001

Sukhadia, N. M., M. V. Dhoble, et al. (1992). Productivity and water-use efficiency of rainy-season crops under different dates of sowing. *Indian Journal of Agronomy* 37(3): 461-465. {a} Dep. Agronomy, Gujarat Agric. Univ., Junagadh

Tarczynski, M. C., D. N. Byrne, et al. (1992). High performance liquid chromatography analysis of carbohydrates of cotton-phloem sap and of honeydew produced by *Bemisia tabaci* feeding on cotton. *Plant Physiology* 98(2): 753-756.

Tariq, M., M. A. Khan, et al. (1992). Correlation and path coefficient analysis in Upland cotton. *Sarhad Journal Of Agriculture* 8(3): 335-341.

Terry, N., C. Carlson, et al. (1992). Rates of selenium volatilization among crop species. *Journal Of Environmental Quality* 21(3): 341-344.

Thaker, V. S., S. Saroop, et al. (1992). Physiological and biochemical changes associated with cotton fiber development: V. Acid invertase and sugars. *Acta Physiologiae Plantarum* 14(1): 11-18.

Thayer, S. S. and O. Bjorkman (1992). Carotenoid distribution and deepoxidation in thylakoid pigment-protein complexes from cotton leaves and bundle-sheath cells of maize. *Photosynthesis Research* 33(3): 213-225. Carnegie Institution of Washington, Dep. Plant Biol., 290 Panama St., Stanford, Calif. 94305-1297, USA

Tingle, F. C. and E. R. Mitchell (1992). Attraction of *Heliothis virescens* (F.) (Lepidoptera: Noctuidae) to volatiles from extracts of cotton flowers. *Journal Of Chemical Ecology* 18(6): 907-914.

Tiwari, V. N., K. C. Mandloi, et al. (1992). Breeding behaviour effects for yield and its components in intra-specific crosses of cotton (*Gossypium hirsutum* L.). *Indian Journal of Agricultural Research* 26(4): 173-178. {a} Dep. Plant Breeding Genetics, JNKVV, Jabalpur, India

Tracy, P. W. and W. P. Sappenfield (1992). Integrated management influence on short-season cotton yield in the Northern Mississippi Delta. *Journal of Production Agriculture* 5(4): 551-555. {a} Dep. Agronomy, Univ. Missouri Delta Res. Center, Portageville, Mo. 63873

Triplett, B. A. and J. E. Mellon (1992). Purification and characterization of anionic peroxidases from cotton (*Gossypium hirsutum* L.). *Plant Science* 81(2): 147-154.

Ungar, E. D., E. Kletter, et al. (1992). Conservative response and stress-damage interactions in cotton reproductive development. *Agronomy Journal* 84(3): 382-386.

Vencill, W. K., L. J. Giraud, et al. (1992). Response of cotton (*Gossypium hirsutum*) to coastal bermudagrass (*Cynodon dactylon*) density in a no-tillage system. *Weed Science* 40(3): 455-459.

Vepraskas, M. J. and D. S. Guthrie (1992). Identifying soils where subsoiling can increase yields of tobacco, corn, and cotton. *Journal of Production Agriculture* 5(4): 482-488. {a} Soil Science Dep., Crop Sci. Dep., North Carolina State Univ., Raleigh, N.C. 27695-7619

Vesmanova, O. Y. and R. G. Dani (1992). Influence of carbohydrate composition of media on clonal micropropagation ability of cotton. *Advances in Plant Sciences* 5(1): 97-99. Lab. Cytoembryol. and Cell Eng., Inst. Exp. Biol. and Plants, Sci. Prod. Establishment "Biologist", Acad. Sci., Uzbek S.S.R., 28 Faizulta Khojaev St., Tashkent 700 143, Russia

Voluevich, E. A. and A. A. Buloichik (1992). Nuclear-cytoplasmic interactions in wheat resistance to fungi pathogens: VI. Nuclear genome influence of alloplasmic lines on quantitative resistance of seedling to clones. *Genetika* 28(11): 68-74. Inst. Genet. Cytol., Acad. Sci. B., Minsk, Beloruss

Wadibhasme, S. S., N. K. Raut, et al. (1992). Correlation of chemical composition of cotton leaves to their reaction to *Alternaria* blight. *Journal of Maharashtra Agricultural Universities* 17(3): 407-409. Punjabrao Krishi Vidyapeeth, Akola-444 104, India

Wallace, T. P. and Z. K. M. El (1992). Reaction of three cotton (*Gossypium hirsutum*) cultivars to single and mixed isolates of *Xanthomonas campestris* pv. *malvacearum*. *Plant Pathology Oxford* 41(5): 569-572. {a} Dep. of Agronomy, P.O. Box 5248, Mississippi State Univ., Mississippi State, Miss. 39762, USA

Wankhade, S. T., S. G. Kharche, et al. (1992). Effect of spacing and varieties on growth, yield and quality of Asiatic cotton (*Gossypium arboreum*) under rainfed conditions. *Indian Journal of Agronomy* 37(3): 523-526. Punjabrao Krishi Vidyapeeth, Akola, Maharashtra 444 104

Wendel, J. F., C. L. Brubaker, et al. (1992). Genetic diversity in *Gossypium hirsutum* and the origin of upland cotton. *American Journal of Botany* 79(11): 1291-1310. {a} Dep. Botany, Iowa State Univ., Ames, Iowa 50011

Wetzel, C. L. R. and W. A. Jensen (1992). Studies of pollen maturation in cotton: The storage reserve accumulation phase. *Sexual Plant Reproduction* 5(2): 117-127.

Williams, W. P. and G. L. Windham (1992). Reaction of a diallel cross of maize to *Meloidogyne incognita* under field conditions. *Field Crops Research* 30(1-2): 167-171.

Willis, G. H., L. L. McDowell, et al. (1992). Washoff of ultra -low-volume -oil-applied insecticides from cotton plants as a function of time between application and rainfall. *Journal Of Environmental Quality* 21(3): 373-377.

Wilson, F. D. and C. R. J. Smith (1992). Pink bollworm (Lepidoptera: Gelechiidae) penetration into and survival in bolls of resistant and susceptible cotton lines. *Journal Of Agricultural Entomology* 9(3): 165-173.

Wilson, F. D., J. L. Szaro, et al. (1992). Behavior of pink bollworm larvae (Lepidoptera: Gelechiidae) on bolls of normal-leaf and okra-leaf cotton isolines under laboratory conditions. *Journal Of Agricultural Entomology* 9(1): 55-64.

Wise, R. R., L. A. Ortiz, et al. (1992). Spatial distribution of photosynthesis during drought in field-grown and acclimated and nonacclimated growth chamber-grown cotton. *Plant Physiology Rockville* 100(1): 26-32. {a} Dep. Agronomy, Univ. Wisconsin, 1575 Linden Drive, Madison, Wis. 53706

Witz, J. A., J. D. J. Lopez, et al. (1992). Influence of sexpheromone trap placement relative to field edge on catch of bollworm males in cotton. *Southwestern Entomologist* 17(1): 1-6.

Wixson, M. B. and D. R. Shaw (1992). Effects of soil-applied AC 263222 on crops rotated with soybean (*Glycine max*). *Weed Technology* 6(2): 276-279.

Wullschleger, S. D. and D. M. Oosterhuis (1992). Canopy leaf area development and age-class dynamics in cotton. *Crop Science* 32(2): 451-456.

- Xiao, Z. A., D. X. Shen, et al. (1992). Relationship between the physiological difference of kiwifruit calli and growth, differentiation of their protoplasts. *Acta Phytophysiologica Sinica* 18(4): 369-375. Dep. Horticulture, Zhejiang Agric. Univ., Hangzhou 310029, China
- Xu, X. and H. M. Taylor (1992). Increase in drought resistance of cotton seedlings treated with mepiquat chloride. *Agronomy Journal* 84(4): 569-574.
- Yigit, A. (1992). Method for culturing *Serangium parcesetosum* Sicard (Coleoptera: Coccinellidae) on *Bemisia tabaci* Genn. (Homoptera: Aleyrodidae). *Zeitschrift fuer Pflanzenkrankheiten und Pflanzenschutz* 99(5): 525-527. Inst. Plant Protection, Univ. Cukurova, Adana, Turkey
- York, A. C. and D. L. Jordan (1992). Cotton (*Gossypium hirsutum*) response to clomazone and insecticide combinations. *Weed Technology* 6(4): 796-800. {a} Dep. Crop Sci., NC State Univ., Raleigh, NC 27695-7620, USA
- Yusupov, T. Y., R. M. Nauruzbaeva, et al. (1992). Detection and characterization of mitochondrial plasmid-like DNA in cotton. *Biopolimery i Kletka* 8(3): 19-23. {a} Inst. Exp. Plant Biol., Acad. Sci. Ukr., Kiev, Ukraine
- Zeng, Y. S., L. P. Yang, et al. (1992). Race-specific resistance protein to fusarium wilt in cotton. *Chinese Journal of Botany* 4(2): 137-142. {a} Shanghai Inst. Biochem., Acad. Sinica, Shanghai 200031, China
- Zeringue, H. J. J. (1992). Effects of C-6-C-10 alkenals and alkanals on eliciting a defence response in the developing cotton boll. *Phytochemistry* 31(7): 2305-2308.
- Zhou, S. Q., D. Q. Qian, et al. (1992). Inheritance of characters and chromosome number in aneusomatic progeny of upland cotton (*Gossypium hirsutum*) of parthenogenetic origin. *Genome* 35(1): 35-38.
- Zhu, B. and D. M. Oosterhuis (1992). Nitrogen distribution within a sympodial branch of cotton. *Journal Of Plant Nutrition* 15(1): 1-14.
- Afek, U., J. A. Menge, et al. (1991). Interaction among mycorrhizae, soil solarization, metalaxyl, and plants in the field. *Plant Disease* 75(7): 665-671.
- Allen, S. J. and K. L. D. West (1991). Predominance of race 18 of *Xanthomonas campestris* pathovar *malvacearum* on cotton in Australia. *Plant Disease* 75(1): 43-44.
- Altman, D. W., D. M. Stelly, et al. (1991). Quantitative trait variation in phenotypically normal regenerants of cotton. *In Vitro Cellular & Developmental Biology Plant* 27p 3: 132-138.
- Ansari, A. H., S. M. Qayyum, et al. (1991). Influence of seedling dates on the yield its components and their interrelation in cotton *Gossypium hirsutum* L. genotypes. *Sarhad Journal Of Agriculture* 7(2): 11-19.
- Aujla, M. S., C. J. Singh, et al. (1991). Evaluation of methods for irrigation of cotton (*Gossypium hirsutum*) in a canal-irrigated area of southwest Punjab, India. *Arid Soil Research And Rehabilitation* 5(3): 225-234.
- Ayub, M., S. Rahman, et al. (1991). Combining ability analysis of some quality characters in various crosses of cotton (*Gossypium hirsutum* L.). *Sarhad Journal Of Agriculture* 7(6): 707-716.
- Barnes, C. J. and T. L. Lavy (1991). Injury and yield response of selected crops to imazaquin and norflurazon residues. *Weed Technology* 5(3): 598-606.
- Basford, K. E., P. M. Kroonenberg, et al. (1991). Three-way methods for multiattribute genotype X environment data: An illustrated partial survey. *Field Crops Research* 27(1-2): 131-158.
- Bashan, Y., H. Levanony, et al. (1991). Association between *Alternaria macrospora* and *Alternaria alternata*, causal agents of cotton leaf blight. *Canadian Journal Of Botany* 69(12): 2603-2607.

- Bashan, Y., H. Levanony, et al. (1991). Wind dispersal of *Alternaria alternata*, a cause of leaf blight of cotton. *Journal Of Phytopathology* 133(3): 225-238.
- Bates, S. L., J. K. Walker, et al. (1991). Studies of boll weevil resistance in two converted, day-neutral cotton race stocks. *Southwestern Entomologist* 16(4): 317-330.
- Bauer, P. J., S. H. Roach, et al. (1991). Cotton genotype response to green-manured annual legumes. *Journal Of Production Agriculture* 4(4): 626-628.
- Bhatade, S. S., S. R. Rajewar, et al. (1991). Phenotypic stability in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 61(11): 829-830.
- Bilger, W. and O. Bjoerkman (1991). Temperature dependence of violaxanthin deepoxidation and non-photochemical fluorescence quenching in intact leaves of *Gossypium hirsutum* L. and *Malva parviflora* L. *Planta* 184(2): 226-234.
- Bordovsky, J. P., W. M. Lyle, et al. (1991). Cottonseed preparation for hydraulic planter. *Agronomy Journal* 83(2): 275-278.
- Boyette, C. D. (1991). Host range and virulence of *Colletotrichum truncatum*, a potential mycoherbicide for hemp sesbania (*Sesbania exaltata*). *Plant Disease* 75(1): 62-64.
- Bozcuk, S. (1991). Effect of salinity on germination and determination of salt tolerance in some culture plants. *Doga Biyoloji Serisi* 15(2): 144-151.
- Brar, G. S., B. L. McMichael, et al. (1991). Hydraulic conductivity of cotton roots as influenced by plant age and rooting medium. *Agronomy Journal* 83(1): 264-266.
- Brown, J. S. (1991). Principal component and cluster analyses of cotton cultivar variability across the USA cotton belt. *Crop Science* 31(4): 915-922.
- Brugnoli, E. and M. Lauteri (1991). Effects of salinity on stomatal conductance, photosynthetic capacity, and carbon isotope discrimination of salt-tolerant (*Gossypium hirsutum* L.) and salt-sensitive (*Phaseolus vulgaris* L.) C-3 non-halophytes. *Plant Physiology* 95(2): 628-635.
- Byrd, J. D. J. and H. D. Coble (1991). Interference of common cocklebur (*Xanthium strumarium*) and cotton (*Gossypium hirsutum*). *Weed Technology* 5(2): 270-278.
- Byrd, J. D. J. and H. D. Coble (1991). Interference of selected weeds in cotton (*Gossypium hirsutum*). *Weed Technology* 5(2): 263-269.
- Cass, Q. B., E. Tiritan, et al. (1991). Gossypol enantiomer ratios in cotton seeds. *Phytochemistry* 30(8): 2655-2658.
- Chapman, K. D. and R. N. Trelease (1991). Intracellular localization of phosphatidylcholine and phosphatidylethanolamine synthesis in cotyledons of cotton seedlings. *Plant Physiology* 95(1): 69-76.
- Chen, X. and S. Zhu (1991). Studies on the inheritance of flaring-bract in upland cotton. *Acta Agronomica Sinica* 17(5): 381-386.
- Chosack, E., B. Rubinstein, et al. (1991). Effect of iron deficiency on electron transport process of plasmalemma-enriched vesicles isolated from cotton roots. *Plant Science* 77(2): 163-172.
- Chu, C. C., T. J. Henneberry, et al. (1991). Pink bollworm in different cottons and effect of plant growth regulators under short-season conditions. *Southwestern Entomologist* 16(2): 132-143.
- Chu, Y. N., C. G. Coble, et al. (1991). Hypocotyl elongation and swelling of cotton as affected by soil temperature, moisture and physical impedance. *Crop Science* 31(2): 410-415.

- Chu, Y. N., C. G. Coble, et al. (1991). Cotton emergence force as affected by soil temperature, moisture, and compression. *Crop Science* 31(2): 405-409.
- Cihacek, L. J. and T. A. Kerby (1991). Effects of residual soil nitrogen and urea on yield and petiole nitrate of cotton. *Journal Of Production Agriculture* 4(2): 193-197.
- Colyer, P. D., S. Micinski, et al. (1991). Effect of planting date on the efficacy of an in-furrow pesticide and the development of cotton seedling disease. *Plant Disease* 75(7): 739-742.
- Colyer, P. D., S. Micinski, et al. (1991). Effect of thrips infestation on the development of cotton seedling diseases. *Plant Disease* 75(4): 380-382.
- Constable, G. A. (1991). Mapping the production and survival of fruit on field-grown cotton. *Agronomy Journal* 83(2): 374-378.
- Cook, C. G. and Z. K. M. El (1991). Evaluation of twelve upland cotton genotypes for resistance to *Phymatotrichum omnivorum*. *Plant Disease* 75(1): 56-58.
- Corbin, B. R. J. and R. E. Frans (1991). Protecting cotton (*Gossypium hirsutum*) from fluometuron injury with seed protectants. *Weed Science* 39(3): 408-411.
- Costa, H. S. and J. K. Brown (1991). Variation in biological characteristics and esterase patterns among populations of *Bemisia tabaci*, and the association of one population with silverleaf symptom induction. *Entomologia Experimentalis Et Applicata* 61(3): 211-220.
- Dani, R. G. (1991). Analysis of combining ability for seed-oil content in cotton (*Gossypium hirsutum* L.). *Acta Agronomica Hungarica* 40(1-2): 123-128.
- Davis, G. D., E. J. Eisenbraun, et al. (1991). Tritium transfer during biosynthesis of cadalene stress compounds in cotton. *Phytochemistry* 30(1): 197-200.
- Delmer, D. P., M. Solomon, et al. (1991). Direct photolabeling with phosphorus-32 UDP-glucose for identification of a subunit of cotton fiber callose synthase. *Plant Physiology* 95(2): 556-563.
- Dhaliwal, Z. S., J. Sigh, et al. (1991). Limitations in the use of thermal summation for describing the activity of pink bollworm, *Pectinophora gossypiella* (Saunders), in relation to phenology of Upland cotton in Punjab (India). *Tropical Agriculture* 68(3): 268-270.
- Duke, S. O., R. N. Paul, et al. (1991). Clomazone causes accumulation of sesquiterpenoids in cotton (*Gossypium hirsutum* L.). *Weed Science* 39(3): 339-346.
- Dzhuraev, Y. F. and A. K. E. Ergashev (1991). Inheritance and localization of some genes controlling cotton fertility. *Tsitologiya I Genetika* 25(3): 30-34.
- Endrizzi, J. E. and D. T. Ray (1991). Linkage analysis of open bud (ob-2) and yellow petal (Y-1) in cotton. *Genome* 34(3): 461-463.
- Endrizzi, J. E. and D. T. Ray (1991). Monosomic and monotelodisomic analysis of 34 mutant loci in cotton. *Journal Of Heredity* 82(1): 53-57.
- Ephrath, J. E. and J. D. Hesketh (1991). The thermal-photoperiod requirement for floral bud growth. *Biotronics* 20: 1-7.
- Esau, K. and R. H. Gill (1991). Distribution of vacuoles and some other organelles in dividing cells. *Botanical Gazette* 152(4): 397-407.
- Finer, J. J., R. W. Saxton, et al. (1991). Growth and disinfestation of 6 different bacteria in embryogenic suspension cultures of cotton. *Plant Cell Reports* 10(8): 380-383.

- Flint, H. M., F. D. Wilson, et al. (1991). Suppression of pink bollworm and effect on beneficial insects of a nectariless okra-leaf cotton germplasm line. *Bulletin Of Entomological Research* 81(4): 379-384.
- Frans, R. E., M. R. McClelland, et al. (1991). Crop and herbicide rotations for johnsongrass (*Sorghum halepense*) control. *Weed Science* 39(4): 660-666.
- Freire, E. C. and J. D. A. N. Moreira (1991). Genetic relationships among "Moco" and other species and races of cotton (*Gossypium* spp.). *Revista Brasileira De Genetica* 14(2): 393-412.
- Georgen, P. G., C. J. Davis, et al. (1991). Root growth and water extraction patterns from a calcic horizon. *Soil Science Society Of America Journal* 55(1): 210-215.
- Gould, J., S. Banister, et al. (1991). Regeneration of *Gossypium hirsutum* and *Gossypium barbadense* from shoot apex tissues for transformation. *Plant Cell Reports* 10(1): 12-16.
- Gridi, P. I. L., E. Cia, et al. (1991). Improvement of Sao Paulo cotton varieties: The origin of the IAC19. *Bragantia* 50(2): 181-194. {a} Secao Algodao, Inst. Agronomico, Caixa Postal 28, 13001 Campinas, SP
- Grossman, K. (1991). Induction of leaf abscission in cotton is a common effect of urea- and adenine-type cytokinins. *Plant Physiology* 95(1): 234-237.
- Gururajan, K. N., S. Henry, et al. (1991). Comparative evaluation of two methods for selecting stable hybrids in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 61(7): 480-483.
- Guthrie, D. S. (1991). Cotton response to starter fertilizer placement and planting dates. *Agronomy Journal* 83(5): 836-839.
- Gutierrez, A. P., S. W. J. Dos, et al. (1991). Modelling the interaction of cotton and the cotton boll weevil: I. A comparison of growth and development of cotton varieties. *Journal Of Applied Ecology* 28(2): 371-397.
- Haigler, C. H., N. R. Rao, et al. (1991). Cultured ovules as models for cotton fiber development under low temperatures. *Plant Physiology* 95(1): 88-96.
- Halloin, J. M., L. S. Lee, et al. (1991). Pre-ripening damage to cottonseed by *Aspergillus flavus* is not influenced by seed coat permeability. *Journal Of The American Oil Chemists' Society* 68(7): 522-523.
- Ham, J. M. and J. L. Heilman (1991). Aerodynamic and surface resistances affecting energy transport in a sparse crop. *Agricultural And Forest Meteorology* 53(4): 267-284.
- Ham, J. M., J. L. Heilman, et al. (1991). Soil and canopy energy balances of a row crop at partial cover. *Agronomy Journal* 83(4): 744-753.
- Hatfield, J. L. and J. J. Burke (1991). Energy exchange and leaf temperature behavior of three plant species. *Environmental And Experimental Botany* 31(3): 295-302.
- He, J., Z. Liang, et al. (1991). Interspecific incompatibility in crosses between the cultivated and the wild species in *Gossypium*. *Acta Genetica Sinica* 18(2): 140-148.
- Hedin, P. A. and J. C. J. McCarty (1991). Effects of kinetin formulations on allelochemicals and agronomic traits of cotton. *Journal Of Agricultural And Food Chemistry* 39(3): 549-553.
- Hendrix, D. L. and R. I. Grange (1991). Carbon partitioning and export from mature cotton leaves. *Plant Physiology* 95(1): 228-233.
- Henning, S. A. and C. H. Cardona (1991). Limiting supplemental irrigation in the Delta States (USA): Impact on net return and cropping patterns. *Journal Of Production Agriculture* 4(1): 98-105.

Hickman, M. V., J. H. Everitt, et al. (1991). Aerial photography and videography for detecting and mapping dicamba injury patterns. *Weed Technology* 5(4): 700-706.

Hillocks, R. J. (1991). *Alternaria* leaf spot of cotton with special reference to Zimbabwe. *Tropical Pest Management* 37(2): 124-128.

Hillocks, R. J. (1991). Screening for resistance to verticillium wilt in Zimbabwe. *Tropical Agriculture* 68(2): 144-148.

Holt, J. S. and D. R. Orcutt (1991). Functional relationships of growth and competitiveness in perennial weeds and cotton (*Gossypium hirsutum*). *Weed Science* 39(4): 575-584.

Howell, C. R. (1991). Biological control of pythium damping-off of cotton with seed-coating preparations of *Gliocladium virens*. *Phytopathology* 81(7): 738-741.

Hutchison, W. D., C. A. Beasley, et al. (1991). Timing insecticide applications for pink bollworm (Lepidoptera: Gelechiidae) management: Comparison of egg and larva treatment thresholds. *Journal Of Economic Entomology* 84(2): 470-475.

Jackson, S. H. (1991). Relationships between normalized leaf water potential and crop water stress index values for acala cotton. *Agricultural Water Management* 20(2): 109-118.

Kamara, L., R. Zartman, et al. (1991). Cotton-root distribution as a function of trickle irrigation emitter depth. *Irrigation Science* 12(3): 141-144.

Keeley, P. E. and R. J. Thullen (1991). Biology and control of black nightshade (*Solanum nigrum*) in cotton (*Gossypium hirsutum*). *Weed Technology* 5(4): 713-722.

Keeley, P. E. and R. J. Thullen (1991). Growth and interaction of bermudagrass (*Cynodon dactylon*) with cotton (*Gossypium hirsutum*). *Weed Science* 39(4): 570-574.

Keeley, P. E. and R. J. Thullen (1991). Growth and interaction of barnyardgrass (*Echinochloa crus galli*) with cotton (*Gossypium hirsutum*). *Weed Science* 39(3): 369-375.

Keeling, J. W., K. T. Siders, et al. (1991). Palmer amaranth (*Amaranthus palmeri*) control in a conservation tillage system for cotton (*Gossypium hirsutum*). *Weed Technology* 5(1): 137-141.

Kennedy, C. W., W. C. J. Smith, et al. (1991). Chemical efficacy of early square removal and subsequent productivity of superokra-leaf cotton. *Crop Science* 31(3): 791-796.

Kharche, S. C. and R. M. Deshpande (1991). Response of dwarf genotypes of American cotton (*Gossypium hirsutum*) to spacing. *Indian Journal Of Agronomy* 36(3): 392-394.

Kirkpatrick, T. L., D. M. Oosterhuis, et al. (1991). Interaction of *Meloidogyne incognita* and water stress in two cotton cultivars. *Journal Of Nematology* 23(4): 462-467.

Koeniger, M. and K. Winter (1991). Carotenoid composition and photon-use efficiency of photosynthesis in *Gossypium hirsutum* L. grown under conditions of slightly suboptimum leaf temperatures and high levels of irradiance. *Oecologia* 87(3): 349-356.

Koganova, T. V., D. K. Shrivastava, et al. (1991). Morphogenic potentials of transformed callus tissue of cotton plants. *Biotekhnologiya*(3): 47-50.

Lakshmanan, P. and P. Vidhyasekaran (1991). *Corticium rolfsii*, a new potential pathogen causing boll rot of *Gossypium hirsutum*. *Zeitschrift Fuer Pflanzenkrankheiten Und Pflanzenschutz* 98(1): 80-83.

Lehle, F. R., A. M. Zegeer, et al. (1991). Ethanolic fermentation in hypoxic cotton seed. *Crop Science* 31(3): 746-750.

Levy, G. J., H. M. Ben, et al. (1991). The effect of polyacrylamide on runoff, erosion, and cotton yield from fields irrigated with moving sprinkler systems. *Irrigation Science* 12(2): 55-60.

Lewis, J. A. and G. C. Papavizas (1991). Biocontrol of cotton damping-off caused by *Rhizoctonia solani* in the field with formulations of *Trichoderma* spp. and *Gliocladium virens*. *Crop Protection* 10(5): 396-402.

Liu, C. M. and D. Y. Yao (1991). Studies on somatic embryogenesis and histological observation in *Gossypium hirsutum* L. cultivar Coker 312. *Acta Botanica Sinica* 33(5): 378-384.

Liu, C. Z., F. X. Cheng, et al. (1991). Artificial germination of *Gossypium* pollen and its utilization in interspecific fertilization in vitro. *Acta Botanica Sinica* 33(7): 523-528.

Lueschen, W. E., S. D. Evans, et al. (1991). Soybean production as affected by tillage in a corn and soybean management system: II. Seed treatment response. *Journal Of Production Agriculture* 4(4): 580-585.

Lukin, Y. L. and E. A. Simonova (1991). Separation of carotenoid pigments by high-performance liquid chromatography in isocratic regime. *Fiziologiya Rastanii* 38(4): 817-822.

Mahadevan, A., B. Parthipan, et al. (1991). Morphological and histological changes in *Gossypium hirsutum* L. and *Ocimum sanctum* L. exposed to methylisocyanate. *Indian Journal Of Experimental Biology* 29(6): 554-557.

Malik, M. N., U. D. Shabab, et al. (1991). Accelerated cotton boll dehiscence with thidiazuron. *Tropical Agriculture* 68(2): 149-150.

Martinez, V. and A. Lauchli (1991). Phosphorus translocation in salt-stressed cotton. *Physiologia Plantarum* 83(4): 627-632.

Mateos, L., J. Berengena, et al. (1991). A comparison between drip and furrow irrigation in cotton at two levels of water supply. *Agricultural Water Management* 19(4): 313-324.

Mateos, L., J. Berengena, et al. (1991). Effects of drip and furrow irrigation on the phenology and yield of cotton. *Investigacion Agraria Produccion Y Proteccion Vegetales* 6(1): 69-80.

McAuslane, H. J., S. B. Vinson, et al. (1991). Stimuli influencing host microhabitat location in the parasitoid *Campoplex sonorensis*. *Entomologia Experimentalis Et Applicata* 58(3): 267-278.

McCray, J. M., M. E. Sumner, et al. (1991). Soil calcium aluminium acidity and penetration resistance with subsoiling, lime and gypsum treatments. *Soil Use And Management* 7(4): 193-199.

McDowell, L. L., G. H. Willis, et al. (1991). Interception and persistence of parathion and fenvalerate on cotton plants as a function of application method. *Pesticide Science* 33(3): 271-280.

Meek, D. W., R. B. Hutmacher, et al. (1991). Heteroscedasticity in whole plant growth curves developed from nonreplicated data. *Agronomy Journal* 83(2): 417-424.

Mellon, J. E. (1991). Purification and characterization of isoperoxidases elicited by *Aspergillus flavus* in cotton ovule cultures. *Plant Physiology* 95(1): 14-20.

Michaelson, M. J., H. J. Price, et al. (1991). Comparison of plant DNA contents determined by Feulgen microspectrophotometry and laser flow cytometry. *American Journal Of Botany* 78(2): 183-188.

Micinski, S., R. G. Luttrell, et al. (1991). Relationship of adult vial test data on pyrethroid resistance to field control of tobacco budworm in cotton. *Southwestern Entomologist Supplement*(15): 173-186.

Minton, E. B. and M. W. Ebelhar (1991). Potassium and aldicarb-disulfoton effects on verticillium wilt, yield, and quality of cotton. *Crop Science* 31(1): 209-212.

- Mo, H. and Z. Li (1991). Augmented NC II design and genetic model test. *Acta Agronomica Sinica* 17(1): 1-9.
- Monks, C. D. and P. A. Banks (1991). Rotational crop response to chlorimuron, clomazone, and imazaquin applied the previous year. *Weed Science* 39(4): 629-633.
- Navasero, R. C. and S. B. Ramaswamy (1991). Morphology of leaf surface trichomes and its influence on egg-laying by *Heliothis virescens*. *Crop Science* 31(2): 342-353.
- Navon, A., M. V. Melamed, et al. (1991). Effects of cotton cultivars on feeding of *Heliothis armigera* and *Spodoptera littoralis* larvae and on oviposition of *Bemisia tabaci*. *Agriculture Ecosystems & Environment* 35(1): 73-80.
- Oliver, L. R., J. M. Chandler, et al. (1991). Influence of geographic region on jimsonweed (*Datura stramonium*) interference in soybeans (*Glycine max*) and cotton (*Gossypium hirsutum*). *Weed Science* 39(4): 585-589.
- Oosterhuis, D. M., R. E. Hampton, et al. (1991). Water deficit effects on the cotton leaf cuticle and the efficiency of defoliants. *Journal Of Production Agriculture* 4(2): 260-265.
- Peng, S. and D. R. Krieg (1991). Single leaf and canopy photosynthesis response to plant age in cotton. *Agronomy Journal* 83(4): 704-708.
- Percy, R. G. and E. L. Turcotte (1991). Early-maturing, short-statured American Pima cotton parents improve agronomic traits of interspecific hybrids. *Crop Science* 31(3): 709-712.
- Pizzinatto, M. A., E. Cia, et al. (1991). Transmission of *Colletotrichum gossypii* var. *cephalosporioides* by cotton seeds. *Summa Phytopathologica* 17(3-4): 207-217.
- Pizzinatto, M. A. and J. O. M. Menten (1991). Pathogenicity of eight *Fusarium* spp. isolated from seeds to cotton seedlings. *Summa Phytopathologica* 17(2): 124-134.
- Pizzinatto, M. A., J. O. M. Menten, et al. (1991). Relationship between density and quality of cotton seeds. *Bragantia* 50(2): 269-289. {a} Secao Fitopatologia, Inst. Agronomico, Caixa Postal 28, 13001 Campinas, SP
- Plaut, Z. and E. Federman (1991). Acclimation of carbon dioxide assimilation in cotton leaves to water stress and salinity. *Plant Physiology* 97(2): 515-522.
- Plekhanov, L. S., L. B. Ivleva, et al. (1991). ATPase activity in plasma membranes from cotton plant roots. *Fiziologiya Rastanii* 38(2): 273-280.
- Polanowski, A. J. and R. L. Obendorf (1991). Soybean isocitrate lyase: Purification, properties, immunoassay and amino-terminal sequence. *Plant Physiology And Biochemistry* 29(2): 119-130.
- Purohit, M. S. and A. D. Deshpande (1991). Effect of inorganic fertilizers on the cotton leaf hopper *Amrasca biguttula biguttula* Ishida in relation to plant protection. *Indian Journal Of Plant Protection* 19(2): 133-138.
- Qi, Z. S., D. Q. Qian, et al. (1991). Induction of parthenogenesis, and chromosome behavior in plants of parthenogenetic origin in cotton (*Gossypium hirsutum*). *Genome* 34(2): 255-260.
- Quisenberry, J. E. and B. L. McMichael (1991). Genetic variation among cotton germplasm for water-use efficiency. *Environmental And Experimental Botany* 31(4): 453-460.
- Radin, J. W., J. R. Mauney, et al. (1991). Effects of nitrogen fertility on water potential of irrigated cotton. *Agronomy Journal* 83(4): 739-743.
- Rahman, S., M. Ahmad, et al. (1991). Genetic architecture of yield components in cotton (*Gossypium hirsutum* L.) I. *Sarhad Journal Of Agriculture* 7(2): 113-128.

- Rao, N. V., A. S. Reddy, et al. (1991). Intraplant distribution of whitefly, *Bemisia tabaci* Genn. on cotton, *Gossypium hirsutum* Linn. *Journal Of Insect Science* 4(1): 32-36.
- Rasulov, B. K. and K. A. Asrorov (1991). Photosynthesis and productivity of cotton plants. *Fiziologiya Rastenii* 38(2): 336-345.
- Rasulov, B. K., V. M. Oya, et al. (1991). Temperature dependence of photosynthesis and its relation to the kinetics of ribulose biphosphate carboxylase and the ribulose-1,5-biphosphate pool. *Fiziologiya Rastenii* 38(1): 15-23.
- Razzouk, S. and W. J. Whittington (1991). Effects of salinity on cotton yield and quality. *Field Crops Research* 26(3-4): 305-314.
- Reddy, V. R., D. N. Baker, et al. (1991). Temperature effects on cotton canopy growth, photosynthesis, and respiration. *Agronomy Journal* 83(4): 699-704.
- Reddy, V. R., K. R. Reddy, et al. (1991). Agroclimatology and modeling: Temperature effect on growth and development of cotton during the fruiting period. *Agronomy Journal* 83(1): 211-217.
- Rikin, A. (1991). Temperature-induced phase shifting of circadian rhythms in cotton seedlings as related to variations in chilling resistance. *Planta* 185(3): 407-414.
- Rodriguez, K. R., D. G. Robertson, et al. (1991). Cotton as a rotation crop for the management of *Meloidogyne arenaria* and *Sclerotium rolfsii* in peanut. *Journal Of Nematology* 23(4 Suppl): 652-657.
- Rooney, W. L. and D. M. Stelly (1991). Preferential transmission and somatic elimination of a *Gossypium sturtianum* chromosome in *Gossypium hirsutum*. *Journal Of Heredity* 82(2): 151-155.
- Rooney, W. L., D. M. Stelly, et al. (1991). Identification of four *Gossypium sturtianum* monosomic alien addition derivatives from a backcrossing program with *Gossypium hirsutum*. *Crop Science* 31(2): 337-341.
- Rosenthal, W. D. and T. J. Gerik (1991). Radiation use efficiency among cotton cultivars. *Agronomy Journal* 83(4): 655-658.
- Russell, S. G., T. J. Monaco, et al. (1991). Influence of soil moisture on phytotoxicity of cinmethylin to various crops. *Weed Science* 39(3): 402-407.
- Sanni, A. I. and D. N. Ogbonna (1991). The production of owoh: A Nigerian fermented seasoning agent from cotton seed (*Gossypium hirsutum* L.). *Food Microbiology* 8(3): 223-230.
- Saranga, Y., J. Rudich, et al. (1991). The relations between leaf water potential of cotton plants and environmental and plant factors. *Field Crops Research* 28(1-2): 39-46.
- Segarra, E., J. W. Keeling, et al. (1991). Tillage and cropping system effects on cotton yield and profitability on the Texas Southern High Plains. *Journal Of Production Agriculture* 4(4): 566-571.
- Shang, X. M., J. Y. Huang, et al. (1991). Buffer capacity of cotton cells and effects of extracellular pH on growth and somatic embryogenesis in cotton cell suspensions. *In Vitro Cellular & Developmental Biology Plant* 27p 3: 147-152.
- Sharma, D. P., K. N. Singh, et al. (1991). Effect of subsurface drain spacing on growth and yield behaviour of cotton on a saline soil. *Journal Of The Indian Society Of Soil Science* 39(4): 757-761.
- Simwat, G. S., A. K. Dhawan, et al. (1991). Effect of insecticides on some parameters of yield in *Gossypium hirsutum* L. *Journal Of Insect Science* 4(2): 190-192.

- Simwat, G. S., A. K. Dhawan, et al. (1991). Evaluation of methomyl alone and in combination with other insecticides for bollworm control on cotton in Punjab. *Journal Of Research Punjab Agricultural University* 28(2): 215-218.
- Simwat, G. S., A. K. Dhawan, et al. (1991). Effect of acid-delinting of cotton seed on the survival of diapausing larvae of pink bollworm. *Journal Of Insect Science* 4(1): 45-49.
- Singh, A. R., S. B. Choulwar, et al. (1991). Hard seededness and quality in relation to picking and locations in cotton. *Seed Research* 19(1): 31-36.
- Singh, D., M. S. Brar, et al. (1991). Response to potassium application and its critical levels for American cotton (*Gossypium hirsutum* L.). *Journal Of The Indian Society Of Soil Science* 39(3): 494-499.
- Singh, D. and R. K. Sahay (1991). Critical leaf-water potential in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 61(11): 822-825.
- Singh, I. P. and D. P. Singh (1991). Estimation of extent of natural crossing in upland cotton (*Gossypium hirsutum*) in Haryana (India). *Indian Journal Of Agricultural Sciences* 61(9): 629-633.
- Singh, M., V. P. Singh, et al. (1991). Improvement of yield and fiber strength in medium and superior-medium staple upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 61(1): 11-15.
- Singh, P., T. H. Singh, et al. (1991). Genetical control of gossypol content in intervarietal crosses of upland cotton. *Crop Improvement* 18(2): 141-143. Cotton Res. Lab., Punjab Agric. Univ., Ludhiana, India
- Singh, T. H., G. S. Chahal, et al. (1991). Extent of natural crossing in cotton in Punjab. *Crop Improvement* 18(1): 54-57.
- Slosser, J. E. (1991). Effects of windbreaks on overwintering of boll weevils. U S Forest Service General Technical Report Rm(204): 13-16.
- Snipes, C. E. (1991). Preemergence velvetleaf (*Abutilon theophrasti*) control in cotton (*Gossypium hirsutum*). *Weed Technology* 5(1): 142-146.
- Snipes, C. E., J. E. Street, et al. (1991). Cotton (*Gossypium hirsutum*) response to simulated triclopyr drift. *Weed Technology* 5(3): 493-498.
- Stark, C. (1991). Osmotic adjustment and growth of salt-stressed cotton as improved by a bioregulator. *Journal Of Agronomy And Crop Science* 167(5): 326-334.
- Starr, J. L. and R. D. Martyn (1991). Reaction of cotton cultivars to *Fusarium* wilt and root-knot nematodes. *Nematologica* 21(1): 51-58.
- Stoilova, A. (1991). Type of development in hybrids of the backcrosses of *Gossypium hirsutum* L. X *Gossypium barbadense* L. to both parental species: I. Characterization of BC-1. *Genetika I Selektsiya* 24(4): 237-243.
- Stoilova, A. (1991). Gamma-irradiation of interspecific hybrids *Gossypium hirsutum* L. X *Gossypium barbadense* L: II. Possibilities for increasing genetic diversity and enhancing form-developing processes. *Genetika I Selektsiya* 24(1): 3-10.
- Stoilova, A. (1991). Gamma-irradiation of the interspecific hybrids *Gossypium hirsutum* L. X *Gossypium barbadense* L: I. Reaction of hybrids to irradiation in the year of treatment. *Genetika I Selektsiya* 23(5): 424-429.
- Subramanian, M. V., G. V. Loney, et al. (1991). Acetolactate synthase inhibiting herbicides bind to the regulatory site. *Plant Physiology* 96(1): 310-313.

- Suttle, J. C. and J. F. Hultstrand (1991). Ethylene-induced leaf abscission in cotton seedlings: The physiological bases for age-dependent differences in sensitivity. *Plant Physiology* 95(1): 29-33.
- Tanaka, M. A. S. and J. O. M. Menten (1991). Comparison of methods for cotton seed inoculation with *Colletotrichum gossypii* var. *cephalosporioides* and *Colletotrichum gossypii*. *Summa Phytopathologica* 17(3-4): 218-226.
- Tariq, M., M. A. Khan, et al. (1991). Identification of parents for hybridization through combining ability analysis in upland cotton. *Sarhad Journal Of Agriculture* 7(5): 633-641.
- Terry, L. I. (1991). Pest and predator populations following early-season cotton insect control in Arizona (USA). *Southwestern Entomologist* 16(1): 51-62.
- Thomas, R. B. and B. R. Strain (1991). Root restriction as a factor in photosynthetic acclimation of cotton seedlings grown in elevated carbon dioxide. *Plant Physiology* 96(2): 627-634.
- Tonhasca, A. J. and R. G. Luttrell (1991). Seasonal history and damage caused by *Neurocolpus nubilus* (Hemiptera: Miridae) on cotton in northern Mississippi (USA). *Environmental Entomology* 20(2): 742-748.
- Tracy, P. W. (1991). Evaluation of urea and urea-ammonium nitrate with and without dicyandiamide as cotton nitrogen sources in Missouri (USA). *Journal Of Fertilizer Issues* 8(1): 1-5.
- Treacy, M. F., J. H. Benedict, et al. (1991). Mineral oil: Enhancement of field efficacy of a pyrethroid insecticide against the boll weevil (Coleoptera: Curculionidae). *Journal Of Economic Entomology* 84(2): 659-663.
- Trolinder, N. L. and X. Shang (1991). In vitro selection and regeneration of cotton resistant to high temperature stress. *Plant Cell Reports* 10(9): 448-452.
- Van, H. H. G., M. S. Hill, et al. (1991). Diallel analyses of cotton (*Gossypium hirsutum*, L.) yield and fibre characteristics over two locations. *Applied Plant Science* 5(2): 68-71.
- Virk, P. S., H. S. Kalsy, et al. (1991). An assessment of the breeding potential of a *Gossypium hirsutum* L. cross. *Crop Improvement* 18(1): 7-13.
- Voo, K. S., C. L. Rugh, et al. (1991). Indirect somatic embryogenesis and plant recovery from cotton (*Gossypium hirsutum* L.). *In Vitro Cellular & Developmental Biology Plant* 27p 3: 117-124.
- Waller, G. D. and A. N. Mamood (1991). Upland and pima cotton as pollen donors for male-sterile upland seed parents. *Crop Science* 31(2): 265-266.
- Wharton, T. F., J. N. Jenkins, et al. (1991). Productivity and tobacco budworm resistance of cotton developed between 1890 and 1986. *Crop Science* 31(1): 29-32.
- Wilson, F. D. (1991). Combining ability for yield characteristics and earliness of pink bollworm-resistant cotton. *Crop Science* 31(4): 922-925.
- Wilson, F. D. and J. K. Brown (1991). Inheritance of response to cotton leaf crumple virus infection in cotton. *Journal Of Heredity* 82(6): 508-509.
- Wilson, F. D., H. M. Flint, et al. (1991). Reduction in insecticide use associated with cotton resistant to pink bollworm. *Crop Science* 31(2): 363-366.
- Wilson, L. T., P. J. Trichilo, et al. (1991). Natural enemies of spider mites (Acari: Tetranychidae) on cotton: Density regulation or casual association? *Environmental Entomology* 20(3): 849-856.
- Wilson, L. T., P. J. Trichilo, et al. (1991). Spider mite (Acari: Tetranychidae) infestation rate and initiation: Effect on cotton yield. *Journal Of Economic Entomology* 84(2): 593-600.

- Winter, K. and M. Koeniger (1991). Dry matter production and photosynthetic capacity in *Gossypium hirsutum* L. under conditions of slightly suboptimum leaf temperatures and high levels of irradiance. *Oecologia* 87(2): 190-197.
- Wullschleger, S. D. and D. M. Oosterhuis (1991). Photosynthesis, transpiration, and water-use efficiency of cotton leaves and fruit. *Photosynthetica* 25(4): 505-515.
- Wullschleger, S. D., D. M. Oosterhuis, et al. (1991). Evidence for light-dependent recycling of respired carbon dioxide by the cotton fruit. *Plant Physiology* 97(2): 574-579.
- Yadav, K. S., S. C. Deshmukh, et al. (1991). Performance of promising varieties of upland cotton (*Gossypium hirsutum*) at different fertility levels and plant densities under rainfed condition. *Indian Journal Of Agronomy* 36(Suppl): 173-176.
- York, A. C., D. L. Jordan, et al. (1991). Insecticides modify cotton (*Gossypium hirsutum*) response to clomazone. *Weed Technology* 5(4): 729-735.
- Yunusova, S. G., S. D. Gusakova, et al. (1991). Comparative study of fatty acid composition in the seeds of some strains of a genetic collection of *Gossypium hirsutum*. *Khimiya Prirodnikh Soedinenii*(2): 173-176.
- Zhang, F., V. Romheld, et al. (1991). Release of zinc mobilizing root exudates in different plant species as affected by zinc nutritional status. *Journal Of Plant Nutrition* 14(7): 675-686.
- Zhang, X., J. Sun, et al. (1991). Somatic embryogenesis and plant regeneration in upland cotton. *Acta Genetica Sinica* 18(5): 461-467.
- Zodape, S. T. and V. D. Chauhan (1991). Effect of pulsed concentrated solar radiation on the activity of catalase and peroxidase in cotton (*Gossypium hirsutum* L.). *Indian Journal Of Plant Physiology* 34(3): 209-214.
- Abdurakhmanova, Z. N., K. A. Aliev, et al. (1990). Photosynthetic carbon metabolism and the transformation of (carbon-14) glycolic acid in cotton (*Gossypium hirsutum* L.) leaf ontogeny. *Fiziologiya Rastenii* 37(4): 675-681.
- Afek, U., E. Rinaldelli, et al. (1990). Mycorrhizal species, root age, and position of mycorrhizal inoculum influence colonization of cotton, onion, and pepper seedlings. *Journal Of The American Society For Horticultural Science* 115(6): 938-942.
- Al, M. A. H. (1990). An ultrastructural study of two cotton lines with intermediate levels of resistance to *Xanthomonas campestris* pathovar *malvacearum*. *Journal Of The University Of Kuwait* 17(1): 127-136.
- Ali, A., R. G. Luttrell, et al. (1990). Effects of temperature and larval diet on development of the fall armyworm (Lepidoptera: Noctuidae). *Annals Of The Entomological Society Of America* 83(4): 725-733.
- Allen, S. J. (1990). *Thielaviopsis basicola*, a new record on cotton in Australia. *Australasian Plant Pathology* 19(1): 24-25.
- Alsaadawi, I. S., F. A. K. Sakeri, et al. (1990). Allelopathic inhibition of *Cynodon dactylon* (L.) Pers. and other plant species by *Euphorbia prostrata* L. *Journal Of Chemical Ecology* 16(9): 2747-2754.
- Altman, D. W., R. D. Stipanovic, et al. (1990). Terpenoids in foliar pigment glands of A, D, and AD genome cottons: Introgression potential for pest resistance. *Journal Of Heredity* 81(6): 447-454.
- Annadurai, R. S., S. Murugesan, et al. (1990). Age-correlated tissue preferences of *Heliothis armigera* (Hubner) and *Spodoptera litura* (F) with special reference to phenolic substances. *Proceedings Of The Indian Academy Of Sciences Animal Sciences* 99(4): 317-326.

- Baehrecke, E. H., S. B. Vinson, et al. (1990). Foraging behavior of *Campoletis sonorensis* in response to *Heliothis virescens* and cotton plants. *Entomologia Experimentalis Et Applicata* 55(1): 47-58.
- Basford, K. E., P. M. Kroonenberg, et al. (1990). Multiattribute evaluation of regional cotton variety trials. *Theoretical And Applied Genetics* 79(2): 225-234.
- Basra, A. S., R. S. Sarlach, et al. (1990). Sucrose hydrolysis in relation to development of cotton (*Gossypium* spp.) fibers. *Indian Journal Of Experimental Biology* 28(10): 985-988.
- Bauer, P. J. and J. T. Cothren (1990). Growth-promoting activity of chlordimeform. *Agronomy Journal* 82(1): 73-75.
- Ben, P. A. and D. N. Baker (1990). Taproot restriction effects on growth, earliness, and dry weight partitioning of cotton. *Crop Science* 30(4): 809-814.
- Bhale, N. L. and M. G. Bhat (1990). Investigations on exploitation of heterosis in cotton (*Gossypium hirsutum* L.) using male sterility. *Indian Journal Of Genetics & Plant Breeding* 50(1): 37-44.
- Botha, J. H. and A. J. Scholtz (1990). Further studies on the effect of spider mites (Acari: Tetranychidae) on cotton yields. *Phytophylactica* 22(1): 77-86.
- Boucounis, T. G., T. Whitwell, et al. (1990). Correlation of bioassay crop growth with cinmethylin and chlorimuron application rates for two soils. *Hortscience* 25(5): 536-538.
- Bourland, F. M. and C. E. J. Watson (1990). COTMAP, a technique for evaluating structure and yield of cotton plants. *Crop Science* 30(1): 224-226.
- Bozcuk, S. (1990). Interaction between salt and kinetin on seed germination of some culture plants. *Doga Turk Botanik Dergisi* 14(2): 139-149.
- Bradow, J. M. and W. J. J. Connick (1990). Volatile seed germination inhibitors from plant residues. *Journal Of Chemical Ecology* 16(3): 645-666.
- Brody, A. K., R. Karban, et al. (1990). Inverse relationship between cotton plant density and verticillium wilt incidence and severity: Evidence for an alternative hypothesis. *Crop Protection* 9(3): 174-176.
- Brouder, S. M. and K. G. Cassman (1990). Root development of two cotton cultivars in relation to potassium uptake and plant growth in a vermiculitic soil. *Field Crops Research* 23(3-4): 187-204.
- Bryson, C. T. (1990). Interference and critical time of removal of hemp sesbania (*Sesbania exaltata*) in cotton (*Gossypium hirsutum*). *Weed Technology* 4(4): 833-837.
- Bryson, C. T. (1990). Differential growth of bermudagrass (*Cynodon dactylon*) biotypes in cotton (*Gossypium hirsutum*). *Weed Technology* 4(3): 612-614.
- Burke, J. J. (1990). Variation among species in the temperature dependence of the reappearance of variable fluorescence following illumination. *Plant Physiology* 93(2): 652-656.
- Burke, J. J., J. L. Hatfield, et al. (1990). A thermal stress index for cotton. *Agronomy Journal* 82(3): 526-530.
- Butler, G. D. J. and T. J. Henneberry (1990). Cottonseed oil and safer insecticidal soap: Effects on cotton and vegetable pests and phytotoxicity. *Southwestern Entomologist* 15(3): 257-264.
- Byzova, M. V., A. S. Kraev, et al. (1990). Molecular characterization of chalcone synthetase gene family of the tetraploid cotton cultivar *Gossypium hirsutum* 108F. *Doklady Akademii Nauk Sssr* 311(3): 742-744.

- Cakmak, I. and H. Marschner (1990). Decrease in nitrate uptake and increase in proton release in zinc deficient cotton, sunflower and buckwheat plants. *Plant And Soil* 129(2): 261-268.
- Callahan, F. E., M. L. Ghirardi, et al. (1990). A novel metabolic form of the 32kDa-D1 protein in the grana-localized reaction center of photosystem II. *Journal Of Biological Chemistry* 265(26): 15357-15360.
- Cassman, K. G., T. A. Kerby, et al. (1990). Potassium nutrition effects of lint yield and fiber quality of Acala cotton. *Crop Science* 30(3): 672-677.
- Chaturvedi, S., H. S. Randhawa, et al. (1990). Cottonseed extract versus pharmamedia for the in vitro mold-yeast conversion of *Blastomyces dermatitidis*. *Journal Of Medical And Veterinary Mycology* 28(2): 139-146.
- Chetverikov, A. G. and V. F. Gachkovskii (1990). Effect of water deficiency and salt stress on energy supply systems in cotton. *Izvestiya Akademii Nauk Sssr Seriya Biologicheskaya*(4): 532-541.
- Choudhari, P. N., D. N. Borole, et al. (1990). Correlation and path analysis in American cotton. *Journal Of Maharashtra Agricultural Universities* 15(3): 289-291.
- Chu, C. C. and L. A. Bariola (1990). In vivo cotton stem and petiole nitrate reductase activity. *Agronomy Journal* 82(2): 221-224.
- Cotty, P. J. (1990). Effect of atoxigenic strains of *Aspergillus flavus* on aflatoxin contamination of developing cottonseed. *Plant Disease* 74(3): 233-235.
- Daigle, D. J., W. J. J. Connick, et al. (1990). Invert emulsions: Carrier and water source for the mycoherbicide, *Alternaria cassiae*. *Weed Technology* 4(2): 327-331.
- Daly, J. C. and G. P. Fitt (1990). Monitoring for pyrethroid resistance in relation to body weight in adult *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). *Journal Of Economic Entomology* 83(3): 705-709.
- De, K. J., B. L. P. De, et al. (1990). The relative sensitivity to plant water stress during the reproductive phase of upland cotton (*Gossypium hirsutum* L.). *Irrigation Science* 11(4): 239-244.
- Dhawan, A. K., G. S. Simwat, et al. (1990). Incidence of different insect pests on LH900 upland cotton sown on different dates. *Journal Of Research Punjab Agricultural University* 27(2): 244-252.
- Dickens, J. C. and E. E. Moorman (1990). Maturation and maintenance of electroantennogram responses to pheromone and host odors in boll weevils, *Anthonomus grandis* Boh. (Coleoptera, Curculionidae) fed their host plant or an artificial diet. *Journal Of Applied Entomology* 109(5): 470-480.
- Diomaiuto, J. and C. Hubac (1990). Morphological and histological repercussions of an increase in drought resistance in cotton plants (*Gossypium hirsutum* L.) produced using far-red light. *Phytomorphology* 40(3-4): 383-394.
- Dugas, W. A. (1990). Comparative measurement of stem flow and transpiration in cotton. *Theoretical And Applied Climatology* 42(4): 215-222.
- Dutt, A. K. and J. L. Saha (1990). Insect pest management of winter planted cotton in coastal rice fallows of West Bengal (India). *Tropical Pest Management* 36(2): 89-92.
- Ephrath, J. E., A. Marani, et al. (1990). Effects of moisture stress on stomatal resistance and photosynthetic rate in cotton (*Gossypium hirsutum* new record): 1. Controlled levels of stress. *Field Crops Research* 23(2): 117-132.
- Ethridge, D. E., R. T. Ervin, et al. (1990). Economic weed control in high plains cotton. *Journal Of Production Agriculture* 3(2): 246-252.

- Finer, J. J. and M. D. McMullen (1990). Transformation of cotton (*Gossypium hirsutum* L.) via particle bombardment. *Plant Cell Reports* 8(10): 586-589.
- Flint, H. M. and N. J. Parks (1990). Infestation of germplasm lines and cultivars of cotton in Arizona (USA) by whitefly nymphs (Homoptera: Aleyrodidae). *Journal Of Entomological Science* 25(2): 223-229.
- Gawel, N. J. and C. D. Robacker (1990). Somatic embryogenesis in two *Gossypium hirsutum* genotypes on semi-solid versus liquid proliferation media. *Plant Cell Tissue And Organ Culture* 23(3): 201-204.
- Gawel, N. J. and C. D. Robacker (1990). Genetic control of somatic embryogenesis in cotton petiole callus cultures. *Euphytica* 49(3): 249-254.
- Geng, S., B. Hu, et al. (1990). Quantification and classification of locational effects on cotton cultivar testing programs. *Agronomy Journal* 82(3): 514-518.
- Gill, K. S., T. H. Singh, et al. (1990). Path analysis of genotype-environment interactions in upland cotton. *Crop Improvement* 17(1): 27-30.
- Golan, G. A., B. Hankamer, et al. (1990). Hydroxyproline and proline content of cell walls of sunflower, peanut and cotton grown under salt stress. *Plant Science* 69(1): 27-32.
- Gordon, T. R. and D. Okamoto (1990). Colonization of crop residue by *Fusarium oxysporum* f. sp. melonis and other species of *Fusarium*. *Phytopathology* 80(4): 381-386.
- Gordon, W. B., D. H. Rickerl, et al. (1990). Effects of nitrogen, dicyandiamide, and tillage on cotton nitrogen concentration and yield. *Agronomy Journal* 82(3): 597-600.
- Goswami, C. L. (1990). Hormonal regulation of fertility-induced changes in stomatal diffusive resistance in waterlogged cotton (*Gossypium hirsutum*, cultivar H-777). *Indian Journal Of Experimental Biology* 28(6): 585-587.
- Green, C. C. and T. W. Culp (1990). Simultaneous improvement of yield, fiber quality, and yarn strength in upland cotton. *Crop Science* 30(1): 66-69.
- Gulin, V. V., K. Y. Saidov, et al. (1990). Heterogeneity of cotton *Gossypium hirsutum* L. cultivar populations for gene biochemical markers. *Genetika* 26(3): 485-490.
- Gulyamov, D. B., E. G. Romanko, et al. (1990). Characteristics of RNA synthesis in chromatin isolated from cotton plant rootlets. *Fiziologiya Rastanii* 37(2): 279-283.
- Hagedorn, C., N. Nelson, et al. (1990). Evaluation of a *Pseudomonas fluorescens* strain for repression of seedling disease in cotton. *Virginia Journal Of Science* 41(4 Part B): 492-500.
- Ham, J. M., J. L. Heilman, et al. (1990). Determination of soil water evaporation and transpiration from energy balance and stem flow measurements. *Agricultural And Forest Meteorology* 52(3-4): 287-302.
- Hampton, R. E., S. D. Wullschleger, et al. (1990). Impact of verticillium wilt on net photosynthesis, respiration and photorespiration in field-grown cotton (*Gossypium hirsutum* L.). *Physiological And Molecular Plant Pathology* 37(4): 271-280.
- He, Z., X. Min, et al. (1990). Study on effects of DPC on endogenous hormone levels and the development of cotton bolls. *Acta Agronomica Sinica* 16(3): 254-258.
- Hendrix, D. L. (1990). Carbohydrates and carbohydrate enzymes in developing cotton ovules. *Physiologia Plantarum* 78(1): 85-92.
- Hitron, O. and B. Zur (1990). Differences in stomatal response within a cotton canopy. *Biotronics* 19: 39-48.

- Hodgson, A. S., G. A. Constable, et al. (1990). A comparison of drip and furrow irrigated cotton on a cracking clay soil: 2. Water use efficiency, waterlogging, root distribution and soil structure. *Irrigation Science* 11(3): 143-148.
- Hons, F. M., N. W. Hopper, et al. (1990). Applied phosphorus and potassium effects on the emergence, yield, and planting seed quality of cotton. *Journal Of Production Agriculture* 3(3): 337-340.
- Howard, D. D. and P. E. Hoskinson (1990). Effect of starter nutrient combinations and nitrogen rate on no-tillage cotton. *Journal Of Fertilizer Issues* 7(1): 6-9.
- Hubbs, C. W. and T. L. Lavy (1990). Dissipation of norflurazon and other persistent herbicides in soil. *Weed Science* 38(1): 81-88.
- Hulugalle, N. R. (1990). Alleviation of soil constraints to crop growth in the upland alfisols and associated soil groups of the West African Sudan savannah by tied ridges. *Soil & Tillage Research* 18(2-3): 231-248.
- Inoue, Y., B. A. Kimball, et al. (1990). Remote estimation of leaf transpiration rate and stomatal resistance based IR thermometry. *Agricultural And Forest Meteorology* 51(1): 21-34.
- Irvine, J. M., J. V. Oakes, et al. (1990). A rapid screen for the detection of specific DNA sequences in plants. *Genetic Analysis Techniques And Applications* 7(2): 25-31.
- Jackson, B. S., T. J. Gerik, et al. (1990). Use of COTTAM for scheduling limited irrigation. *Journal Of Production Agriculture* 3(4): 420-425.
- Jacoby, P. W., C. H. Meadors, et al. (1990). Effects of triclopyr, clopyralid, and picloram on growth and production of cotton. *Journal Of Production Agriculture* 3(3): 297-301.
- Jenkins, J. N., J. C. J. McCarty, et al. (1990). Fruiting efficiency in cotton: Boll size and boll set percentage. *Crop Science* 30(4): 857-860.
- Jenkins, J. N., J. C. J. McCarty, et al. (1990). Effectiveness of fruiting sites in cotton: Yield. *Crop Science* 30(2): 365-369.
- Jones, R. G., P. J. Bauer, et al. (1990). Effect of reduced rates of ethephon on late-season insect oviposition and feeding sites in cotton. *Journal Of Entomological Science* 25(2): 246-252.
- Kabar, K. (1990). Comparison of Kinetin and Gibberellic Acid Effects on Seed Germination under Saline Conditions. *Phyton Horn* 30(2): 291-298. Atatürk Universitesi, K. Karabakir Egitim Fakultesi, Biyoloji Bolumu, TR-25240 Erzurum, Turkey
- Katageri, I. S., S. N. Kadapa, et al. (1990). A case study on exploitation of heterosis using multiple pest tolerance in *Gossypium hirsutum* L. *Indian Journal Of Genetics & Plant Breeding* 50(4): 314-319.
- Kennedy, C. W., W. C. J. Smith, et al. (1990). Computer programs for recording and analyzing cotton fruit location maps. *Journal Of Agronomic Education* 19(2): 167-171.
- Kerby, T. A., K. G. Cassman, et al. (1990). Genotypes and plant densities for narrow-row cotton systems: II. Leaf area and dry-matter partitioning. *Crop Science* 30(3): 649-653.
- Kerby, T. A., K. G. Cassman, et al. (1990). Genotypes and plant densities for narrow-row cotton system: I. Height, nodes, earliness, and location of yield. *Crop Science* 30(3): 644-649.
- Klich, M. A. (1990). The degree of susceptibility of nectary-inoculated cotton flowers and bolls to subsequent seed infection by *Aspergillus flavus* is determined at or before anthesis. *Applied And Environmental Microbiology* 56(8): 2499-2502.

- Kohel, R. J. and S. C. McMichael (1990). Immature fiber mutant of upland cotton. *Crop Science* 30(2): 419-421.
- Kojnov, G. and A. Stoilova (1990). Hybridization between the species *Gossypium hirsutum* L. and *Gossypium barbadense* L: A promising method for developing valuable forms and cultivars. *Genetika I Seleksiya* 23(4): 359-366.
- Kuznetsov, V. V., B. T. Khydyrov, et al. (1990). Common systems of cotton plant resistance to salinity and high temperatures: Facts and hypotheses. *Fiziologiya Rastenii* 37(5): 987-996.
- Kuznetsov, V. V., B. V. Roshchupkin, et al. (1990). Effect of salinization on interaction between the initial and adaptive resistance in plants. *Doklady Akademii Nauk Sssr* 314(2): 509-512.
- Ladyman, J. A. R., J. R. Goss, et al. (1990). Activity of a novel chemical hybridizing agent in field-grown cotton. *Field Crops Research* 23(3-4): 177-186.
- Lakshmanan, P., R. Jeyarajan, et al. (1990). A boll rot of cotton caused by *Corynespora cassicola* in Tamil Nadu, India. *Phytoparasitica* 18(2): 171-174.
- Lawrence, G. W., K. S. McLean, et al. (1990). Response of *Rotylenchulus reniformis* to nematicide applications on cotton. *Journal Of Nematology* 22(4 Suppl): 707-711.
- Lu, Z., H. Cheng, et al. (1990). The genetic effects of selfing and studies on the methods of cultivar maintenance in cotton (*Gossypium hirsutum* L.). *Scientia Agricultura Sinica* 23(1): 69-75.
- Malek, F. and P. Zandi (1990). Determination of free gossypol in Iranian cottonseed and cottonseed cake. *Food Chemistry* 37(4): 289-296.
- Malik, M. N. A., F. I. Chaudhry, et al. (1990). Effect of PIX on yield and growth of cotton (*Gossypium hirsutum* L.). *Sarhad Journal Of Agriculture* 6(1): 67-70.
- Malik, M. N. A. and M. I. Makhdum (1990). Phosphorus fertilizer experiments with cotton in Bahawalpur region (Pakistan). *Sarhad Journal Of Agriculture* 6(5): 521-528.
- Malik, M. N. A., M. I. Makhdum, et al. (1990). Response of cotton to different doses of phosphatic fertilizer at Sahiwal (Punjab) (Pakistan). *Sarhad Journal Of Agriculture* 6(2): 169-174.
- Markova, I. A., A. K. Tonkikh, et al. (1990). Detection of auxin-binding sites in cotton seedlings. *Fiziologiya Rastenii* 37(4): 727-732.
- Masud, S. M., R. D. Lacewell, et al. (1990). Economic implications and simulations of fleahopper and bollworm management decisions for cotton in the Texas Coastal Bend (USA). *Journal Of Production Agriculture* 3(2): 200-208.
- McAuslane, H. J., S. B. Vinson, et al. (1990). Effect of host diet on flight behavior of the parasitoid *Campoletis sonorensis* (Hymenoptera: Ichneumonidae). *Journal Of Entomological Science* 25(4): 562-570.
- McMillan, K. D. and A. Rikin (1990). Relationships between circadian rhythm of chilling resistance and acclimation to chilling in cotton seedlings. *Planta* 182(3): 455-460.
- Meek, B. D., W. R. Detar, et al. (1990). Infiltration rate as affected by an alfalfa and no-till cotton cropping system. *Soil Science Society Of America Journal* 54(2): 505-508.
- Melero, V. J. M. and D. R. M. Jimenez (1990). Etiology, incidence, and distribution of cotton seedling damping-off in southern Spain. *Plant Disease* 74(8): 597-600.
- Mennen, H., B. Jacoby, et al. (1990). Is sodium proton antiport ubiquitous in plant cells? *Journal Of Plant Physiology* 137(2): 180-183.

Mercer, K. L., J. A. Pawlak, et al. (1990). Distance-of-influence of devil's claw (*Proboscidea louisianica*) on cotton (*Gossypium hirsutum*). *Weed Technology* 4(1): 87-91.

Micinski, S., P. D. Colyer, et al. (1990). Effects of planting date and early-season pest control on yield in cotton. *Journal Of Production Agriculture* 3(4): 597-602.

Mohod, V. K. and S. B. Parde (1990). Evaluation of isolines of upland cotton for some of the commercially important plant traits. *Asian Journal Of Plant Science* 2(2): 49-53.

Morris, D. A. and D. K. Small (1990). Auxin transport and the regulation of petiole abscission in cotton (*Gossypium hirsutum* L.): A comparison of IAA and phenylacetic acid. *Plant Growth Regulation* 9(3): 201-214.

Morrison, J. E. J., T. J. Gerik, et al. (1990). A no-tillage farming system for clay soils. *Journal Of Production Agriculture* 3(2): 219-227.

Morrow, M. R. and D. R. Krieg (1990). Cotton management strategies for a short growing season environment: Water-nitrogen considerations. *Agronomy Journal* 82(1): 52-56.

Mueller, F. and S. Jiang (1990). Uptake, distribution and metabolism of carbon-14 labelled carbendazim in *Gossypium hirsutum*. *Zeitschrift Fuer Pflanzenkrankheiten Und Pflanzenschutz* 97(2): 216-229.

Muhammad, N. and J. E. Jones (1990). Genetics of resistance to reniform nematode in upland cotton. *Crop Science* 30(1): 13-16.

Mullins, G. L. and C. H. Burmester (1990). Dry matter, nitrogen, phosphorus, and potassium accumulation by four cotton varieties. *Agronomy Journal* 82(4): 729-736.

Narbutth, E. V. and R. J. Kohel (1990). Inheritance and linkage analysis of a new fiber mutant in cotton. *Journal Of Heredity* 81(2): 131-133.

Nayyar, H., K. Kaur, et al. (1990). Activity profiles of pyruvate orthophosphate dikinase and phosphoenolpyruvate carboxylase in developing cotton fibers. *Beitraege Zur Biologie Der Pflanzen* 65(3): 319-324.

Ni, W., R. N. Trelease, et al. (1990). Two temporally synthesized charge subunits interact to form the five isoforms of cottonseed (*Gossypium hirsutum*) catalase. *Biochemical Journal* 269(1): 233-238.

Noe, J. P. (1990). Efficacy of fumigant nematicides to control *Hoplolaimus columbus* on cotton. *Journal Of Nematology* 22(4 Suppl): 718-723.

Norman, M. A., R. A. Liebl, et al. (1990). Site of clomazone action in tolerant soybean and susceptible cotton photomixotrophic cell suspension cultures. *Plant Physiology* 94(2): 704-709.

Norman, M. A., R. A. Liebl, et al. (1990). Uptake and metabolism of clomazone in tolerant-soybean and susceptible-cotton photomixotrophic cell suspension cultures. *Plant Physiology* 92(3): 777-784.

Palumbo, J. C., T. F. Watson, et al. (1990). Globemallow, *Sphaeralcea* spp., as reproductive hosts for the boll weevil (Coleoptera: Curculionidae) in Arizona (USA). *Journal Of Economic Entomology* 83(2): 392-397.

Pan, X., M. Liu, et al. (1990). Meteorological environmental and genotypical controls of cotton photosynthesis. *Acta Agronomica Sinica* 16(4): 317-323.

Panchabhavi, K. S., K. A. Kulkarni, et al. (1990). Comparative efficiency of techniques for assessing loss due to insect pests in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 60(4): 252-254.

- Patterson, M. G., C. D. Monks, et al. (1990). Effects of chlorimuron applied postemergence to cotton (*Gossypium hirsutum*). *Weed Technology* 4(2): 314-317.
- Pettigrew, W. T., J. D. Hesketh, et al. (1990). Carbon dioxide saturated photosynthesis as affected by phosphate stress. *Photosynthetica* 24(2): 209-216.
- Pettigrew, W. T., J. D. Hesketh, et al. (1990). A vapor pressure deficit effect on crop canopy photosynthesis. *Photosynthesis Research* 24(1): 27-34.
- Pike, C. S., H. A. Norman, et al. (1990). Effects of acclimation to low temperature and to water stress on photosynthesis and on physical and chemical properties of lipids from thylakoids of cucumber and cotton. *Plant Science* 68(2): 189-196.
- Price, H. J., D. M. Stelly, et al. (1990). Molecular cytogenetic mapping of a nucleolar organizer region in cotton. *Journal Of Heredity* 81(5): 365-370.
- Radin, J. W. (1990). Responses of transpiration and hydraulic conductance to root temperature in nitrogen- and phosphorus-deficient cotton seedlings. *Plant Physiology* 92(3): 855-857.
- Raj, M. F., P. G. Shah, et al. (1990). Residues of synthetic pyrethroids in cotton seed, oil and lint. *Indian Journal Of Plant Protection* 18(2): 191-196.
- Ramsden, L. and H. Meier (1990). Phosphorylation of proteins in plasma membrane fractions from developing cotton fibers. *Journal Of Plant Physiology* 136(3): 313-317.
- Rao, N. V., A. S. Reddy, et al. (1990). Incidence of whitefly (*Bemisia tabaci*) in relation to leaf characters of upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 60(9): 619-624.
- Rao, T. N., Y. S. Nerkar, et al. (1990). Identification of cultivars of cotton by SDS polyacrylamide gel electrophoresis (SDS-PAGE) of soluble seed proteins. *Plant Varieties & Seeds* 3(1): 7-14.
- Rasulov, B. K., T. R. Pyarnik, et al. (1990). Carbon dioxide exchange components and photosynthetic carbon metabolism in cotton leaves under conditions of inhibited assimilate export. *Fiziologiya Rastenii* 37(1): 12-21.
- Rawitz, E., H. Lior, et al. (1990). The effect of drip-line placement and residue incorporation on the growth of drip-irrigated cotton. *Soil & Tillage Research* 16(1-2): 227-232.
- Reddy, V. R., D. N. Baker, et al. (1990). Temperature and mepiquat chloride effects on cotton canopy architecture. *Agronomy Journal* 82(2): 190-195.
- Riffle, M. S., D. S. Murray, et al. (1990). Soil-water relations and interference between devil's claw (*Proboscidea louisianica*) and cotton (*Gossypium hirsutum*). *Weed Science* 38(1): 39-44.
- Rojas, R. R., J. E. Leggett, et al. (1990). Dormancy in southwestern cotton boll weevils: A comparative physiological study of *Anthonomus grandis* in wild and domesticated cotton bolls. *Journal Of Insect Physiology* 36(8): 601-606.
- Sanam'Yan, M. F. and D. A. Musaev (1990). Detection and cytological study of the aneuploid and euploid plants with translocations of chromosomes in cotton *Gossypium hirsutum* L. *Genetika* 26(3): 506-514.
- Schmitt, D. P. and J. E. Bailey (1990). Chemical control of *Hoplolaimus columbus* on cotton and soybean. *Journal Of Nematology* 22(4 Suppl): 689-694.
- Singh, D. and R. K. Sahay (1990). Proline accumulation in relation to yield reduction and biomass recovery under stress conditions in cotton (*Gossypium* spp.). *Indian Journal Of Agricultural Sciences* 60(11): 739-741.

- Singh, J., Z. S. Dhaliwal, et al. (1990). Temporal changes in the dispersion of populations of three homopterous insect pests of upland cotton. *Insect Science And Its Application* 11(1): 73-78.
- Singh, M., T. H. Singh, et al. (1990). Genetic analysis of lint yield and its components in cotton. *Crop Improvement* 17(1): 64-67.
- Singh, M., V. P. Singh, et al. (1990). Development of high-yielding and better-quality varieties of upland cotton (*Gossypium hirsutum*) resistant to bacterial blight (*Xanthomonas campestris* pathovar *malvacearum*). *Indian Journal Of Agricultural Sciences* 60(7): 439-442.
- Singh, P. and S. S. Narayanan (1990). Intraspecific heterosis in diploid and tetraploid cottons. *Indian Journal Of Genetics & Plant Breeding* 50(4): 396-399.
- Singh, T. and L. S. Brar (1990). Efficacy of different herbicides for weed control in American cotton. *Journal Of Research Punjab Agricultural University* 27(4): 562-566.
- Smith, B. S., D. S. Murray, et al. (1990). Velvetleaf (*Abutilon theophrasti*) interference with cotton (*Gossypium hirsutum*). *Weed Technology* 4(4): 799-803.
- Smith, B. S., J. A. Pawlak, et al. (1990). Interference from established stands of silverleaf nightshade (*Solanum elaeagnifolium*) on cotton (*Gossypium hirsutum*) lint yield. *Weed Science* 38(2): 129-133.
- Snipes, C. E., M. E. Kurtz, et al. (1990). Productivity of cotton following rice. *Journal Of Production Agriculture* 3(2): 209-211.
- Sobreira, D. G., J. D. C. Machado, et al. (1990). Occurrence of fungi and physiological conditions of some seed lots of cotton (*Gossypium hirsutum* L.) produced in the State of Minas Gerais, season 1985/86. *Ciencia E Pratica* 14(3): 317-325.
- Stelly, D. M. (1990). Localization of the Le-2 locus of cotton (*Gossypium hirsutum* L.). *Journal Of Heredity* 81(3): 193-197.
- Stelly, D. M., K. C. Kautz, et al. (1990). Pollen fertility of some simple and compound translocations of cotton. *Crop Science* 30(4): 952-955.
- Stoilova, A. (1990). An express method for inducing mutations in remote cotton hybrids. *Genetika I Seleksiya* 23(6): 545-549.
- Subramanian, M. V., H. Y. Hung, et al. (1990). Properties of mutant acetolactate synthases resistant to triazolopyrimidine sulfonanilide. *Plant Physiology* 94(1): 239-244.
- Sun, X., I. Rao, et al. (1990). Potassium nutrition of cotton plant in relation to potassium status in soils. *Acta Pedologica Sinica* 27(2): 166-171.
- Tan, L. and Z. Liu (1990). Researches on selection and varietal traits of Zhongmian 12. *Scientia Agricultura Sinica* 23(3): 12-19.
- Temple, P. J. (1990). Water relations of differentially irrigated cotton exposed to ozone. *Agronomy Journal* 82(4): 800-805.
- Tianzheng, Z. and P. Jiaju (1990). A male -sterile line with "virescent" marker character in Upland cotton. *Euphytica* 48(3): 233-238.
- Turley, R. B. and R. N. Trelease (1990). Purification of plant peroxisomes in iso-osmotic metrizamide. *Physiologia Plantarum* 79(3): 570-578.
- Turley, R. B. and R. N. Trelease (1990). Development and regulation of three glyoxysomal enzymes during cotton seed maturation and growth. *Plant Molecular Biology* 14(2): 137-146.

- Uthamasamy, S., P. Sivasubramanian, et al. (1990). Monitoring of whitefly (*Bemisia tabaci*) on upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 60(11): 744-746.
- Van, H. H. G., L. J. B. Rossouw, et al. (1990). Yield and quality of cotton (*Gossypium hirsutum* L.) in a ratoon cropping system. *Applied Plant Science* 4(2): 67-69.
- Wali, B. M. and V. R. Koraddi (1990). Studies on crop geometry and plant density in rainfed cotton: II. Dry matter production and nutrient uptake. *Mysore Journal Of Agricultural Sciences* 24(2): 189-194.
- Wallace, T. P. and Z. K. M. El (1990). Quantitative analysis of resistance in cotton to three new isolates of the bacterial blight pathogen. *Theoretical And Applied Genetics* 79(4): 443-448.
- Wang, K. and M. Li (1990). The karyotype variation and evolution of D genome in *Gossypium*. *Acta Agronomica Sinica* 16(3): 200-207.
- Wendel, J. F. and R. G. Percy (1990). Allozyme diversity and introgression in the Galapagos Islands (Pacific Ocean) endemic *Gossypium darwinii* and its relationship to continental *Gossypium barbadense*. *Biochemical Systematics And Ecology* 18(7-8): 517-528.
- Westerman, R. B. and D. S. Murray (1990). Cotton (*Gossypium hirsutum*) response to glyphosate spot applied. *Weed Technology* 4(4): 759-764.
- White, R. H. and A. D. Worsham (1990). Control of legume cover crops in no-till corn (*Zea mays*) and cotton (*Gossypium hirsutum*). *Weed Technology* 4(1): 57-62.
- Wiegand, C. L. and A. J. Richardson (1990). Use of spectral vegetation indices to infer leaf area, evapotranspiration and yield: II. Results. *Agronomy Journal* 82(3): 630-636.
- Wiesenborn, W. D. and T. C. Baker (1990). Upwind flight to cotton flowers by *Pectinophora gossypiella* (Lepidoptera: Gelechiidae). *Environmental Entomology* 19(3): 490-493.
- Wilkinson, M. C., P. A. Wheatley, et al. (1990). Higher plant heat-shock protein 70: Purification and immunochemical analysis. *Phytochemistry* 29(10): 3073-3080.
- Wilson, F. D. (1990). Relative resistance of cotton lines to pink bollworm. *Crop Science* 30(3): 500-504.
- Wullschleger, S. D. and D. M. Oosterhuis (1990). Photosynthetic and respiratory activity of fruiting forms within the cotton canopy. *Plant Physiology* 94(2): 463-469.
- Wullschleger, S. D. and D. M. Oosterhuis (1990). Photosynthesis of individual field-grown cotton leaves during ontogeny. *Photosynthesis Research* 23(2): 163-170.
- Yakir, D., M. J. Deniro, et al. (1990). Effects of water stress on oxygen, hydrogen and carbon isotope ratios in two species of cotton plants. *Plant Cell And Environment* 13(9): 949-956.
- Yakir, D., M. J. Deniro, et al. (1990). Natural deuterium and oxygen-18 enrichment in leaf water of cotton plants grown under wet and dry conditions: Evidence for water compartmentation and its dynamics. *Plant Cell And Environment* 13(1): 49-56.
- Yakubova, M. M., K. I. Bobodzhanova, et al. (1990). The functioning of supracomplexes from *Gossypium* and *Arabidopsis* chloroplasts. *Biologicheskie Nauki*(1): 36-41.
- Yu, S. and Z. Huang (1990). Inheritance analysis of earliness components of short season cotton varieties in *Gossypium hirsutum*. *Scientia Agricultura Sinica* 23(6): 48-54.
- Yuan, C. X. and J. Ding (1990). Effects of water stress on the content of IAA and the activities of IAA oxidase and peroxidase in cotton leaves. *Acta Phytophysiological Sinica* 16(2): 179-180.

- Zhang, S., J. T. Cothren, et al. (1990). Mepiquat chloride seed treatment and germination temperature effects on cotton growth, nutrient partitioning, and water use efficiency. *Journal Of Plant Growth Regulation* 9(4): 195-200.
- Zhou, N. and Y. Xi (1990). Peroxidase isozyme study on the origin of tetraploid species in *Gossypium*. *Acta Genetica Sinica* 17(4): 294-300.
- Zodape, S. T. and S. D. Gomkale (1990). Method of increasing cotton yield using pulsed concentrated solar radiation. *Indian Journal Of Plant Physiology* 33(2): 141-144.
- Zubkova, N. F., Z. V. Bukashkina, et al. (1990). Study of the defoliating effects of cytokinins. *Fiziologiya Rastenii* 37(2): 535-541.
- Akey, D. H. and G. D. J. Butler (1989). Developmental rates and fecundity of apterous *Aphis gossypii* on seedlings of *Gossypium hirsutum*. *Southwestern Entomologist* 14(3): 295-301.
- Alabi, S. O. and C. A. Echekwu (1989). Genotype X environment interactions in cotton and their implications for future variety testing in the northern cotton-growing zone of Nigeria. *Samaru Journal Of Agricultural Research* 6(June): 44-47.
- Ali, A., R. G. Luttrell, et al. (1989). Distribution of fall armyworm (Lepidoptera: Noctuidae) egg masses on cotton. *Environmental Entomology* 18(5): 881-885.
- Ansari, A. H., A. M. Khushk, et al. (1989). Effect of different planting dates on the growth and yield of cotton (*Gossypium hirsutum* L.) cultivars. *Pakistan Journal Of Scientific And Industrial Research* 32(7): 474-477.
- Aripdzhanov, S. A., K. A. Khakimov, et al. (1989). Biosynthesis of nucleic acids and proteins in cotton cells during callusogenesis phase. *Biopolimery I Kletka* 5(6): 104-107.
- Baehrecke, E. H., H. J. Williams, et al. (1989). Electroantennogram responses of *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) to chemicals in cotton (*Gossypium hirsutum* L.). *Journal Of Chemical Ecology* 15(1): 37-46.
- Barnes, C. J., A. J. Goetz, et al. (1989). Effects of imazaquin residues on cotton (*Gossypium hirsutum*). *Weed Science* 37(6): 820-824.
- Ben, H. M., Z. Plaut, et al. (1989). Cotton canopy and drying effects on runoff during irrigation with moving sprinkler systems. *Agronomy Journal* 81(5): 752-757.
- Bergey, D. R., D. M. Stelly, et al. (1989). In situ hybridization of biotinylated DNA probes to cotton meiotic chromosomes. *Stain Technology* 64(1): 25-38.
- Beyrouthy, C. A. and D. M. Oosterhuis (1989). The porous-membrane technique for root studies of field-grown crops. *Plant And Soil* 116(2): 265-272.
- Bhale, N. L., R. G. Dani, et al. (1989). Variation among plants for seed-oil and lint characteristics of cotton (*Gossypium hirsutum* L.). *Indian Journal Of Genetics & Plant Breeding* 49(2): 265-268.
- Bilger, W., O. Bjorkman, et al. (1989). Light-induced spectral absorbance changes in relation to photosynthesis and the epoxidation state of xanthophyll cycle components in cotton leaves. *Plant Physiology* 91(2): 542-551.
- Botha, J. H., A. I. Greeff, et al. (1989). Preliminary screening of cotton plants for resistance to spider mite damage in South Africa. *Phytophylactica* 21(4): 379-384.
- Bradburne, J. A., M. J. Kasperbauer, et al. (1989). Reflected far-red light effects on chlorophyll and light-harvesting chlorophyll protein (LHC-II) contents under field conditions. *Plant Physiology* 91(3): 800-803.

Brody, A. K. and R. Karban (1989). Demographic analysis of induced resistance against spider mites (Acari: Tetranychidae) in cotton. *Journal Of Economic Entomology* 82(2): 462-465.

Buchala, A. J., S. Roulin, et al. (1989). Polysaccharides in the culture medium of cotton *Gossypium hirsutum* L. ovules cultured in vitro. *Plant Cell Reports* 8(1): 25-28.

Buriro, R. S., S. M. Qayyum, et al. (1989). Inter and intra row spacings effects on the growth and yield of cotton (*Gossypium hirsutum* L.). *Sarhad Journal Of Agriculture* 5(5): 473-478.

Burke, J. J. and D. R. Upchurch (1989). Leaf temperature and transpirational control in cotton. *Environmental And Experimental Botany* 29(4): 487-492.

Butler, G. D. J., D. L. Coudriet, et al. (1989). Sweet-potato whitefly: Host plant preference and repellent effect of plant-derived oils on cotton, squash, lettuce and cantaloupe. *Southwestern Entomologist* 14(1): 9-16.

Carvalho, L. P. D. and J. D. A. N. Moreira (1989). A selection criterion for earliness in herbaceous cotton. *Pesquisa Agropecuaria Brasileira* 24(7): 829-834.

Cassman, K. G., T. A. Kerby, et al. (1989). Differential response of two cotton cultivars to fertilizer and soil potassium. *Agronomy Journal* 81(6): 870-876.

Cassman, K. G., B. A. Roberts, et al. (1989). Soil potassium balance and cumulative cotton response to annual potassium additions on a vermiculitic soil. *Soil Science Society Of America Journal* 53(3): 805-812.

Castner, E. P., D. S. Murray, et al. (1989). Interference of hogpotato (*Hoffmanseggia glauca*) with cotton (*Gossypium hirsutum*). *Weed Science* 37(5): 688-694.

Chapman, K. D., R. B. Turley, et al. (1989). Relationship between cottonseed malate synthase aggregation behavior and suborganellar location in glyoxysomes and endoplasmic reticulum. *Plant Physiology* 89(1): 352-359.

Chen, Z. X., S. J. Li, et al. (1989). Plantlet regeneration from protoplasts isolated from an embryogenic suspension culture of cotton (*Gossypium hirsutum* L.). *Acta Botanica Sinica* 31(12): 966-969.

Chu, C. C., L. E. Graham, et al. (1989). Nitrate reductase activity and nitrate concentration in cotton plant leaf blades and petioles. *Agronomy Journal* 81(4): 577-581.

Cotty, P. J. (1989). Effects of cultivar and boll age on aflatoxin in cottonseed after inoculation with *Aspergillus flavus* at stimulated exit holes of the pink bollworm. *Plant Disease* 73(6): 489-492.

Croxford, A. C., P. J. Edwards, et al. (1989). Temporal and spatial variation in palatability of soybean and cotton leaves following wounding. *Oecologia* 79(4): 520-525.

Dani, R. G. (1989). Heterosis and combining ability for oil content and other economic traits in cotton (*Gossypium hirsutum* L.). *Indian Journal Of Genetics & Plant Breeding* 49(1): 47-52.

Dani, R. G. (1989). Genotype X environment interactions for seed-oil and protein content in cotton (*Gossypium hirsutum* L.). *Indian Journal Of Genetics & Plant Breeding* 49(2): 237-240.

Dani, R. G. and R. J. Kohel (1989). Maternal effects and generation mean analysis of seed-oil content in cotton (*Gossypium hirsutum* L.). *Theoretical And Applied Genetics* 77(4): 569-575.

Davidonis, G. H. (1989). Fiber development in preanthesis cotton ovules. *Physiologia Plantarum* 75(2): 290-294.

- De, B. L. P., K. J. De, et al. (1989). The effect of plant density on yield and quality of cotton (*Gossypium hirsutum* L.) in the Vaalharts and Sandvet irrigation areas (South Africa). *South African Journal Of Plant And Soil* 6(3): 154-160.
- De, K. J., J. J. Human, et al. (1989). The effect of delaying irrigation on yield and quality of cotton (*Gossypium hirsutum* L.) in the Vaalharts irrigation area (South Africa). *South African Journal Of Plant And Soil* 6(3): 179-182.
- De, P. F. M. D., L. G. R. Ferreira, et al. (1989). Physiological and biochemical changes of two cotton cultivars (*Gossypium hirsutum* L.) under water stress. *Revista Brasileira De Botanica* 12(1-2): 1-10.
- Degrandi, H. G. and F. Morales (1989). Identification and distribution of pollinating honey bees (Hymenoptera: Apidae) on sterile male cotton. *Journal Of Economic Entomology* 82(2): 580-583.
- Demmig, A. B., K. Winter, et al. (1989). Photosynthetic characteristics and the ratios of chlorophyll, beta-carotene, and the components of the xanthophyll cycle upon a sudden increase in growth light regime in several plant species. *Botanica Acta* 102(4): 319-325.
- Dhandapani, N., A. A. Kareem, et al. (1989). Life table studies of *Spodoptera litura* (F.) on two varieties of cotton. *Journal Of Entomological Research* 13(1-2): 60-63.
- Dhawan, A. K., G. S. Simwat, et al. (1989). Effect of late sowing of different varieties of upland cotton on bollworm incidence and yield of seed cotton. *Journal Of Research Punjab Agricultural University* 26(3): 423-430.
- Dickens, J. C. (1989). Green leaf volatiles enhance aggregation pheromone of boll weevil, *Anthonomus grandis*. *Entomologia Experimentalis Et Applicata* 52(3): 191-204.
- Diubate, M. and K. S. Karev (1989). Effect of fertilization and plant density on the yield and quality of Beli izvor cotton cultivar. *Pochvoznanie I Agrokimiya* 24(6): 40-45.
- Dos, S. W. J., A. P. Gutierrez, et al. (1989). Economic damage caused by the cotton stem borer in southern Brazil. *Pesquisa Agropecuaria Brasileira* 24(3): 297-306.
- Dugas, W. A. (1989). Plant spacing effects on net irradiance extinction in cotton. *Theoretical And Applied Climatology* 40(1-2): 93-100.
- Durant, J. A. (1989). Yield response of cotton cultivars to early-season applications of chlordimeform and aldicarb. *Journal Of Economic Entomology* 82(2): 626-632.
- El, H. L., T. P. T. Anh, et al. (1989). Enzymatic breakdown of polar lipids in cotton leaves under water stress: I. Degradation of monogalactosyl-diacylglycerol. *Plant Physiology And Biochemistry* 27(4): 495-502.
- Endrizzi, J. E., F. R. H. Katterman, et al. (1989). DNA hybridization and time of origin of three species of *Gossypium*. *Evolutionary Trends In Plants* 3(2): 115-120.
- Ephrath, J. E., D. Shiteinberg, et al. (1989). *Alternaria alternata* in cotton (*Gossypium hirsutum*) cultivar Acala: Effects on gas exchange, yield components and yield accumulation. *Netherlands Journal Of Plant Pathology* 95(3): 157-166.
- Fender, S. E. and M. A. O'Connell (1989). Heat shock protein expression in thermotolerant and thermosensitive lines of cotton. *Plant Cell Reports* 8(1): 37-40.
- Fleischer, S. H., M. J. Gaylor, et al. (1989). Roadside management of annual fleabane (*Erigeron annuus*) and wild carrot (*Daucus carota*). *Weed Technology* 3(1): 72-75.

- Flint, H. M. and N. J. Parks (1989). Effect of azadirachtin from the neem tree on immature sweetpotato whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae) and other selected pest species on cotton. *Journal Of Agricultural Entomology* 6(4): 211-216.
- Flint, H. M., F. D. Wilson, et al. (1989). Causes of square shed in cotton in central Arizona (USA). *Southwestern Entomologist* 14(3): 271-278.
- Francey, Y., J. P. Jaquet, et al. (1989). The biosynthesis of beta-glucans in cotton (*Gossypium hirsutum* L.) fibers of ovules cultured in vitro. *Journal Of Plant Physiology* 134(4): 485-491.
- Furbeck, S. M., F. M. Bourland, et al. (1989). Comparison of the hot water and accelerated ageing techniques for deterioration of cottonseed. *Seed Science And Technology* 17(2): 255-262.
- Galau, G. A. and T. A. Wilkins (1989). Alloplasmic male sterility in AD allotetraploid *Gossypium hirsutum* upon replacement of its resident A cytoplasm with that of D species *Gossypium harknessii*. *Theoretical And Applied Genetics* 78(1): 23-30.
- Geever, R. F., F. R. H. Katterman, et al. (1989). DNA hybridization analyses of a *Gossypium* allotetraploid and two closely related diploid species. *Theoretical And Applied Genetics* 77(4): 553-559.
- Gill, S. S., T. H. Singh, et al. (1989). Variability in morphological and yield component traits of upland cotton (*Gossypium hirsutum* L.) under different environments. *Journal Of Research Punjab Agricultural University* 26(2): 175-179.
- Giller, Y. E. (1989). Variability of the molecular organization of the pigment system photosynthetic apparatus. *Biologicheskii Zhurnal Armenii* 42(9-10): 890-899.
- Goodell, P. B. and H. Ferris (1989). Influence of environmental factors on the hatch and survival of *Meloidogyne incognita*. *Journal Of Nematology* 21(3): 328-334.
- Guinn, G. and D. L. Brummett (1989). Fruiting of cotton: IV. Nitrogen, abscisic acid, indole-3-acetic acid, and cutout. *Field Crops Research* 22(4): 257-266.
- Guinn, G. and D. L. Brummett (1989). Changes in amide-linked and ester IAA in cotton fruiting forms during their development. *Plant Physiology* 89(3): 941-944.
- Gulick, S. H., K. G. Cassman, et al. (1989). Exploitation of soil potassium in layered profiles by root systems of cotton and barley. *Soil Science Society Of America Journal* 53(1): 146-153.
- Gulin, V. V., K. Y. Saidov, et al. (1989). Inheritance of benzidinase in the cotton *Gossypium hirsutum*. *Doklady Akademii Nauk Sssr* 307(1): 223-225.
- Guthrie, D. S. and J. T. Cothren (1989). Enhanced ethylene evolution and electrolyte leakage from methomyl-treated cotton (*Gossypium hirsutum*). *Field Crops Research* 19(4): 241-252.
- Guthrie, D. S. and A. C. York (1989). Cotton (*Gossypium hirsutum*) development and yield following fluometuron postemergence applied. *Weed Technology* 3(3): 501-504.
- Harman, W. L., G. J. Michels, et al. (1989). A conservation tillage system for profitable cotton production in the central Texas (USA) high plains. *Agronomy Journal* 81(4): 615-618.
- Hasnam and E. Sulistyowati (1989). Performance of cotton varieties under intercropping with mungbean. *Indonesian Journal Of Crop Science* 4(1): 15-26.
- Hatfield, J. L. (1989). Aerodynamic properties of partial canopies. *Agricultural And Forest Meteorology* 46(1-2): 15-22.

- Henneberry, T. J. and T. E. Clayton (1989). Oviposition patterns, percent egg hatch, and larval mortality of pink bollworm (Lepidoptera: Gelechiidae) on cotton plants. *Journal Of Economic Entomology* 82(3): 932-934.
- Hofmann, W. C. and P. T. Else (1989). The response of cotton to foliar applications of a cytokinin-containing compound. *Applied Agricultural Research* 4(1): 25-29.
- Hossain, S., I. H. Mian, et al. (1989). Efficacy of three nematicides and two oil cakes for control of root-knot nematode (*Meloidogyne incognita*) on potato seedlings. *Journal Of The Faculty Of Agriculture Kyushu University* 34(1-2): 115-121.
- Huang, G., Y. Gou, et al. (1989). Studies on a nuclear male sterility maintainer line of cotton. *Scientia Agricultura Sinica* 22(6): 13-17.
- Hughes, D. W. and G. A. Galau (1989). Temporally modular gene expression during cotyledon development. *Genes & Development* 3(3): 358-369.
- Hussain, R. R., T. M. Piracha, et al. (1989). Effect of pre- and post-emergence application of herbicides on weed growth and seed cotton yield of American cotton (*Gossypium hirsutum* L.) cultivar B557. *Pakistan Journal Of Scientific And Industrial Research* 32(5): 336-338.
- Idso, S. B., S. G. Allen, et al. (1989). Problems with porometry: Measuring net photosynthesis by leaf chamber techniques. *Agronomy Journal* 81(3): 475-479.
- Ilango, K. and S. Uthamasamy (1989). Influence of spacings and fertilizer levels on the incidence of bollworm-boll rot complex in three varieties of cotton (*Gossypium hirsutum*). *Applied Agricultural Research* 4(3): 173-178.
- Kadapa, S. N. (1989). Development of cotton (*Gossypium hirsutum* L.) variety JK 119 producing higher yield than DCH 32 hybrid. *Indian Journal Of Genetics & Plant Breeding* 49(3): 413-420.
- Kadapa, S. N. (1989). Development of compact cotton strains for high lint productivity in *Gossypium hirsutum* L.: I. Breeding procedure and selection. *Indian Journal Of Genetics & Plant Breeding* 49(3): 313-320.
- Karimov, K. K. and S. V. Dontsova (1989). Synthesis of nucleic acids in germinating cotton seeds as affected by abscisic acid. *Fiziologiya Rastenii* 36(2): 332-338.
- Katageri, I. S. and S. N. Kadapa (1989). Heterosis for yield and its component characters in bollworm tolerant *Gossypium hirsutum* L. X *Gossypium barbadense* L. cotton hybrids. *Indian Journal Of Genetics & Plant Breeding* 49(1): 107-112.
- Keeley, P. E. and R. J. Thullen (1989). Growth and interaction of johnsongrass (*Sorghum halepense*) with cotton (*Gossypium hirsutum*). *Weed Science* 37(3): 339-344.
- Keeley, P. E. and R. J. Thullen (1989). Growth and competition of black nightshade (*Solanum nigrum*) and Palmer amaranth (*Amaranthus palmeri*) with cotton (*Gossypium hirsutum*). *Weed Science* 37(3): 326-334.
- Keeling, J. W. and J. R. Abernathy (1989). Response of cotton (*Gossypium hirsutum*) to repeated applications of dinitroaniline herbicides. *Weed Technology* 3(3): 527-530.
- Keeling, J. W. and J. R. Abernathy (1989). Preemergence weed control in a conservation tillage cotton (*Gossypium hirsutum*) cropping system on sandy soils. *Weed Technology* 3(1): 182-185.
- Keeling, J. W., C. G. Henniger, et al. (1989). Horseweed (*Conyza canadensis*) control in conservation tillage cotton (*Gossypium hirsutum*). *Weed Technology* 3(2): 399-401.

- Keeling, W., E. Segarra, et al. (1989). Evaluation of conservation tillage cropping systems for cotton on the Texas (USA) southern high plains. *Journal Of Production Agriculture* 2(3): 269-273.
- Kerby, T. A., M. Keeley, et al. (1989). Weather and seed quality variables to predict cotton seedling emergence. *Agronomy Journal* 81(3): 415-419.
- Khan, A. A., M. M. Ali, et al. (1989). Manifestation of heterosis and combining ability in upland cotton. *Pakistan Journal Of Scientific And Industrial Research* 32(9): 629-632.
- Kloth, R. H. (1989). Changes in the level of tubulin subunits during development of cotton (*Gossypium hirsutum*) fiber. *Physiologia Plantarum* 76(1): 37-41.
- Kohel, R. J. and A. E. Percival (1989). Genetic analysis of the rex mutant in cotton. *Journal Of Heredity* 80(1): 78-80.
- Lakshmanan, P. and S. Mohan (1989). Twig and stem blight of cotton caused by *Botryodiplodia theobromae*. *Plant Pathology* 38(2): 290-292.
- Lamoreaux, R. J., V. L. Corbin, et al. (1989). Adsorption and deactivation of norflurazon by activated charcoal. *Weed Technology* 3(2): 297-302.
- Li, Y. C., A. K. Alva, et al. (1989). Response of cotton cultivars to aluminum in solutions with varying silicon concentrations. *Journal Of Plant Nutrition* 12(7): 881-892.
- Luckett, D. J. (1989). Colchicine mutagenesis is associated with substantial heritable variation in cotton. *Euphytica* 42(1-2): 177-182.
- Mahler, R. J. (1989). Sulfur effects on cotton cultivars grown in a greenhouse. *Journal Of Plant Nutrition* 12(2): 187-206.
- McCarty, J. C. J. and J. E. Jones (1989). Boll weevil (Coleoptera: Curculionidae) nonpreference for primitive cotton. *Journal Of Economic Entomology* 82(1): 298-300.
- McConnell, J. S., B. S. Frizzell, et al. (1989). Effects of soil compaction and subsoil tillage of two alfisols on the growth and yield of cotton. *Journal Of Production Agriculture* 2(2): 140-146.
- McDowell, L. L., G. H. Willis, et al. (1989). Nitrogen and phosphorus yields in run-off from silty soils in the Mississippi Delta, USA. *Agriculture Ecosystems & Environment* 25(2-3): 119-138.
- Mellon, J. E. and B. A. Triplett (1989). De novo synthesis of peroxidase in cotton ovule culture medium. *Physiologia Plantarum* 77(3): 302-307.
- Meredith, W. R. J. and R. Wells (1989). Potential for increasing cotton yields through enhanced partitioning to reproductive structures. *Crop Science* 29(3): 636-639.
- Meron, M., D. W. Grimes, et al. (1989). Shoot resistance to water flow in cotton. *Journal Of Experimental Botany* 40(217): 919-924.
- Mert, H. H. (1989). Photosynthesis and photorespiration in two cultivars of cotton under salt stress. *Biologia Plantarum Prague* 31(5): 413-414. Biol. Dep., Fac. Educ., Dokuz Eylul Univ., Buca, Izmir, Turkey
- Milam, M. R., J. N. Jenkins, et al. (1989). Breeding upland cotton for resistance to the tarnished plant bug. *Field Crops Research* 21(3-4): 227-238.
- Miller, J. E., R. P. Patterson, et al. (1989). Response of soluble sugars and starch in field-grown cotton to ozone, water stress, and their combination. *Environmental And Experimental Botany* 29(4): 477-486.

Mitra, J. and K. Raghu (1989). Effects of DDT on the growth of crop plants. *Environmental Pollution* 61(2): 157-170.

Myles, E. L. J. and J. E. Endrizzi (1989). Aneuploids induced by deficiencies of chromosome 9 and analysis of the time of nondisjunction in cotton. *Genome* 32(1): 12-18.

Nanthagopal, R. and S. Uthamasamy (1989). Life tables for American bollworm, *Heliothis armigera* Hubner on four species of cotton under field conditions. *Insect Science And Its Application* 10(4): 521-530.

Nanthagopal, R. and S. Uthamasamy (1989). Life tables for spotted bollworm, *Earias vitella* (Fabricius), on four species of cotton. *Crop Protection* 8(2): 133-136.

Navasero, R. C. and G. W. Elzen (1989). Responses of *Microplitis croceipes* to host and nonhost plants of *Heliothis virescens* in a wind tunnel. *Entomologia Experimentalis Et Applicata* 53(1): 57-64.

Nayyar, H., K. Kaur, et al. (1989). Regulation of differential fiber development in cotton by endogenous plant growth regulators. *Proceedings Of The Indian National Science Academy Part B Biological Sciences* 55(5-6): 463-468.

Nie, R. Z. and M. X. Li (1989). Cytological studies on semi-wild races of *Gossypium hirsutum*. *Acta Botanica Sinica* 31(5): 333-342.

Pandey, A., S. T. Shende, et al. (1989). Effect of *Azotobacter chroococcum* seed inoculation on its establishment in rhizosphere, on growth and yield and yield attributing parameters of cotton (*Gossypium hirsutum*). *Zentralblatt Fuer Mikrobiologie* 144(8): 595-604.

Pandey, A., S. T. Shende, et al. (1989). *Azotobacter chroococcum* seed inoculation for cotton (*Gossypium hirsutum*): Effect on seed quality. *Zentralblatt Fuer Mikrobiologie* 144(8): 605-610.

Parrott, W. L., J. N. Jenkins, et al. (1989). Relationship between gossypol gland density on cotton squares and resistance to tobacco budworm (Lepidoptera: Noctuidae) larvae. *Journal Of Economic Entomology* 82(2): 589-592.

Parrott, W. L., G. H. McKibben, et al. (1989). Feeding response of the boll weevil (Coleoptera: Curculionidae) to ester extracts of host plants. *Journal Of Economic Entomology* 82(2): 449-453.

Patterson, D. T. and M. T. Highsmith (1989). Competition of spurred anoda (*Anoda cristata*) and velvetleaf (*Abutilon theophrasti*) with cotton (*Gossypium hirsutum*) during simulated drought and recovery. *Weed Science* 37(5): 658-664.

Peng, Y. J. and S. Qian (1989). Embryological studies on interspecific hybrids of *Gossypium hirsutum* X *Gossypium raimondii* Ulbr. *Acta Agronomica Sinica* 15(3): 243-250.

Pennington, D. A. and L. Heatherley (1989). Effect of changing solar radiation on canopy-air temperatures of cotton and soybean. *Agricultural And Forest Meteorology* 46(1-2): 1-14.

Pizzamiglio, M. A., A. P. Gutierrez, et al. (1989). Phenological patterns and sampling decision rules for arthropods fields in Parana, Brazil: Before boll weevil. *Pesquisa Agropecuaria Brasileira* 24(3): 337-346.

Price, N. S., R. W. Roncadori, et al. (1989). Cotton root growth as influenced by phosphorus nutrition and vesicular-arbuscular mycorrhizas. *New Phytologist* 111(1): 61-66.

Puech, S. I., T. C. Hsiao, et al. (1989). Water-stress effects on the carbon exchange rates of three upland cotton (*Gossypium hirsutum*) cultivars in the field. *Field Crops Research* 21(3-4): 239-256.

Radin, J. W. and M. A. Matthews (1989). Water transport properties of cortical cells in roots of nitrogen-and phosphorus-deficient cotton seedlings. *Plant Physiology* 89(1): 264-268.

- Rahman, S., M. Ahmad, et al. (1989). Genetic analysis of seed oil and protein in various crosses of *Gossypium hirsutum* L. *Sarhad Journal Of Agriculture* 5(4): 387-394.
- Rakhimov, M. M., R. A. Akhmedzhanov, et al. (1989). Hydrolytic and transalkylating functions of phospholipase D from cotton seeds. *Fiziologiya Rastenii* 36(3): 502-511.
- Ramalho, F. S. and F. M. M. Jesus (1989). Evaluation of electrodynamic and conventional insecticides against cotton boll weevil and pink bollworm. *International Pest Control* 31(3): 56-58.
- Ramalho, F. S., V. L. B. Silva, et al. (1989). Effects of insecticide residues on *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae). *Pesquis a Agropecuaria Brasileira* 24(3): 315-320.
- Rashid, A., A. Mahmood, et al. (1989). Heterosis and heterobeltiosis studies for quality characters in intraspecific crosses of *Gossypium hirsutum* L. *Pakistan Journal Of Scientific And Industrial Research* 32(5): 333-335.
- Reddy, K. C. K., G. R. M. Sankar, et al. (1989). A basis for simultaneous optimization of chemical and organic nitrogen under optimum calcium:nitrogen conditions. *Indian Journal Of Agricultural Sciences* 59(2): 102-106.
- Reddy, V. R., B. Acock, et al. (1989). Seasonal leaf area-leaf weight relationships in the cotton canopy. *Agronomy Journal* 81(1): 1-4.
- Reddy, V. R., D. N. Baker, et al. (1989). Analysis of effects of atmospheric carbon dioxide and ozone on cotton yield trends. *Journal Of Environmental Quality* 18(4): 427-432.
- Riffle, M. S., D. S. Murray, et al. (1989). Duration and intensity of unicorn plant (*Proboscidea louisianica*) interference with cotton (*Gossypium hirsutum*). *Weed Technology* 3(2): 313-316.
- Rodriguez, K. R., D. G. Robertson, et al. (1989). Crops uncommon to Alabama (USA) for the management of *Meloidogyne arenaria* in peanut. *Journal Of Nematology* 21(4 Suppl): 712-716.
- Rooney, W. L. and D. M. Stelly (1989). Allelic composition of cotton at the Le-1 and Le-2 loci. *Crop Science* 29(3): 707-712.
- Rush, C. M. and T. J. Gerik (1989). Relationship between postharvest management of grain sorghum and phymatotrichum root rot in the subsequent cotton crop. *Plant Disease* 73(4): 304-305.
- Safarov, K. S., A. K. Kasymov, et al. (1989). A low-molecular-weight intracellular factor of organic nature from cotton seedling roots and its uncoupling effect in oxidative phosphorylation. *Fiziologiya Rastenii* 36(3): 512-519.
- Sappenfield, W. P. and J. N. Ward (1989). Selecting cotton cultivar types for southeast Missouri environments. *Journal Of Production Agriculture* 2(2): 161-164.
- Schaefer, C. and O. Bjoerkman (1989). Relationship between efficiency of photosynthetic energy conversion and chlorophyll fluorescence quenching in upland cotton (*Gossypium hirsutum* L.). *Planta* 178(3): 367-376.
- Seshadri, V. (1989). Effect of plant density and growth-regulator on growth and yield of two hybrids of cotton (*Gossypium hirsutum* X *Gossypium barbadense*). *Indian Journal Of Agricultural Sciences* 59(2): 107-109.
- Seshadri, V. and K. Natarajan (1989). Effect of different intercrops on growth and yield of upland cotton (*Gossypium hirsutum*) and their economics. *Indian Journal Of Agricultural Sciences* 59(4): 227-230.
- Seshadrinathan, A. R. and H. K. H. Rao (1989). Effect of soil amender Fertilade on growth, development and yield of upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 59(5): 312-316.

- Shepherd, R. L. and M. G. Huck (1989). Progression of root-knot nematode symptoms and infection on resistant and susceptible cottons. *Journal Of Nematology* 21(2): 235-241.
- Silva, L. C., J. T. A. D. Costa, et al. (1989). Stomatal density and distribution in leaves of *Gossypium* sp. *Pesquisa Agropecuaria Brasileira* 24(7): 835-840.
- Singh, D. and R. K. Sahay (1989). Effect of pix and kaolin on growth and yield of upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 59(4): 247-250.
- Singh, K. and S. Singh (1989). Different cotton (*Gossypium hirsutum*) based cropping systems, and their fertilizer-use efficiency, yield and economic returns. *Indian Journal Of Agricultural Sciences* 59(5): 286-290.
- Singh, M., V. P. Singh, et al. (1989). Improvement of yield through increased boll weight in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 59(3): 141-144.
- Singh, R. and A. D. Taneja (1989). Influence of phytochemicals and leaf pubescence of some malvaceous plants on development, survival and oviposition of cotton leafhopper. *Zeitschrift Fuer Angewandte Zoologie* 76(3): 357-368.
- Slosser, J. E., J. R. Price, et al. (1989). Evaluation of furrow diking and early -season insecticide applications on boll weevils (Coleoptera: Curculionidae), bollworms (Lepidoptera: Noctuidae), and cotton yield in the Texas Rolling Plains (USA). *Journal Of Economic Entomology* 82(2): 599-607.
- Smith, E. M. and F. C. Wehner (1989). In vivo and in vitro effects of alternative crops on *Rhizoctonia solani* associated with crater disease of wheat. *Phytophylactica* 21(1): 61-64.
- Sneh, B., A. M. Ichielevich, et al. (1989). Mechanism of seedling protection induced by a hypovirulent isolate of *Rhizoctonia solani*. *Canadian Journal Of Botany* 67(7): 2135-2141.
- Sneh, B., A. M. Ichielevich, et al. (1989). Comparative anatomy of colonization of cotton hypocotyls and roots by virulent and hypovirulent isolates of *Rhizoctonia solani*. *Canadian Journal Of Botany* 67(7): 2142-2149.
- Sohi, H. S., K. S. Aulakh, et al. (1989). Prevalence of *Xanthomonas campestris* pathovar *malvacearum* in cotton seeds and its control by seed treatment. *Annals Of Biology* 5(1): 29-32.
- Sohu, V. S., G. S. Chahal, et al. (1989). Genetic analysis of bloom rate in upland cotton (*Gossypium hirsutum* L.). *Journal Of Genetics & Breeding* 43(4): 191-194.
- Sproos, B. B., J. Rotem, et al. (1989). The relationship between infections of the cotyledons of *Gossypium barbadense* and *Gossypium hirsutum* with *Alternaria macrospora* and cotyledon abscission. *Physiological And Molecular Plant Pathology* 35(4): 293-300.
- Steiner, J. L. (1989). Tillage and surface residue effects on evaporation from soils. *Soil Science Society Of America Journal* 53(3): 911-916.
- Stelly, D. M., D. W. Altman, et al. (1989). Cytogenetic abnormalities of cotton somaclones from callus cultures. *Genome* 32(5): 762-770.
- Stewart, S. D. and W. L. Sterling (1989). Susceptibility of cotton fruiting forms to insects, boll rot, and physical stress. *Journal Of Economic Entomology* 82(2): 593-598.
- Stoilova, A. and N. Savova (1989). Inheritance of fiber length in *Gossypium hirsutum* X *Gossypium barbadense* crosses. *Genetika I Seleksiya* 22(3): 198-204.
- Tanaka, M. A. S., J. O. M. Menten, et al. (1989). Artificial inoculation of cotton seeds with *Colletotrichum gossypii* var. *cephalosporioides* and seed infection as a function of exposure time to the pathogen. *Summa Phytopathologica* 15(3-4): 232-237.

- Thomson, N. J. and D. J. Luckett (1989). Heterosis and combining ability effects on cotton: I. Combining ability. *Australian Journal Of Agricultural Research* 39(6): 973-990.
- Tiwana, U. S., A. S. Brar, et al. (1989). Effect of sowing dates and nitrogen application on growth and yield of upland cotton (*Gossypium hirsutum* L.) in Punjab (India). *Journal Of Research Punjab Agricultural University* 26(3): 373-378.
- Tomar, S. P. S., S. S. Tomar, et al. (1989). Effect of soil and foliar application of nitrogen and phosphorus on yield of seed cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 59(8): 506-508.
- Trolinder, N. L. and C. Xhixian (1989). Genotype specificity of the somatic embryogenesis response in cotton. *Plant Cell Reports* 8(3): 133-136.
- Ullrich, E. C. I., A. Sanz, et al. (1989). Evaluation of arsenate- and vanadate-associated changes of electrical membrane potential and phosphate transport in *Lemna gibba* gl. *Journal Of Experimental Botany* 40(210): 119-128.
- Ull'Masov, T. N., M. K. Gulov, et al. (1989). Cloning of chloroplast *psbA* and *rbcL* genes from cotton *Gossypium hirsutum*. *Molekulyarnaya Genetika Mikrobiologiya I Virusologiya*(8): 17-20.
- Umbeck, P., W. Swain, et al. (1989). Inheritance and expression of genes for kanamycin and chloramphenicol resistance in transgenic cotton plants. *Crop Science* 29(1): 196-201.
- Ungar, E. D., E. Kletter, et al. (1989). Early season development of floral buds in cotton. *Agronomy Journal* 81(4): 643-649.
- Wali, B. M. and V. R. Koraddi (1989). Biometrical studies in rainfed cotton: I. Growth and yield. *Mysore Journal Of Agricultural Sciences* 23(4): 441-446.
- Wilson, F. D. (1989). Inheritance of undulate leaf mutant of upland cotton. *Journal Of Heredity* 80(6): 482-483.
- Wilson, F. D. (1989). Yield, earliness, and fiber properties of cotton carrying combined traits for pink bollworm resistance. *Crop Science* 29(1): 7-12.
- Wullschleger, S. D. and D. M. Oosterhuis (1989). Water use efficiency as a function of leaf age and position within the cotton canopy. *Plant And Soil* 120(1): 79-86.
- Wullschleger, S. D. and D. M. Oosterhuis (1989). The occurrence of an internal cuticle in cotton (*Gossypium hirsutum* L.) leaf stomates. *Environmental And Experimental Botany* 29(2): 229-236.
- Zhang, T., J. Pan, et al. (1989). Genetic identification of a genetic male-sterile line associated with virescent indicative character in upland cotton. *Scientia Agricultura Sinica* 22(4): 17-21.
- Ahmad, J. and S. A. Aziz (1988). Studies on the application of Dyar's law to the different nymphal instars of *Dysdercus cingulatus* Fab. *Indian Journal Of Entomology* 50(2): 220-222.
- Akey, A. D., G. D. Butler, et al. (1988). A plant root warmer for studying phytophagous insect development and fecundity at temperatures below host plant tolerance. *Annals Of The Entomological Society Of America* 81(6): 950-953.
- Akhtar, M. A. and I. U. Khan (1988). Bacterial blight of cotton and its control. *Pakistan Journal Of Agricultural Research* 9(2): 202-204.
- Akhtar, M. R., M. Iqbal, et al. (1988). Farmer's low and high yielding wheat technologies: Cotton zone of the Punjab (Pakistan). *Pakistan Journal Of Agricultural Research* 9(3): 321-324.

- Akhtar, M. S., A. U. H. Khan, et al. (1988). Effect of salinity-sodicity interaction on growth and leaf composition of cotton. *Pakistan Journal Of Agricultural Research* 9(4): 458-462.
- Alm, D. M., D. R. Pike, et al. (1988). Leaf area development in some crop and weed species. *Biotronics* 17: 29-40.
- Altman, D. W. (1988). Exogenous hormone applications at pollination for in vitro and in vivo production of cotton interspecific hybrids. *Plant Cell Reports* 7(4): 257-261.
- Ansari, A. H., A. M. Khuskh, et al. (1988). Effect of post emergence weedicides on the yield and yield component of cotton *Gossypium hirsutum* L. *Pakistan Journal Of Scientific And Industrial Research* 31(4): 285-287.
- Azevedo, D. M. P. D., L. B. D. Nobrega, et al. (1988). Selectivity and efficiency of herbicide mixtures on weed control. *Pesquisa Agropecuaria Brasileira* 23(8): 861-868.
- Bai, Q. Y., Y. L. Zhai, et al. (1988). Phosalone applied to cotton (*Gossypium hirsutum* L.): Its deposition and persistence on foliages, soils and final residues in seeds. *Journal Of Environmental Science And Health Part B Pesticides Food Contaminants And Agricultural Wastes* 23(1): 33-44.
- Baker, J., C. Steele, et al. (1988). Sequence and characterization of 6 Lea proteins and their genes from cotton. *Plant Molecular Biology* 11(3): 277-292.
- Baker, J. M. and B. C. H. M. Van (1988). Water transfer through cotton plants connecting soil regions of differing water potential. *Agronomy Journal* 80(6): 993-997.
- Balasubramanian, G., S. Chelliah, et al. (1988). Life-table and intrinsic rate of increase of *Spodoptera litura* Fabricius on castor and cotton. *Entomon* 13(2): 141-146.
- Batson, W. E. and L. E. Trevathan (1988). Suitability and efficacy of ground corncobs as a carrier of *Fusarium solani* spores. *Plant Disease* 72(3): 222-225.
- Beltrao, N. E. D. M., D. M. P. D. Azevedo, et al. (1988). Spacing and sowing density on upland cotton in south-west of Bahia state (Brazil). *Pesquisa Agropecuaria Brasileira* 23(8): 903-910.
- Berger, L. A., B. E. Vaissiere, et al. (1988). *Bombus* spp. (Hymenoptera: Apidae) as pollinators of male-sterile upland cotton on the Texas High Plains (USA). *Environmental Entomology* 17(5): 789-794.
- Bhardwaj, S. N., A. D. Saini, et al. (1988). Contribution of area, thickness and conductance of leaf in biomass production in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 58(2): 100-103.
- Bhardwaj, S. N. and K. P. Singh (1988). Relation between specific leaf weight, leaf conductance, carbon exchange rate and chlorophyll contents in genotypes of upland cotton (*Gossypium hirsutum* Linn.). *Indian Journal Of Plant Physiology* 31(1): 102-104.
- Bilapate, G. G. (1988). Investigations on *Heliothis armigera* (Hubner) in Marathwada (India): XIII. Growth and development on different host plants. *Journal Of Maharashtra Agricultural Universities* 13(2): 186-188.
- Blair, L. C., C. J. Chastain, et al. (1988). Initiation and characterization of a cotton (*Gossypium hirsutum* L.) photoautotrophic cell suspension culture. *Plant Cell Reports* 7(4): 266-269.
- Bland, W. L. and W. A. Dugas (1988). Root length density from minirhizotron observations. *Agronomy Journal* 80(2): 271-275.
- Bourland, F. M., G. Kaiser, et al. (1988). Rapid deterioration of cotton, *Gossypium hirsutum* L., seed using hot water. *Seed Science And Technology* 16(3): 673-684.

- Bridges, D. C. and J. M. Chandler (1988). Influence of cultivar height on competitiveness of cotton (*Gossypium hirsutum*) with johnsongrass (*Sorghum halepense*). *Weed Science* 36(5): 616-620.
- Brown, S. M. and T. Whitwell (1988). Influence of tillage on horseweed, *Conyza canadensis*. *Weed Technology* 2(3): 269-270.
- Burke, J. J., J. R. Mahan, et al. (1988). Crop-specific thermal kinetic windows in relation to wheat and cotton biomass production. *Agronomy Journal* 80(4): 553-556.
- Butler, G. D. J., D. L. Coudriet, et al. (1988). Toxicity and repellency of soybean and cottonseed oils to the sweet-potato whitefly and the cotton aphid on cotton in greenhouse studies. *Southwestern Entomologist* 13(2): 81-86.
- Butler, G. D. J., W. D. Hutchison, et al. (1988). Effect of aldicarb treatments to cotton on *Bemisia tabaci* and *Aphis gossypii* populations in Israel. *Southwestern Entomologist* 13(2): 87-94.
- Byrd, J. D. J. and A. C. York (1988). Interactions of carbaryl and dimethoate with sethoxydim. *Weed Technology* 2(4): 433-436.
- Cakmak, I. and H. Marschner (1988). Zinc-dependent changes in ESR signals, NADPH oxidase and plasma membrane permeability in cotton roots. *Physiologia Plantarum* 73(1): 182-186.
- Cakmak, I. and H. Marschner (1988). Increase in membrane permeability and exudation in roots of zinc deficient plants. *Journal Of Plant Physiology* 132(3): 356-361.
- Castrejon, S. A. (1988). Evaluation of lines and varieties of the cotton plant (*Gossypium hirsutum* L.) for tolerance to Cotton Rust (*Puccinia cacabata* Arth. and Holow). *Revista Mexicana De Fitopatologia* 6(1): 73-75.
- Castrejon, S. A. (1988). Induced resistance: II. Response of the cotton plant (*Gossypium hirsutum* L.) to artificial inoculation with low-virulence *Verticillium dahliae* Kleb. in a field infested with high-virulence races. *Revista Mexicana De Fitopatologia* 6(2): 180-184.
- Castrejon, S. A. and A. J. Galindo (1988). Induced resistance: I. Resistance to *Verticillium dahliae* race T-9 in cotton (*Gossypium hirsutum* L.) variety Coker 310, induced by *Verticillium albo atrum* in greenhouse conditions. *Revista Mexicana De Fitopatologia* 6(2): 177-179.
- Cathey, G. W. and W. R. J. Meredith (1988). Cotton response to planting date and mepiquat chloride. *Agronomy Journal* 80(3): 463-466.
- Chang, J. F., J. H. Benedict, et al. (1988). Pheromone production of the boll weevil in response to seven cotton genotypes grown in two environments. *Environmental Entomology* 17(6): 921-925.
- Chang, J. F., J. H. Benedict, et al. (1988). Volatile monoterpenes collected from the air surrounding flower buds of seven cotton genotypes. *Crop Science* 28(4): 685-688.
- Cohen, Y., M. Fuchs, et al. (1988). Calibrated heat pulse method for determining water uptake in cotton. *Agronomy Journal* 80(3): 398-402.
- Combrink, N. J. J. (1988). An hypothesis concerning the development of the red leaf disorder in cotton (*Gossypium hirsutum* L.). *South African Journal Of Plant And Soil* 5(2): 110-111.
- Constable, G. A. and I. J. Rochester (1988). Nitrogen application to cotton on clay soil: Timing and soil testing. *Agronomy Journal* 80(3): 498-502.
- Dale, J. E. (1988). Effect of AC 94337 on young grain sorghum (*Sorghum bicolor*) and rotational crops. *Weed Technology* 2(4): 456-459.

- Davis, D. D., F. L. Carter, et al. (1988). Mixed plantings of upland male sterile and Pima restorer for increased bee pollination in production of F-1 interspecific hybrid cotton seed. *Southwestern Entomologist* 13(2): 113-118.
- Dhawan, A. K., A. S. Sidhu, et al. (1988). Assessment of avoidable loss in cotton (*Gossypium hirsutum* and *Gossypium arboreum*) due to sucking pests and bollworms. *Indian Journal Of Agricultural Sciences* 58(4): 290-292.
- Dimitrova, L. and M. Bozhinov (1988). Variability of cotton fiber properties caused by the genotype and some environmental factors. *Rasteniev"Dni Nauki* 25(9): 27-34.
- Duhoon, S. S. and P. V. Verma (1988). Effect of gamma -rays on heterosis in cotton (*Gossypium arboreum* and *Gossypium hirsutum*) hybrids. *Indian Journal Of Agricultural Sciences* 58(3): 184-187.
- Finer, J. J. (1988). Plant regeneration from somatic embryogenic suspension cultures of cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 7(6): 399-402.
- Fleischer, S. J., M. J. Gaylor, et al. (1988). Dispersal of *Lygus lineolaris* (Heteroptera: Miridae) adults through cotton following nursery host destruction. *Environmental Entomology* 17(3): 533-541.
- Flint, H. M., N. J. Curtice, et al. (1988). Development of pink bollworm populations (Lepidoptera: Gelechiidae) on nectaried and nectariless deltapine cotton in field cages. *Environmental Entomology* 17(2): 306-308.
- Francl, L. J., T. D. Wyllie, et al. (1988). Influence of crop rotation on population density of *Macrophomina phaseolina* in soil infested with *Heterodera glycines*. *Plant Disease* 72(9): 760-764.
- Fridlyand, L. E., V. L. Kaler, et al. (1988). Assessment of the internal parameters of photosynthetic apparatus in the cotton leaf by carbon dioxide exchange curves. *Fiziologiya Rastanii* 35(1): 24-30.
- Galau, G. A., H. W. Bass, et al. (1988). Restriction fragment length polymorphisms in diploid and allotetraploid *Gossypium*: Assigning the late embryogenesis -abundant (Lea) alleles in *Gossypium hirsutum*. *Molecular & General Genetics* 211(2): 305-314.
- Graves, D. A. and J. M. Stewart (1988). Chronology of the differentiation of cotton (*Gossypium hirsutum* L.) fiber cells. *Planta* 175(2): 254-258.
- Graves, D. A. and J. M. Stewart (1988). Analysis of the protein constituency of developing cotton fibers. *Journal Of Experimental Botany* 39(198): 59-69.
- Greeff, A. I. and J. J. Human (1988). The effect of date of planting on the fiber properties of four cotton cultivars grown under irrigation. *South African Journal Of Plant And Soil* 5(4): 167-172.
- Green, J. D., D. S. Murray, et al. (1988). Soil water relations of silverleaf nightshade (*Solanum elaeagnifolium*) with cotton (*Gossypium hirsutum*). *Weed Science* 36(6): 740-746.
- Guinn, G. and D. L. Brummett (1988). Changes in abscisic acid and IAA before and after anthesis relative to changes in abscission rates of cotton fruiting forms. *Plant Physiology* 87(3): 629-631.
- Guinn, G. and D. L. Brummett (1988). Changes in free and conjugated IAA and abscisic acid in young cotton fruits and their abscission zones in relation to fruit retention during and after moisture stress. *Plant Physiology* 86(1): 28-31.
- Gutierrez, J. C., G. Garcia, et al. (1988). Identification of (*Gossypium hirsutum* L.). *Investigacion Agraria Produccion Y Proteccion Vegetales* 3(3): 297-314.
- Halevy, J. and T. Markovitz (1988). Foliar NPKS application to cotton from full flowering to boll filling. *Fertilizer Research* 15(3): 247-252.

- Hande, Y. K. and M. S. Rane (1988). Reaction of some important cotton cultures against individual races of the bacterial blight pathogen. *Journal Of Maharashtra Agricultural Universities* 13(1): 17-20.
- Hartung, W., J. W. Radin, et al. (1988). Abscisic acid movement into the apoplastic solution of water-stressed cotton leaves: Role of apoplastic pH. *Plant Physiology* 86(3): 908-913.
- Heagle, A. S., J. E. Miller, et al. (1988). Injury and yield response of cotton to chronic doses of ozone and soil moisture deficit. *Journal Of Environmental Quality* 17(4): 627-635.
- Hedin, P. A., J. C. McCarty, et al. (1988). Plant bioregulator induced increases in the protein content of cotton plant tissues. *Journal Of Agricultural And Food Chemistry* 36(4): 742-745.
- Hedin, P. A., W. L. Parrott, et al. (1988). Elucidating mechanisms of tobacco budworm resistance to allelochemicals by dietary tests with insecticide synergists. *Pesticide Biochemistry And Physiology* 32(1): 55-61.
- Hodgson, A. S. and D. A. Macleod (1988). Seasonal and soil fertility effects on the response of waterlogged cotton to foliar-applied nitrogen fertilizer. *Agronomy Journal* 80(2): 259-265.
- Hodgson, R. H. and R. H. Snyder (1988). Thidiazuron effects on Malvaceae, corn, *Zea mays*; and soybean, *Glycine max*. *Weed Technology* 2(3): 342-349.
- Hofmann, W. C., D. L. Kittock, et al. (1988). Planting seed density in relation to cotton emergence and yield. *Agronomy Journal* 80(5): 834-836.
- Hossain, T. and M. M. Ahmedova (1988). Effects of electromagnetic field on chromosomal behavior of cotton (*Gossypium* spp.). *Bangladesh Journal Of Botany* 17(2): 127-130.
- Hsu, C. L. and B. C. Mullin (1988). A new protocol for isolation of mitochondrial DNA from cotton seedlings. *Plant Cell Reports* 7(5): 356-360.
- Huete, A. R. (1988). A soil-adjusted vegetation index (SAVI). *Remote Sensing Of Environment* 25(3): 295-310.
- Huete, A. R. and R. D. Jackson (1988). Soil and atmosphere influences on the spectra of partial canopies. *Remote Sensing Of Environment* 25(1): 89-106.
- Idso, S. B. (1988). Three phases of plant response to atmospheric carbon dioxide enrichment. *Plant Physiology* 87(1): 5-7.
- Jaeger, C. H., J. D. Goeschl, et al. (1988). Short-term responses of phloem transport to mechanical perturbation. *Physiologia Plantarum* 72(3): 588-594.
- Jalali, S. K., S. P. Singh, et al. (1988). Influence of the food plants on the degree of parasitism of larvae of *Heliothis armigera* by *Cotesia kazak*. *Entomophaga* 33(1): 65-72.
- Ji, D. and J. Zhu (1988). Genetic analysis of oil and amino acid content in shelled seed of upland cotton hybrids. *Acta Agronomica Sinica* 14(1): 1-6.
- Kalisa, P. and E. A. Makram (1988). Effect of sowing date on the yield of rain grown cotton in Burundi. *Annals Of Agricultural Science* 33(2): 881-890.
- Kasymova, G. F., N. I. Koryakina, et al. (1988). Electrophoretic study of storage cottonseed proteins. *Fiziologiya I Biokhimiya Kul'Turnykh Rastenii* 20(3): 263-270.
- Kittock, D. L., E. T. Turcotte, et al. (1988). Estimation of heat tolerance improvement in recent American Pima cotton cultivars. *Journal Of Agronomy And Crop Science* 161(5): 305-309.

- Kleifeld, Y., T. Blumenfeld, et al. (1988). The use of fomesafen for pre-emergence weed control in cotton. *Phytoparasitica* 16(2): 133-144.
- Knapp, S. J. and T. S. Cox (1988). S-1 family recurrent selection in autogamous crop based on dominant genetic male-sterility. *Crop Science* 28(2): 227-231.
- Koryakina, N. I., G. F. Kasymova, et al. (1988). Amino acid composition of storage proteins from cotton seeds. *Fiziologiya I Biokhimiya Kul'Turnykh Rastenii* 20(4): 393-397.
- Kumar, V. (1988). Response of late-sown cotton (*Gossypium hirsutum* L.) to dates of sowing and plant population in the Nigerian savanna. *Crop Research* 1(1): 58-68.
- Kunce, C. M., R. N. Trelease, et al. (1988). Purification and biosynthesis of cottonseed (*Gossypium hirsutum* L.) catalase. *Biochemical Journal* 251(1): 147-156.
- Lauter, D. J., A. Meiri, et al. (1988). Isoosmotic regulation of cotton and peanut at saline concentrations of potassium and sodium. *Plant Physiology* 87(4): 911-916.
- Lima, E. F., J. M. F. C. Carvalho, et al. (1988). Survival of *Colletotrichum gossypii* var. *cephalosporioides* on cotton seed (*Gossypium hirsutum* latifolium). *Fitopatologia Brasileira* 13(3): 247-248.
- Makram, E. (1988). Cotton response to mepiquat chloride in the Republic of Burundi. *Annals Of Agricultural Science* 33(2): 931-940.
- Makram, E. A. (1988). Effect of foliar nutrition feeding by commercial formulations of macro and micro-elements on growth, yield and some fiber properties of the cotton cultivar Pan 575 (*Gossypium hirsutum* + L.). *Annals Of Agricultural Science* 33(2): 903-916.
- Makram, E. A. (1988). Cotton skip-row studies in Burundi. *Annals Of Agricultural Science* 33(2): 891-902.
- Makram, E. A. (1988). Effect of foliar nutrition feeding rates and soil fertilization on growth, yield and some fiber properties of cotton cultivar Pan 575 (*Gossypium hirsutum* L.). *Annals Of Agricultural Science* 33(2): 917-930.
- Martin, W. R. J., P. W. Morgan, et al. (1988). Cotton fleahopper and associated microorganisms as components in the production of stress ethylene by cotton. *Plant Physiology* 87(1): 280-285.
- Mauney, J. R. and D. L. Hendrix (1988). Responses of glasshouse grown cotton to irrigation with carbon dioxide-saturated water. *Crop Science* 28(5): 835-838.
- Mehla, A. S., B. R. Mor, et al. (1988). Genetic analysis of earliness characters in upland cotton (*Gossypium hirsutum* L.). *Crop Improvement* 15(2): 156-159.
- Miller, J. E., R. P. Patterson, et al. (1988). Growth of cotton under chronic ozone stress at two levels of soil moisture. *Journal Of Environmental Quality* 17(4): 635-643.
- Milyaeva, E. L., Z. R. Khaitova, et al. (1988). Peculiarities of growth, development and bud differentiation in a cotton mutant. *Fiziologiya Rastenii* 35(2): 218-225.
- Misra, R. K., A. M. Alston, et al. (1988). Role of root hairs in phosphorus depletion from a macrostructured soil. *Plant And Soil* 107(1): 11-18.
- Nadarajan, N. and S. R. S. Rangasamy (1988). Inheritance of the fuzzless-lintless character in cotton (*Gossypium hirsutum*). *Theoretical And Applied Genetics* 75(5): 728-730.
- Nakayama, F. S. and B. A. Kimball (1988). Soil carbon dioxide distribution and flux within the open-top chamber. *Agronomy Journal* 80(3): 394-398.

- Naqvi, K. M. and A. G. Khanzada (1988). Preliminary studies on the effectiveness of oil-based ULV insecticide formulations against cotton jassid and thrips. *Pakistan Journal Of Agricultural Research* 9(1): 100-105.
- Naqvi, K. M., A. G. Khanzada, et al. (1988). Effectiveness of different insecticides to control brown thrips on cotton. *Pakistan Journal Of Agricultural Research* 9(2): 185-188.
- Navon, A., M. Zur, et al. (1988). Effects of cotton leaf surface alkalinity on feeding of *Spodoptera littoralis* larvae. *Journal Of Chemical Ecology* 14(3): 839-844.
- Nayyar, H., A. S. Basra, et al. (1988). Carbon-14 bicarbonate fixation into lipids of elongating cotton fibers in two *Gossypium* spp. *Journal Of Nuclear Agriculture And Biology* 17(4): 243-245.
- Nelson, E. B. (1988). Biological control of pythium seed rot and preemergence damping-off of cotton with *Enterobacter cloacae* and *Erwinia herbicola* applied as seed treatments. *Plant Disease* 72(2): 140-142.
- Oosterhuis, D. M. and M. J. Urwiler (1988). Cotton main-stem leaves in relation to vegetative development and yield. *Agronomy Journal* 80(1): 65-67.
- Ortegon, E. J., N. C. Torres, et al. (1988). Study on the longevity, habits, progeny and preliminary evaluation of *Meteorus laphygmae* (Viereck), parasite of *Spodoptera* sp. *Revista Colombiana De Entomologia* 14(1): 7-12.
- Pandey, S. S. and V. V. R. Subrahmanyam (1988). Biosaturation of cyclopropane fatty acids in germinating cottonseed. *Phytochemistry* 27(12): 3963-3964.
- Pandey, S. S. and V. V. R. Subrahmanyam (1988). Lipid changes in maturing and germinating cottonseeds. *Phytochemistry* 27(11): 3405-3410.
- Papadopoulos, I. and Y. Stylianou (1988). Trickle irrigation of cotton with treated sewage effluent. *Journal Of Environmental Quality* 17(4): 574-580.
- Patterson, D. T. (1988). Growth and water relations of cotton (*Gossypium hirsutum*), spurred anoda (*Anoda cristata*), and velvetleaf (*Abutilon theophrasti*) during simulated drought and recovery. *Weed Science* 36(3): 318-324.
- Patterson, D. T., M. T. Highsmith, et al. (1988). Effects of temperature and carbon dioxide concentration on the growth of cotton (*Gossypium hirsutum*), spurred anoda (*Anoda cristata*), and velvetleaf (*Abutilon theophrasti*). *Weed Science* 36(6): 751-757.
- Peeters, M. C., S. Voets, et al. (1988). Pattern of nucleolar growth in differentiating cotton fibers (*Gossypium hirsutum* L.). *Annals Of Botany* 62(4): 377-382.
- Peng, C. and J. Guo (1988). A study of genetic divergence of quantitative characters in upland cotton (*Gossypium hirsutum* L.). *Scientia Agricultura Sinica* 21(4): 26-32.
- Qi, G. R., G. J. Cao, et al. (1988). Studies on the sites expressing evolutionary changes in the structure of eukaryotic 5S ribosomal RNA. *Journal Of Molecular Evolution* 27(4): 336-340.
- Qian, S., J. Huang, et al. (1988). Investigation on interspecific hybridization of *Gossypium* species. *Acta Agronomica Sinica* 14(2): 96-102.
- Qiu, X., Y. Pai, et al. (1988). Studies on the photosynthetic characteristics of canopy in cotton cultivars (*Gossypium hirsutum* L.). *Acta Agronomica Sinica* 14(4): 315-321.
- Radin, J. W. and D. H. Hendrix (1988). The apoplastic pool of abscisic acid in cotton leaves in relation to stomatal closure. *Planta* 174(2): 180-186.

Rakhmankulov, C., P. Abidov, et al. (1988). Component composition of water-soluble proteins of the cotton *Gossypium hirsutum* seeds. *Fiziologiya I Biokhimiya Kul'Turnykh Rastenii* 20(5): 477-480.

Rakhmatov, N. A., A. P. Ibragimov, et al. (1988). Comparative characteristics of ribosomes and ribosomal proteins in cotton chloroplasts and cytoplasm. *Fiziologiya Rastenii* 35(4): 805-808.

Ram, C., B. S. Chhabra, et al. (1988). Correlation among laboratory tests and field emergence in cotton. *Seed Research* 16(1): 47-50.

Ramalho, F. S. and F. M. M. Jesus (1988). Distribution of boll weevil (*Anthonomus grandis* Boheman) eggs within cotton plants. *Tropical Agriculture* 65(3): 245-248.

Reddy, V. R. and D. N. Baker (1988). Estimation of parameters for the cotton simulation model GOSSYM: Cultivar differences. *Agricultural Systems* 26(2): 111-122.

Reeves, D. W., J. T. Touchton, et al. (1988). Effect of nitrogen and dicyandiamide on growth and water relations of cotton. *Soil Science Society Of America Journal* 52(1): 281-285.

Rickerl, D. H., W. B. Gordon, et al. (1988). Winter legume and tillage effects on cotton growth and soil ecology. *Soil & Tillage Research* 11(1): 63-72.

Riffle, M. S., W. E. Thilsted, et al. (1988). Germination and seed production of unicorn plant (*Proboscidea louisianica*). *Weed Science* 36(6): 787-791.

Rodriguez, G. B. and J. R. Barrow (1988). Pollen selection for heat tolerance in cotton. *Crop Science* 28(5): 857-859.

Rodriguez, K. R., P. S. King, et al. (1988). Potential of crops uncommon to Alabama (USA) for management of root-knot and soybean cyst nematodes. *Annals Of Applied Nematology* 2: 116-120.

Rotem, J., J. Eidt, et al. (1988). Relative effects of *Alternaria alternata* and *Alternaria macrospora* on cotton crops in Israel. *Plant Pathology* 37(1): 16-19.

Rotem, J., U. Wendt, et al. (1988). The effect of sunlight on symptom expression of *Alternaria alternata* on cotton. *Plant Pathology* 37(1): 12-15.

Sahay, R. K., N. L. Bhale, et al. (1988). Involvement of ethylene with new wilt of cotton (*Gossypium* species). *Indian Journal Of Agricultural Sciences* 58(12): 885-887.

Schmid, K. M. and G. W. Patterson (1988). Distribution of cyclopropanoid fatty acids in malvaceous plant parts. *Phytochemistry* 27(9): 2831-2834.

Sharma, J. P. (1988). Evaluation of some synthetic pyrethroids and carbaryl for the control of bollworms and their effect on yield parameters of cotton. *Entomon* 13(3-4): 215-222.

Shende, S. T., M. Singh, et al. (1988). Effect of seed bacterization with *Azotobacter chroococcum* on yield of upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 58(3): 206-209.

Sheng, C. F. (1988). Economic threshold of the third generation of cotton bollworm in north China. *Acta Entomologica Sinica* 31(1): 36-41.

Singh, J., Z. S. Dhaliwal, et al. (1988). Suitability of different plant parts of some host species for neonate survival of *Heliothis armigera* (Hubner). *Annals Of Biology* 4(1-2): 33-37.

Singh, M., S. N. Bhardwaj, et al. (1988). Yield improvement through simultaneous improvement of yield components in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 58(6): 425-428.

- Singh, M., V. P. Singh, et al. (1988). Outcrossing for improvement in yield and quality of upland cotton (*Gossypium hirsutum*) in north India. *Indian Journal Of Agricultural Sciences* 58(2): 110-113.
- Singh, V. V., M. K. Meshram, et al. (1988). Evolution of upland cotton (*Gossypium hirsutum*) lines with multiracial resistance to bacterial blight caused by *Xanthomonas campestris* pathovar *malvacearum*. *Indian Journal Of Agricultural Sciences* 58(4): 255-258.
- Standley, J., E. A. Clarke, et al. (1988). Modification by crop rotations of the nitrogen fertilizer requirements for irrigated cotton and soil test calibration. *Queensland Journal Of Agricultural And Animal Sciences* 45(2): 129-138.
- Stark, S. B. and K. R. Hopper (1988). *Chrysoperla carnea* predation on *Heliothis virescens* larvae parasitized by *Microplitis croceipes*. *Entomologia Experimentalis Et Applicata* 48(1): 69-72.
- Stegink, S. J. and K. C. Vaughn (1988). Norflurazon (SAN-9789) reduces abscisic acid levels in cotton seedlings: A glandless isolate is more sensitive than its glanded counterpart. *Pesticide Biochemistry And Physiology* 31(3): 269-275.
- Stelly, D. M., J. A. Lee, et al. (1988). Proposed schemes for mass-extraction of doubled haploids of cotton. *Crop Science* 28(6): 885-890.
- Subbaraj, R., K. S. Sundararajan, et al. (1988). Temperature compensated nature of circadian rhythms in tropics. *Indian Journal Of Experimental Biology* 26(4): 277-279.
- Susin, J., M. Clemente, et al. (1988). Evaluation of taxonomic characters coming from structures of epidermic origin for intraspecific differentiation in *Gossypium hirsutum* L. *Agronomie* 8(1): 61-70.
- Suttle, J. C. (1988). Disruption of the polar auxin transport system in cotton seedlings following treatment with the defoliant thidiazuron. *Plant Physiology* 86(1): 241-245.
- Temple, P. J., R. S. Kupper, et al. (1988). Injury and yield responses of differentially irrigated cotton to ozone. *Agronomy Journal* 80(5): 751-755.
- Thangaraju, D., S. Parameswari, et al. (1988). Insecticidal control of bollworms in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 58(7): 533-536.
- Trolinder, N. L. and J. R. Goodin (1988). Somatic embryogenesis in cotton (*Gossypium*): II. Requirements for embryo development and plant regeneration. *Plant Cell Tissue And Organ Culture* 12(1): 43-54.
- Trolinder, N. L. and J. R. Goodin (1988). Somatic embryogenesis in cotton (*Gossypium*): I. Effects of source of explant and hormone regime. *Plant Cell Tissue And Organ Culture* 12(1): 31-42.
- Tyagi, A. P. and P. A. S. Oling'Otie (1988). Screening for resistance to bacterial blight in Upland cotton (*Gossypium Hirsutum* L.). *Plant Breeding* 101(2): 164-166.
- Upchurch, D. R. and J. R. Mahan (1988). Maintenance of constant leaf temperature by plants: II. Experimental observations in cotton. *Environmental And Experimental Botany* 28(4): 359-366.
- Van, H. H. G., S. W. H. Van, et al. (1988). Comparison of cotton (*Gossypium hirsutum* L.) yield and fiber properties over locations and seasons. *South African Journal Of Plant And Soil* 5(2): 75-78.
- Wells, R. (1988). Response of leaf ontogeny and photosynthetic activity to reproductive growth in cotton. *Plant Physiology* 87(1): 274-279.
- Wells, R., W. R. J. Meredith, et al. (1988). Heterosis in upland cotton: II. Relationship of leaf area to plant photosynthesis. *Crop Science* 28(3): 522-525.

- Wilcut, J. W., M. G. Patterson, et al. (1988). Efficacy and economics of pendimethalin herbicide combinations for weed control in cotton (*Gossypium hirsutum* L.). *Applied Agricultural Research* 3(4): 203-208.
- Wilson, F. D. and J. L. Szaro (1988). A rapid field technique for infesting cotton plants with pink bollworm (Lepidoptera: Gelechiidae) eggs. *Journal Of Economic Entomology* 81(3): 959-962.
- Wullschleger, S. D., M. A. Dixon, et al. (1988). Field measurement of leaf water potential with a temperature-corrected in situ thermocouple psychrometer. *Plant Cell And Environment* 11(3): 199-204.
- Yoo, K. H., J. T. Touchton, et al. (1988). Runoff, sediment and nutrient losses from various tillage systems of cotton. *Soil & Tillage Research* 12(1): 13-24.
- Zaki, M. S., M. R. Hussain, et al. (1988). Response of seed cotton yield and technological properties of cotton to some herbicides. *Pakistan Journal Of Agricultural Research* 9(4): 448-452.
- Zeringue, H. J. J. (1988). Production of carpel wall phytoalexins in the developing cotton boll. *Phytochemistry* 27(11): 3429-3432.
- Zhang, X., J. Sun, et al. (1988). Studies on *Gossypium* distant hybrid ovule culture in vitro. *Scientia Agricultura Sinica* 21(6): 53-58.
- Zhong, H. and A. Lauchli (1988). Incorporation of carbon-14 glucose into cell wall polysaccharides of cotton roots: Effects of sodium chloride and calcium chloride. *Plant Physiology* 88(3): 511-514.
- Zimmerman, T. W. and C. D. Robacker (1988). Media and gelling agent effect on cotton callus initiation from excised seed hypocotyls. *Plant Cell Tissue And Organ Culture* 15(3): 269-274.
- Altman, D. W., D. M. Stelly, et al. (1987). Introgression of the glanded-plant and glandless-seed trait from *Gossypium sturtianum* Willis into cultivated upland cotton using ovule culture. *Crop Science* 27(5): 880-884.
- Armentrout, V. N. and A. J. Downer (1987). Infection cushion development by *Rhizoctonia solani* on cotton. *Phytopathology* 77(4): 619-623.
- Armentrout, V. N., A. J. Downer, et al. (1987). Factors affecting infection cushion development by *Rhizoctonia solani* on cotton. *Phytopathology* 77(4): 623-630.
- Atuahene, C. C., A. Donkoh, et al. (1987). The use of raw cottonseed meal (CSM) in layer diets. *Journal Of Animal Production Research* 7(1): 1-8.
- Azevedo, D. M. P. D., N. E. D. M. Beltrao, et al. (1987). Mixtures of herbicides against weeds in annual cotton. *Pesquisa Agropecuaria Brasileira* 22(8): 797-802.
- Babu, V. R., S. N. Prasad, et al. (1987). Evaluation of cotton genotypes for tolerance to saline water irrigations. *Indian Journal Of Agronomy* 32(3): 229-231.
- Baker, S. H. (1987). Effects of tillage practices on cotton double cropped with wheat. *Agronomy Journal* 79(3): 513-516.
- Bar, T. A. and J. Rudich (1987). Osmotic adjustment of cotton to moderate potassium-chloride stress and subsequent water stress during early stages of development. *Agronomy Journal* 79(1): 166-171.
- Basu, A. K. and M. G. Bhat (1987). Variability for harvest index in upland cotton. *Indian Journal Of Agricultural Sciences* 57(4): 236-240.
- Bhatt, J. G. (1987). Leaf growth, reproduction growth and yield in cotton (*Gossypium hirsutum* L.). *Journal Of Agronomy And Crop Science* 159(4): 264-268.

- Bhatt, J. G. and E. Appukuttan (1987). Changes in concentrations of major nutrients in developing bolls of hybrid cotton. *Indian Journal Of Plant Physiology* 30(2): 160-164.
- Birajdar, J. M., K. R. Pawar, et al. (1987). Studies on planting pattern, spacing and intercropping in hybrid-4, cotton under rainfed conditions. *Journal Of Maharashtra Agricultural Universities* 12(1): 67-69.
- Bozhinov, M. and L. Dimitrova (1987). A study of the economic and technological characteristics of some foreign and Bulgarian cotton cultivars. *Rasteniev"Dni Nauki* 24(9): 51-59.
- Bridges, D. C. and J. M. Chandler (1987). Effect of herbicide and weed height on johnsongrass (*Sorghum halepense*) control and cotton (*Gossypium hirsutum*) yield. *Weed Technology* 1(3): 207-211.
- Bridges, D. C. and J. M. Chandler (1987). Influence of johnsongrass (*Sorghum halepense*) density and period of competition on cotton yield. *Weed Science* 35(1): 63-67.
- Brown, J. K., J. D. Mihail, et al. (1987). Effects of cotton leaf crumple virus on cotton inoculated at different growth stages. *Plant Disease* 71(8): 699-703.
- Brown, S. M., J. M. Chandler, et al. (1987). Weed control in a conservation tillage rotation in the Texas Blacklands (USA). *Weed Science* 35(5): 695-699.
- Bryson, C. T. (1987). Interference of hemp sesbania (*Sesbania exaltata*) with cotton (*Gossypium hirsutum*). *Weed Science* 35(3): 314-318.
- Bucheli, P., A. J. Buchala, et al. (1987). Autolysis in vitro of cotton (*Gossypium hirsutum*) fiber cell walls. *Physiologia Plantarum* 70(4): 633-638.
- Byrd, J. D. J. and A. C. York (1987). Annual grass control in cotton (*Gossypium hirsutum*) with fluazifop, sethoxydim, and selected dinitroaniline herbicides. *Weed Science* 35(3): 388-394.
- Byrd, J. D. J. and A. C. York (1987). Interaction of fluometuron and MSMA with sethoxydim and fluazifop. *Weed Science* 35(2): 270-276.
- Cakmak, I. and H. Marschner (1987). Mechanism of phosphorus-induced zinc deficiency in cotton: III. Changes in physiological availability of zinc in plants. *Physiologia Plantarum* 70(1): 13-20.
- Carter, C. H. and P. E. Keeley (1987). Selective control of johnsongrass (*Sorghum halepense*) in cotton (*Gossypium hirsutum*) with foliar herbicides. *Weed Science* 35(3): 418-421.
- Cathey, G. W. and J. C. Bailey (1987). Evaluation of chlordimeform for cotton yield enhancement. *Journal Of Economic Entomology* 80(3): 670-674.
- Chang, C. W. and R. D. Ryan (1987). Effects of water stress on starch and sucrose metabolism in cotton leaves. *Starch Staerke* 39(3): 84-87.
- Chlan, C. A., K. Borroto, et al. (1987). Developmental biochemistry of cottonseed embryogenesis and germination: XIX. Sequences and genomic organization of the alpha globulin (vicilin) genes of cottonseed. *Plant Molecular Biology* 9(6): 533-546.
- Cole, D. J. and W. J. Owen (1987). Influence of monooxygenase inhibitors on the metabolism of the herbicides chlortoluron and metolachlor in cell suspension cultures. *Plant Science* 50(1): 13-20.
- Combrink, N. J. J. and E. A. Davies (1987). Effect of calcium, potassium and boron supply on growth of cotton (*Gossypium hirsutum* L.) and tobacco (*Nicotiana tabacum* L.). *South African Journal Of Plant And Soil* 4(3): 143-144.
- Contolini, C. S. and M. Y. Menzel (1987). Early development of duplication-deficiency ovules in upland cotton. *Crop Science* 27(2): 345-348.

- Correll, J. C., T. R. Gordon, et al. (1987). Host range, specificity, and biometrical measurements of *Leveillula taurica* in California (USA). *Plant Disease* 71(3): 248-251.
- Cramer, G. R., J. Lynch, et al. (1987). Influx of sodium, potassium, and calcium into roots of salt-stressed cotton seedlings: Effects of supplemental calcium. *Plant Physiology* 83(3): 510-516.
- Dani, R. G. and R. J. Kohel (1987). Effects of time of boll set on seed-oil content in upland cotton. *Indian Journal Of Agricultural Sciences* 57(6): 391-394.
- De, V. D. A. (1987). Alternative hosts for the black root rot fungus *Thielaviopsis basicola* of tobacco. *Phytophylactica* 19(3): 347-348.
- De, V. D. A. (1987). Influence of rotation crops on the tobacco fungus black shank (*Phytophthora nicotianae* var. *nicotianae*). *Phytophylactica* 19(1): 115-116.
- Devey, M. E. and M. L. Roose (1987). Genetic analysis of *Verticillium* wilt tolerance in cotton using pedigree data from three crosses. *Theoretical And Applied Genetics* 74(1): 162-167.
- Dickens, J. C. and H. R. Agee (1987). Photosensitivity and spectral response of boll weevils reared on natural host and artificial larval diets supplemented with carotenoids. *Journal Of Applied Entomology* 103(2): 185-192.
- Eck, H. V., A. C. Mathers, et al. (1987). Plant water stress at various growth stages and growth and yield of soybeans. *Field Crops Research* 17(1): 1-16.
- Ehleringer, J. R. and S. D. Hammond (1987). Solar tracking and photosynthesis in cotton leaves. *Agricultural And Forest Meteorology* 39(1): 25-36.
- Elzen, G. W., H. J. Williams, et al. (1987). Comparative flight behavior of parasitoids *Campoletis sonorensis* and *Microplitis croceipes*. *Entomologia Experimentalis Et Applicata* 45(2): 175-180.
- Fallieri, J. and J. N. Jenkins (1987). The use of haploids of cotton in interspecific crosses. *Pesquisa Agropecuaria Brasileira* 22(1): 83-88.
- Firoozabady, E., D. L. Deboer, et al. (1987). Transformation of cotton (*Gossypium hirsutum* L.) by *Agrobacterium tumefaciens* and regeneration of transgenic plants. *Plant Molecular Biology* 10(2): 105-116.
- Fornazier, M. J. and O. Nakano (1987). Chemicals efficiency on the control of the pink bollworm, using a new experimental methodology. *Pesquisa Agropecuaria Brasileira* 22(9-10): 923-928.
- Fryrear, D. W. (1987). Ridging reduces wind damage to cotton. *Applied Agricultural Research* 1(5): 311-314.
- Galau, G. A., N. Bijaisoradat, et al. (1987). Accumulation kinetics of cotton late embryogenesis -abundant messenger RNA species and storage protein messenger RNA species: Coordinate regulation during embryogenesis and the role of abscisic acid. *Developmental Biology* 123(1): 198-212.
- Galau, G. A. and D. W. Hughes (1987). Coordinate accumulation of homeologous transcripts of seven cotton *Lea* gene families during embryogenesis and germination. *Developmental Biology* 123(1): 213-221.
- Garg, H. R., T. H. Singh, et al. (1987). Genetic analysis through triple-test cross in the F-2 population of upland cotton. *Indian Journal Of Agricultural Sciences* 57(10): 701-704.
- Garg, H. R., T. H. Singh, et al. (1987). Genetical analysis of quantitative traits in two backcross populations of upland cotton (*Gossypium hirsutum* L.). *Genetica Agraria* 41(4): 419-426.
- Genty, B., J. M. Briantais, et al. (1987). Effects of drought on primary photosynthetic processes of cotton leaves. *Plant Physiology* 83(2): 360-364.

- Gerik, T. J., J. E. J. Morrison, et al. (1987). Effects of controlled-traffic on soil physical properties and crop rooting. *Agronomy Journal* 79(3): 434-438.
- Gesos, K. F. and O. Nagymetov (1987). Inheritance of fiber yield components in intraspecific cotton hybrids and an evaluation of the combining ability of parental forms in the diallel crossing system. *Sel'Skokhozyaistvennaya Biologiya*(1): 76-79.
- Gigou, J. (1987). The importance of phosphorus deficiency for annual crops in the Ivory Coast. *Agronomie Tropicale Riz Et Riziculture Et Cultures Vivrieres Tropicales* 42(1): 20-28.
- Gigou, J. (1987). Crop adaptation in central Ivory Coast. *Agronomie Tropicale Riz Et Riziculture Et Cultures Vivrieres Tropicales* 42(1): 1-12.
- Gigou, J. and P. F. Chabalier (1987). Utilization of nitrogenous fertilizer by annual crops in the Ivory Coast. *Agronomie Tropicale* 42(3): 171-179.
- Gill, M. S. and Y. P. S. Bajaj (1987). Hybridization between diploid (*Gossypium arboreum*) and tetraploid (*Gossypium hirsutum*) cotton through ovule culture. *Euphytica* 36(2): 625-630.
- Graham, C. T. J., J. N. Jenkins, et al. (1987). Effects of mepiquat chloride on natural plant resistance to tobacco budworm in cotton. *Crop Science* 27(2): 360-361.
- Green, J. D., D. S. Murray, et al. (1987). Full-season interference of silverleaf nightshade (*Solanum elaeagnifolium*) with cotton (*Gossypium hirsutum*). *Weed Science* 35(6): 813-818.
- Grimes, D. W., H. Yamada, et al. (1987). Climate-normalized cotton leaf water potentials for irrigation scheduling. *Agricultural Water Management* 12(4): 293-304.
- Grisham, M. P., W. L. Sterling, et al. (1987). Characterization of the induction of stress ethylene synthesis in cotton caused by the cotton fleahopper (Hemiptera: Miridae) and its microorganisms. *Annals Of The Entomological Society Of America* 80(3): 411-416.
- Guinn, G. and D. L. Brummett (1987). Concentrations of abscisic acid and IAA in cotton fruits and their abscission zones in relation to fruit retention. *Plant Physiology* 83(1): 199-202.
- Gupta, S. P. and T. H. Singh (1987). Heterosis and inbreeding depression for seed cotton yield and some seed and fiber attributes in upland cotton (*Gossypium hirsutum* L.). *Crop Improvement* 14(1): 14-17.
- Gutierrez, A. P., F. Schulthess, et al. (1987). Energy acquisition and allocation in plants and insects: A hypothesis for the possible role of hormones in insect feeding patterns. *Canadian Entomologist* 119(2): 109-130.
- Guye, M. G. and J. M. Wilson (1987). The effects of chilling and chill-hardening temperatures on stomatal behavior in a range of chill-sensitive species and cultivars. *Plant Physiology And Biochemistry* 25(6): 717-722.
- Halevy, J., A. Marani, et al. (1987). Growth and nitrogen, phosphorus, potassium uptake of high-yielding cotton grown at different nitrogen levels in a permanent-plot experiment. *Plant And Soil* 103(1): 39-44.
- Hassidim, M., B. Rubinstein, et al. (1987). Generation of a membrane potential by electron transport in plasmalemma-enriched vesicles of cotton and radish. *Plant Physiology* 85(4): 872-875.
- Hatfield, J. L., J. E. Quisenberry, et al. (1987). Use of canopy temperatures to identify water conservation in cotton germplasm. *Crop Science* 27(2): 269-273.
- Heilman, J. L. and M. R. Kress (1987). Effects of vegetation on spectral irradiance at the soil surface. *Agronomy Journal* 79(5): 765-768.

- Hendrix, D. L. and K. K. Peelen (1987). Artifacts in the analysis of plant tissues for soluble carbohydrates. *Crop Science* 27(4): 710-715.
- Hocking, P. J., D. C. Reicosky, et al. (1987). Effects of intermittent waterlogging on the mineral nutrition of cotton. *Plant And Soil* 101(2): 211-222.
- Hofmann, W. C., M. M. Karpiscak, et al. (1987). Response of cotton, alfalfa, and cantaloupe to foliar-deposited salt in an arid environment. *Journal Of Environmental Quality* 16(3): 267-272.
- Hopper, N. W. and H. R. Hinton (1987). Electrical conductivity as a measure of planting seed quality in cotton. *Agronomy Journal* 79(1): 147-152.
- Huete, A. R. (1987). Soil-dependent spectral response in a developing plant canopy. *Agronomy Journal* 79(1): 61-68.
- Hughes, D. W. and G. A. Galau (1987). Translation efficiency of Lea messenger RNAs in cotton embryos: Minor changes during embryogenesis and germination. *Plant Molecular Biology* 9(4): 301-314.
- Hussain, M. M., U. Melcher, et al. (1987). Replication of cauliflower mosaic virus DNA in leaves and suspension culture protoplasts of cotton. *Plant Physiology* 83(3): 633-639.
- Idso, S. B., B. A. Kimball, et al. (1987). Atmospheric carbon dioxide enrichment effects on cotton midday foliage temperature: Implications for plant water use and crop yield. *Agronomy Journal* 79(4): 667-672.
- Jagtap, D. R. and A. K. Kolhe (1987). Graphical and combining ability analyses in upland cotton. *Indian Journal Of Agricultural Sciences* 57(7): 456-464.
- Kapoor, H. C., S. Prakash, et al. (1987). Regulation of in vivo nitrate reductase activity in cotton (*Gossypium hirsutum*) leaves in light and dark and the possible role of cytokinin. *Indian Journal Of Biochemistry And Biophysics* 24(6): 326-328.
- Karve, A. D., A. C. Bhalerao, et al. (1987). Photomorphogenic effect of crowding on growth and dry-matter production of crops. *Indian Journal Of Agricultural Sciences* 57(2): 112-116.
- Keeley, P. E., C. H. Carter, et al. (1987). Yellow nutsedge (*Cyperus esculentus*) control with herbicides. *Weed Technology* 1(2): 133-139.
- Keeley, P. E., R. J. Thullen, et al. (1987). Repeated annual applications of herbicides for control of rhizome johnsongrass (*Sorghum halepense*) in cotton (*Gossypium hirsutum*). *Weed Science* 35(1): 75-80.
- Kennedy, C. W., M. T. Ba, et al. (1987). Differences in root and shoot growth and soil moisture extraction between cotton cultivars in an acid subsoil. *Plant And Soil* 101(2): 241-246.
- Khan, I. A., M. A. Khan, et al. (1987). Evaluation of parent varieties in the hybrid cotton production program in Pakistan. *Pakistan Journal Of Scientific And Industrial Research* 30(11): 836-840.
- Klich, M. A. (1987). Relation of plant water potential at flowering to subsequent cottonseed infection by *Aspergillus flavus*. *Phytopathology* 77(5): 739-741.
- Kohel, R. J. and C. R. Benedict (1987). Growth analysis of cottons with differing maturities. *Agronomy Journal* 79(1): 31-34.
- Kumar, H., K. Kaur, et al. (1987). The relationship of hydrolases to fiber development in *Gossypium hirsutum* L. *Proceedings Of The Indian National Science Academy Part B Biological Sciences* 53(3): 259-262.

- Kumar, H., K. Kaur, et al. (1987). The relationship of the hydrolases to fiber development in *Gossypium hirsutum* L. *Proceedings Of The Indian National Science Academy Part B Biological Sciences* 53(3): 259-266.
- Kumar, V., N. P. Mehta, et al. (1987). Influence of stratified harvests and processing intervals on the quality of cotton seed. *Seed Research* 15(1): 1-8.
- Lamoureux, G. L. and D. G. Rusness (1987). EPTC metabolism in corn, cotton, and soybean: Identification of a novel metabolite derived from the metabolism of a glutathione conjugate. *Journal Of Agricultural And Food Chemistry* 35(1): 1-7.
- Lax, A. R., K. C. Vaughn, et al. (1987). Structural and physiological studies of a plastome cotton mutant with slow sorting out. *Journal Of Heredity* 78(3): 147-152.
- Lee, J. A. (1987). Induction of adventitious shoots in cotton. *Crop Science* 27(2): 349-350.
- Loper, G. M., J. Olvey, et al. (1987). Concentration of the systemic gametocide, TD-1123, in cotton nectar, and honeybee response. *Crop Science* 27(3): 558-561.
- Lusby, W. R., J. E. Oliver, et al. (1987). Free and esterified sterols of cotton buds and anthers. *Lipids* 22(2): 80-83.
- Luster, D. G. and R. C. Fites (1987). An abbreviated method for the isolation of mitochondria from storage tissues of oil seedlings. *Physiologia Plantarum* 70(4): 622-625.
- Mane, S. S., N. L. Bhale, et al. (1987). Evaluation of single-cross, multiple-cross and modified back-cross methods based on recombination and character association in upland cotton. *Indian Journal Of Agricultural Sciences* 57(5): 318-321.
- Masner, P., M. Angst, et al. (1987). Fenoxycarb, an insect growth regulator with juvenile hormone activity: A candidate for *Heliothis virescens* (F.) control on cotton. *Pesticide Science* 18(2): 89-94.
- McCarty, J. C. J., J. N. Jenkins, et al. (1987). Genetic resistance to boll weevil oviposition in primitive cotton. *Crop Science* 27(2): 263-264.
- McCarty, J. C. J. and W. L. McGovern (1987). Oviposition preference in two laboratory colonies of boll weevils (Coleoptera: Curculionidae). *Journal Of Economic Entomology* 80(2): 366-368.
- McDowell, L. L., G. H. Willis, et al. (1987). Fenvalerate wash-off from cotton plants by rainfall. *Pesticide Science* 21(2): 83-92.
- McMichael, B. L., J. E. Quisenberry, et al. (1987). Lateral root development in exotic cottons. *Environmental And Experimental Botany* 27(4): 499-502.
- Menges, R. M. (1987). Weed seed population dynamics during six years of weed management systems in crop rotations on irrigated soil. *Weed Science* 35(3): 328-332.
- Menzel, M. Y. and B. J. Dougherty (1987). Transmission of duplication-deficiencies from cotton translocations is unrelated to map lengths of the unbalanced segments. *Genetics* 116(2): 321-330.
- Menzel, M. Y., K. L. Richmond, et al. (1987). A revised map of chromosome 16 in *Gossypium hirsutum*. *Genome* 29(6): 823-827.
- Mercer, K. L., D. S. Murray, et al. (1987). Interference of unicorn plant (*Proboscidea louisianica*) with cotton (*Gossypium hirsutum*). *Weed Science* 35(6): 807-812.
- Meredith, W. R. J. and R. Wells (1987). Sub okra leaf influence on cotton yield. *Crop Science* 27(1): 47-48.

- Mersie, W. and M. Singh (1987). Comparison of norflurazon absorption by excised roots of three plant species. *Pesticide Biochemistry And Physiology* 28(1): 114-120.
- Meyer, W. S., D. C. Reicosky, et al. (1987). A portable chamber for measuring canopy gas exchange of crops subject to different root zone conditions. *Agronomy Journal* 79(1): 181-184.
- Meyer, W. S., D. C. Reicosky, et al. (1987). Physiological responses of cotton to a single waterlogging at high and low nitrogen levels. *Plant And Soil* 102(2): 161-170.
- Moholkar, N. D. and M. V. Thombre (1987). Effects of picking sequences on seed germination, seedling characters and their correlations in cotton. *Journal Of Maharashtra Agricultural Universities* 12(2): 155-157.
- Mshiu, E. P., J. H. Young, et al. (1987). Resistance of cotton varieties to *Heliothis zea*. *Southwestern Entomologist* 12(3): 183-190.
- Narayanan, S. S., P. Singh, et al. (1987). Disruptive selection for earliness in upland cotton. *Indian Journal Of Agricultural Sciences* 57(8): 544-546.
- Narayanan, S. S., P. Singh, et al. (1987). Disruptive selection for genetic improvement of upland cotton. *Indian Journal Of Agricultural Sciences* 57(7): 449-452.
- Nazirov, R. N. (1987). Productivity of the radiomutant cotton cultivars and of their initial forms. *Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni*: 46-48.
- Nehra, D. S., S. K. Varma, et al. (1987). Effect of ascorbic acid on fruiting and yield of cotton. *Indian Journal Of Plant Physiology* 30(4): 429-431.
- Oosterhuis, D. M. (1987). A technique to measure the components of root water potential using screen-caged thermocouple psychrometers. *Plant And Soil* 103(2): 285-288.
- Padaganur, G. M. and M. K. Basavaraj (1987). Fungicidal control of *Alternaria* blight of cotton in transition belt of Karnataka (India). *Indian Journal Of Agricultural Sciences* 57(6): 445-447.
- Patil, V. N. and M. R. Patil (1987). Effect of genotype and harvest date on fungal invasion and deterioration of cotton seed prior to harvest. *P.K.*: 133-138.
- Paul, A. V. N., B. Parshad, et al. (1987). An artificial diet for *Pectinophora gossypiella* (Saund.) and *Earias vitella* Fab. bollworms of cotton. *Indian Journal Of Agricultural Sciences* 57(3): 189-192.
- Pavlovskaya, N. E. (1987). Regular features of changes in oxidoreductase activity in the process of cotton seed formation. *Fiziologiya I Biokhimiya Kul'Turnykh Rastenii* 19(5): 449-455.
- Peeters, M. C., S. Voets, et al. (1987). Nucleolar size at early stages of cotton fiber development in relation to final fiber dimension. *Physiologia Plantarum* 71(4): 436-440.
- Pierce, M. and M. Essenberg (1987). Localization of phytoalexins in fluorescent mesophyll cells isolated from bacterial blight-infected cotton cotyledons and separated from other cells by fluorescence-activated cell sorting. *Physiological And Molecular Plant Pathology* 31(2): 273-290.
- Plaut, Z., M. L. Mayoral, et al. (1987). Effect of altered sink:source ratio on photosynthetic metabolism of source leaves. *Plant Physiology* 85(3): 786-791.
- Pushnik, J. C., G. W. Miller, et al. (1987). Influences of ultraviolet (UV)-blue light radiation on the growth of cotton: II. Photosynthesis, leaf anatomy, and iron reduction. *Journal Of Plant Nutrition* 10(17): 2283-2298.

- Radin, J. W., B. A. Kimball, et al. (1987). Photosynthesis of cotton plants exposed to elevated levels of carbon dioxide in the field. *Photosynthesis Research* 12(3): 191-204.
- Ramalho, F. D. S., W. L. Parrot, et al. (1987). Distribution of the tobacco budworm (*Heliothis virescens* (F.)); Lepidoptera, Noctuidae) eggs within cotton plants. *Ciencia E Cultura* 39(1): 61-65.
- Randhawa, P. S., N. J. Singh, et al. (1987). Bacterial flora of cotton seeds and biocontrol of seedling blight caused by *Xanthomonas campestris* pathovar *malvacearum*. *Seed Science And Technology* 15(1): 65-72.
- Rao, E. S., D. S. Kumari, et al. (1987). Allelopathic potential of *Hyptis suaveolens* Poit. on seed germination of weeds and crops. *Indian Botanical Reporter* 6(2): 77-78.
- Reddy, V. R., D. N. Baker, et al. (1987). Application of GOSSYM to yield decline in cotton: I. Systems analysis of effects of herbicides on growth, development, and yield. *Agronomy Journal* 79(1): 42-47.
- Rodriguez, K. R., H. Ivey, et al. (1987). Peanut-cotton rotations for the management of *Meloidogyne arenaria*. *Journal Of Nematology* 19(4): 484-486.
- Saka, K., F. R. Katterman, et al. (1987). Cell regeneration and sustained division of protoplasts from cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 6(6): 470-472.
- Schaefer, N. L., F. M. Melhuish, et al. (1987). The effect of an intermittent water-table gradient on soil and xylem nitrate in cotton. *Plant And Soil* 97(1): 71-78.
- Seth, S., B. P. S. Lather, et al. (1987). Stability parameters in upland cotton. *Indian Journal Of Agricultural Sciences* 57(6): 429-433.
- Sharopova, N. R., M. F. Abzalov, et al. (1987). Genetic control of esterase, superoxide dismutase and 6-phosphogluconate dehydrogenase in the cotton *Gossypium hirsutum*. *Doklady Akademii Nauk Sssr* 296(4): 994-997.
- Sharopova, N. R., A. A. Pomortsev, et al. (1987). Genetic control of phosphoglucomutase and NADP-dependent malate dehydrogenase in cotton. *Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni*: 7-9.
- Shende, S. T., M. Singh, et al. (1987). Effect of seed bacterization with *Azotobacter chroococcum* on germination and seedling height of upland cotton. *Indian Journal Of Agricultural Sciences* 57(3): 183-185.
- Shinde, Y. M., M. V. Thomre, et al. (1987). Relationship between genetic make-up and phenotypic stability in cotton (*Gossypium hirsutum*). *Indian Journal Of Genetics & Plant Breeding* 47(2): 189-194.
- Shoemaker, R. C., S. E. Christofferson, et al. (1987). Storage protein accumulation patterns in somatic embryos of cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 6(1): 12-15.
- Shukla, U. C. and H. Raj (1987). Influence of genotypical variability on zinc response in cotton (*Gossypium hirsutum* L.). *Plant And Soil* 104(1): 151-154.
- Silberbush, M. and A. J. Ben (1987). The effect of salinity of parameters of potassium and nitrate uptake of cotton. *Communications In Soil Science And Plant Analysis* 18(1): 65-82.
- Singh, H. and P. K. Chhuneja (1987). Performance of high-volume, low-volume and ultra-low-volume sprays for the control of *Amrasca biguttula biguttula* (Ishida) and *Bemisia tabaci* Genn. infesting cotton. *Indian Journal Of Agricultural Sciences* 57(5): 360-364.
- Singh, H. and B. Ram (1987). Effect of different host plants on the development of red cotton bug, *Dysdercus koenigii* (Fab.). *Indian Journal Of Entomology* 49(3): 345-350.

- Singh, K. P., S. N. Bhardwaj, et al. (1987). Parameters controlling crop growth rate and their relationship with photosynthetically active radiation and temperature in upland cotton. *Indian Journal Of Agricultural Sciences* 57(11): 828-832.
- Singh, M., T. H. Singh, et al. (1987). Combining ability analysis for fiber quality characters in upland cotton. *Crop Improvement* 14(2): 136-140.
- Singh, V., S. S. Verma, et al. (1987). Nitrogen use economy through integrated weed control in cotton. *Indian Journal Of Agronomy* 32(3): 226-228.
- Sippell, D. W., O. S. Bindra, et al. (1987). Resistance to whitefly (*Bemisia tabaci*) in cotton (*Gossypium hirsutum*) in the Sudan. *Crop Protection* 6(3): 171-178.
- Slama, F. (1987). Causes of sodium exclusion from leaves of plants sensitive to sodium chloride. *Agronomie* 7(7): 517-522.
- Snipes, C. E., J. E. Street, et al. (1987). Interference periods of common cocklebur (*Xanthium strumarium*) with cotton (*Gossypium hirsutum*). *Weed Science* 35(4): 529-532.
- Soomro, A. W. and S. A. Waring (1987). Effect of temporary flooding on cotton growth and nitrogen nutrition in soils with different organic matter levels. *Australian Journal Of Agricultural Research* 38(1): 91-100.
- Stark, S. B. and F. Whitford (1987). Functional response of *Chrysopa carnea* (Neuroptera: Chrysopidae) larvae feeding on *Heliothis virescens* (Lepidoptera: Noctuidae) eggs on cotton in field cages. *Entomophaga* 32(5): 521-528.
- Stoilova, A. (1987). Correlations between total and commercial productivity in interspecific hybrids of cotton (*Gossypium hirsutum* X *Gossypium barbadense*). *Genetika I Seleksiya* 20(6): 517-521.
- Thaker, V. S., S. Saroop, et al. (1987). Physiological and biochemical changes associated with cotton fiber development: IV. Glycosidases and beta-1,3-glucanase activities. *Annals Of Botany* 60(5): 579-586.
- Thi, A. T. P., F. C. Borrel, et al. (1987). Effects of drought on (1-carbon-14)-oleic and (1-carbon-14)-linoleic acid desaturation in cotton leaves. *Physiologia Plantarum* 69(1): 147-150.
- Thombre, M. V., S. V. Pathade, et al. (1987). Genetic analysis for boll number, boll weight and seed cotton yield. *Journal Of Maharashtra Agricultural Universities* 12(3): 306-308.
- Thomson, N. J., P. E. Reid, et al. (1987). Effects of the okra leaf, nectariless, frego bract and glabrous conditions on yield and quality of cotton lines. *Euphytica* 36(2): 545-554.
- Tiwari, V. N., K. C. Mandloi, et al. (1987). Heterosis and inbreeding depression in intraspecific crosses of upland cotton. *Indian Journal Of Agricultural Sciences* 57(5): 313-317.
- Trolinder, N. L. and J. R. Goodin (1987). Somatic embryogenesis and plant regeneration in cotton (*Gossypium hirsutum* L.). *Plant Cell Reports* 6(3): 231-234.
- Tyagi, A. P. (1987). Correlation studies on yield and fiber traits in upland cotton (*Gossypium hirsutum* L.). *Theoretical And Applied Genetics* 74(2): 280-283.
- Ungar, E. D., D. Wallach, et al. (1987). Cotton response to bud and boll removal. *Agronomy Journal* 79(3): 491-497.
- Usmanov, P. D., I. G. Mednik, et al. (1987). Genotypic specificity in plant responses to middle-wave UV radiation. *Fiziologiya Rastanii* 34(4): 720-729.

- Van, H. H. G., S. W. H. Van, et al. (1987). Release of cotton breeding lines (*Gossypium hirsutum* L.) adapted to South African conditions. *South African Journal Of Plant And Soil* 4(4): 203-205.
- Van, W. R. J., G. C. Prinsloo, et al. (1987). A preliminary survey of plant-parasitic nematode genera in the cotton-producing areas of South Africa. *Phytophylactica* 19(3): 259-260.
- Vavasseur, A., G. Lasceve, et al. (1987). Ion movements during stomatal opening in darkness after illumination under nitrogen. *Physiologia Plantarum* 71(4): 471-476.
- Whitwell, T., S. M. Brown, et al. (1987). Influence of application date on harvest aids for cotton. *Applied Agricultural Research* 2(1): 15-19.
- Wilson, F. D. (1987). Pink bollworm resistance, lint yield, and earliness of cotton isolines in a resistant genetic background. *Crop Science* 27(5): 957-960.
- Wilson, F. D. (1987). Inheritance of pink filament in cotton. *Journal Of Heredity* 78(4): 223-224.
- Wilson, F. D. and R. L. Shepherd (1987). Plant pubescence, genetic background, and seasonal effects on agronomic and fiber properties of upland cotton. *Crop Science* 27(5): 865-868.
- Wilson, R. F., J. J. Burke, et al. (1987). Plant morphological and biochemical responses to field water deficits: II. Responses of leaf glycerolipid composition in cotton. *Plant Physiology* 84(2): 251-254.
- Wullschleger, S. D. and D. M. Oosterhuis (1987). Electron microscope study of cuticular abrasion on cotton leaves in relation to water potential measurements. *Journal Of Experimental Botany* 38(189): 660-667.
- Yatsu, L. Y. and H. W. Kircher (1987). Electron microscopic cytochemistry of cyclopropanoids in cottonseed (*Gossypium hirsutum* L.) tissues. *Lipids* 22(6): 397-400.
- Yokoyama, V. Y., B. E. Mackey, et al. (1987). Relation of cotton protein and tannin to methyl parathion treatment and cigarette beetle (Coleoptera: Anobiidae) growth on foliage. *Journal Of Economic Entomology* 80(4): 834-839.
- Yunuskhonov, S. H. (1987). Isolation of a protein with diaphorase activity from cotton seeds and the study of some properties of the protein. *Khimiya Prirodnikh Soedinenii*(3): 416-421.
- Zade, V. R. and A. M. Dhopte (1987). Anatomy and pigment analysis of red leaf in cotton. *P.K.*: 107-111.
- Zaffaroni, E., M. D. S. Diniz, et al. (1987). Yield stability of sole and intercropping systems in the northeast of Brazil. *Pesquisa Agropecuaria Brasileira* 22(4): 393-400.
- Zeringue, H. J. (1987). A possible relationship between phytoalexin production in the cotton leaf and a phytotoxic response. *Phytochemistry* 26(4): 975-978.
- Zhou, R. H. and X. D. Lin (1987). Isolation of levo gossypol from natural plant. *Acta Pharmaceutica Sinica* 22(8): 603-607.
- Abd, E. G. A. A., T. A. E. El, et al. (1986). Comparative study of some Egyptian and American cotton varieties in relation to hill spacings. *Egyptian Journal Of Agronomy* 11(1-2): 71-78.
- Agnihotri, N. P., K. P. Srivastava, et al. (1986). Relative efficacy of some synthetic pyrethroids and other commonly used insecticides against bollworms and their residues in cotton. *International Journal Of Tropical Agriculture* 4(2): 168-174.
- Ahmad, Z., M. R. Attique, et al. (1986). An estimate of the loss in cotton (*Gossypium hirsutum*) yield in Pakistan attributable to the jassid *Amrasca devastans*. *Crop Protection* 5(2): 105-108.

- Ananthkrishnan, T. N., K. P. Sanjayan, et al. (1986). Host plant preferences in *Cyrtacanthacris ranacea* Stoll in some malvaceous hosts in terms of food utilization. *Proceedings Of The Indian National Science Academy Part B Biological Sciences* 52(3): 351-357.
- Anderson, J. M. (1986). Sucrose release from soybean (*Glycine max* cultivar Ransom II) leaf slices. *Physiologia Plantarum* 66(2): 319-327.
- Bailey, B. A. and F. M. Bourland (1986). The influence of seed quality on response of cotton (*Gossypium hirsutum*) seedlings to the preplant herbicide trifluralin. *Field Crops Research* 13(4): 375-382.
- Bajaj, Y. P. S. and M. S. Gill (1986). Micropropagation and germplasm preservation of cotton (*Gossypium* spp.) through shoot tip and meristem culture. *Indian Journal Of Experimental Biology* 24(9): 581-583.
- Barbosa, F. R., W. A. Moreira, et al. (1986). Toxicity of pesticides and herbicides used on cotton plants to the entomopathogenic fungus *Nomuraea rileyi*. *Pesquisa Agropecuaria Brasileira* 21(6): 641-644.
- Barrow, J. R. (1986). The conditions required to isolate and maintain viable cotton (*Gossypium hirsutum* L.) microspores. *Plant Cell Reports* 5(6): 405-408.
- Beltrao, N. E. D. M., L. B. D. Nobrega, et al. (1986). Intercropping of upland cotton (*Gossypium hirsutum* var. *latifolium*) with cowpea (*Vigna unguiculata* ssp. *unguiculata*) cultivars. *Pesquisa Agropecuaria Brasileira* 21(3): 271-284.
- Beltrao, N. E. D. M., D. J. Vieira, et al. (1986). The importance of planting configuration and the sorghum (*Sorghum bicolor*) cultivar intercropped with annual cotton (*Gossypium hirsutum latifolium*) cultivar CNPA-2H. *Pesquisa Agropecuaria Brasileira* 21(2): 173-186.
- Bhardwaj, S. N. and Kalindi (1986). Relation between photosynthetic rate, translocation of assimilates and boll weight in upland cotton (*Gossypium hirsutum*, Linn.). *Indian Journal Of Plant Physiology* 29(2): 133-136.
- Bhardwaj, S. N., M. Singh, et al. (1986). Leaf size, its thickness and conductance in the genotypes of upland cotton. *Indian Journal Of Agricultural Sciences* 56(12): 840-843.
- Borroto, K. E. and L. Dure, III (1986). The expression of chloroplast genes during cotton (*Gossypium hirsutum*) embryogenesis. *Plant Molecular Biology* 7(2): 105-114.
- Boyette, C. D. and H. L. Walker (1986). Evaluation of *Fusarium lateritium* as a biological herbicide for controlling velvetleaf (*Abutilon theophrasti*) and prickly sida (*Sida spinosa*). *Weed Science* 34(1): 106-109.
- Breen, P. J., J. D. Hesketh, et al. (1986). Field measurements of leaf photosynthesis of 3-carbon pathway and 4-carbon pathway species under high irradiance and enriched carbon dioxide. *Photosynthetica* 20(3): 281-285.
- Burichenko, V. K., T. P. Kopitsya, et al. (1986). Histones from cotton: Molecular characteristics of histone H1. *Khimiya Prirodnikh Soedinenii*(4): 491-497.
- Butler, G. D. J., J. K. Brown, et al. (1986). Effect of cotton (*Gossypium hirsutum*) seedling infection by cotton-leaf crumple virus on subsequent growth and yield. *Journal Of Economic Entomology* 79(1): 208-211.
- Butler, G. D. J., B. A. Kimball, et al. (1986). Populations of *Bemisia tabaci* (Homoptera: Aleyrodidae) on cotton (*Gossypium hirsutum* cultivar Deltapine 61) grown in open-top field chambers enriched with carbon dioxide. *Environmental Entomology* 15(1): 61-63.
- Cakmak, I. and H. Marschner (1986). Mechanism of phosphorus-induced zinc deficiency in cotton (*Gossypium hirsutum* cultivar Deltapine 15/21): I. Zinc deficiency-enhanced uptake rate of phosphorus. *Physiologia Plantarum* 68(3): 483-490.

- Canuto, V. T. B., N. N. Canuto, et al. (1986). Evaluation of the quality of cotton seeds stored in different regions of Pernambuco, Brazil. *Pesquisa Agropecuaria Brasileira* 21(8): 793-799.
- Carmi, A. (1986). Effects of root zone volume and plant density on the vegetative and reproductive development of cotton (*Gossypium hirsutum*). *Field Crops Research* 13(1): 25-32.
- Chakravarthy, A. K. and A. S. Sidhu (1986). Resistance to insect pest damage in four cotton varieties in Ludhiana(India). *Insect Science And Its Application* 7(5): 647-652.
- Chang, C. W. and J. A. Wetmore (1986). Effects of water stress on starch and its metabolizing enzyme activities in cotton (*Gossypium hirsutum* cultivar Coker 100) leaves. *Starch Staerke* 38(9): 311-313.
- Chenault, E. W., A. F. Wiese, et al. (1986). An economic analysis of incorporation methods for preplant herbicides on clay loam soil. *Weed Science* 34(3): 419-422.
- Chlan, C. A., J. B. Pyle, et al. (1986). Developmental biochemistry of cottonseed embryogenesis and germination XVIII Complementary DNA and amino acid sequences of members of the storage protein families. *Plant Molecular Biology* 7(6): 475-490.
- Cohen, M. L. and W. D. Steinmetz (1986). Foliar washoff of pesticides by rainfall. *Environmental Science & Technology* 20(5): 521-523.
- Correia, J. S. and J. D. Vendramim (1986). Influence of cotton cultivars on the biology of *Heliothis virescens* (Lepidoptera-Noctuidae). *Anais Da Sociedade Entomologica Do Brasil* 15(1): 47-60.
- Cramer, G. R., A. Lauchli, et al. (1986). Effects of sodium chloride and calcium chloride on ion activities in complex nutrient solutions and root growth of cotton (*Gossypium hirsutum*). *Plant Physiology* 81(3): 792-797.
- D'Aoust, A. L. and C. Hubac (1986). Effect of phytochrome state on solute leakage from leaf discs of cotton plants. *Canadian Journal Of Botany* 64(12): 2973-2975.
- Davidoff, B. and H. M. Selim (1986). Goodness of fit for eight water infiltration models. *Soil Science Society Of America Journal* 50(3): 759-764.
- Davies, E. A. and N. J. J. Combrink (1986). Effect of calcium, potassium and boron supply on water stress induced chlorophyll loss from tobacco (*Nicotiana tabacum*) and cotton (*Gossypium hirsutum*) leaf discs. *South African Journal Of Plant And Soil* 3(1): 51-53.
- Devey, M. E. and A. A. Rosielle (1986). Relationship between field and greenhouse ratings for tolerance to verticillium wilt on cotton (*Gossypium hirsutum*). *Crop Science* 26(1): 1-4.
- Dilday, R. H. (1986). Development of a cotton plant with glandless seeds, and glanded foliage and fruiting forms. *Crop Science* 26(3): 639-641.
- Dugger, W. M. and R. L. Palmer (1986). Incorporation of UDPglucose into cell wall glucans and lipids by intact cotton (*Gossypium hirsutum* cultivar Acala SJ1) fibers. *Plant Physiology* 81(2): 464-470.
- Echekwu, C. A. and A. T. Obilana (1986). Genetics of some agronomic and fiber traits in upland cotton. *Samaru Journal Of Agricultural Research* 4(1-2): 63-70.
- Elzen, G. W., H. J. Williams, et al. (1986). Wind tunnel flight responses by hymenopterous parasitoid *Campoletis sonorensis* to cotton cultivars and lines. *Entomologia Experimentalis Et Applicata* 42(3): 285-290.
- Erdman, M. D. and W. G. Monson (1986). In vitro digestibility of selected agricultural wastes at various moisture levels treated with microwave energy. *Journal Of Agricultural And Food Chemistry* 34(5): 889-892.

- Flint, H. M., N. J. Cutice, et al. (1986). A comparison of related nectaried and nectariless cottons for control of the pink bollworm in field plots treated with gossypure, insecticides, or untreated. *Journal Of Agricultural Entomology* 3(4): 362-368.
- Francl, L. J. and V. H. Dropkin (1986). *Heterodera glycines* population dynamics and relation of initial population to soybean (*Glycine max* cultivar Forrest) yield. *Plant Disease* 70(8): 791-795.
- Gafurov, A. T., S. S. Faizullaev, et al. (1986). Autocompetition and economical by valuable traits in cotton. *Sel'Skokhozyaistvennaya Biologiya*(5): 50-54.
- Galau, G. A., D. W. Hughes, et al. (1986). Abscisic acid induction of cloned cotton (*Gossypium hirsutum*) late embryogenesis-abundant (Lea) messenger RNA species. *Plant Molecular Biology* 7(3): 155-170.
- Gawel, N. J., A. P. Rao, et al. (1986). Somatic embryogenesis from leaf and petiole callus cultures of *Gossypium hirsutum* L. *Plant Cell Reports* 5(6): 457-459.
- Gawel, N. J. and C. D. Robacker (1986). Effect of pollen-style interaction on the pollen tube growth of *Gossypium hirsutum*. *Theoretical And Applied Genetics* 72(1): 84-87.
- Gazyants, S. M. and D. M. Laiskhram (1986). Genetic analysis of photosynthetic processes during interspecific cotton hybridization. *Sel'Skokhozyaistvennaya Biologiya*(5): 72-75.
- Gigou, J. (1986). Nutrient balance under a sorghum-cotton rotation in lysimeters. *Agronomie Tropicale Riz Et Riziculture Et Cultures Vivrieres Tropicales* 41(2): 149-153.
- Gould, J. H., R. L. Palmer, et al. (1986). Isolation and culture of cotton (*Gossypium hirsutum* cultivar Acala SJ-2) ovule epidermal protoplasts (prefiber cells) and analysis of the regenerated wall. *Plant Cell Tissue And Organ Culture* 6(1): 47-60.
- Gould, J. H., R. L. Palmer, et al. (1986). Isolation and culture of anucleate protoplasts from cotton (*Gossypium hirsutum*) fiber; assessment of viability and analysis of regenerated wall polymers. *Plant Cell Tissue And Organ Culture* 6(1): 61-72.
- Gulin, V. V., V. A. Kal'Chenko, et al. (1986). Biochemical polymorphism in inbred lines of cotton (*Gossypium hirsutum* L.). *Genetika* 22(8): 2152-2158.
- Gunaseelan, T. and R. Krishnaswami (1986). Genetic analysis of characters associated with yield in *Gossypium hirsutum* L. *International Journal Of Tropical Agriculture* 4(3): 258-267.
- Gunaseelan, T. and R. Krishnaswami (1986). Genetic analysis of some useful characters in *Gossypium hirsutum* L. *International Journal Of Tropical Agriculture* 4(3): 268-275.
- Gupta, S. P. and T. H. Singh (1986). Combining ability in the F-1 and advanced generations of a diallel cross in upland cotton (*Gossypium hirsutum* L.). *Indian Journal Of Genetics & Plant Breeding* 46(2): 345-347.
- Gupta, S. P. and T. H. Singh (1986). Diallel analysis of earliness and plant height in upland cotton. *Crop Improvement* 13(2): 122-125.
- Heagle, A. S., W. W. Heck, et al. (1986). Injury and yield response of cotton (*Gossypium hirsutum*) to chronic doses of ozone and sulfur dioxide. *Journal Of Environmental Quality* 15(4): 375-382.
- Hearn, A. B. (1986). Effect of preceding crop on the nitrogen requirements of irrigated cotton (*Gossypium hirsutum*) on a vertisol. *Field Crops Research* 13(2): 159-175.
- Hendrix, D. L. and S. C. Huber (1986). Diurnal fluctuations in cotton (*Gossypium hirsutum* cultivar Coker 100) leaf carbon export, carbohydrate content, and sucrose synthesizing enzymes. *Plant Physiology* 81(2): 584-586.

- Higgins, J. M., R. H. Walker, et al. (1986). Coffee senna (*Cassia occidentalis*) competition with cotton (*Gossypium hirsutum*). *Weed Science* 34(1): 52-56.
- Hons, F. M. and B. L. McMichael (1986). Planting pattern effects on yield, water use and root growth of cotton (*Gossypium hirsutum*). *Field Crops Research* 13(2): 147-158.
- Hopper, K. R. and E. G. King (1986). Linear functional response of *Microplitis croceipes* (Hymenoptera: Braconidae) to variation in *Heliothis* spp. (Lepidoptera: Noctuidae) density in the field. *Environmental Entomology* 15(3): 476-480.
- Hubac, C., D. Guerrier, et al. (1986). Effect of far-red light on malate and potassium contents in cotton (*Gossypium hirsutum* cultivar Bou) leaves: Relation to drought resistance. *Physiologia Plantarum* 66(1): 37-40.
- Hue, N. V., G. R. Craddock, et al. (1986). Effect of organic acids on aluminum toxicity in subsoils. *Soil Science Society Of America Journal* 50(1): 28-34.
- Imamaliev, A. I., P. Y. Popova, et al. (1986). Changes in structural polysaccharide content in the cell wall of the cotton fiber and the formation of the physicommechanical properties of the fiber. *Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni:* 18-20.
- Isamil, F. M. and E. F. A. Al (1986). Comparative study for the relative importance of characters contributing to seed-cotton yield in American cotton (*Gossypium hirsutum* cultivar McNair 220) and Egyptian cotton (*Gossypium barbadense* cultivar Giza 75). *Journal Of Agronomy And Crop Science* 156(2): 128-132.
- Jackson, B. S. and G. F. Arkin (1986). Quantifying cultivar differences in cotton (*Gossypium hirsutum*). *Field Crops Research* 14(1): 75-86.
- Jagtap, D. R. (1986). Combining ability in upland cotton. *Indian Journal Of Agricultural Sciences* 56(12): 833-840.
- Janat, M. M. and J. L. Stroehlein (1986). Yield, dry matter production and phosphorus uptake of drip irrigated cotton (*Gossypium hirsutum* cultivar DPL 90). *Journal Of Fertilizer Issues* 3(4): 124-128.
- Jenkins, J. N., W. L. Parrott, et al. (1986). Performance of cottons (*Gossypium hirsutum*) when infested with tobacco budworm (*Heliothis virescens*). *Crop Science* 26(1): 93-95.
- Johnson, L. F. and J. H. Doyle (1986). Relationships of seedling disease of cotton (*Gossypium hirsutum*) to characteristics of loessial soils in Tennessee (USA). *Phytopathology* 76(3): 286-290.
- Kalsy, H. S., H. R. Garg, et al. (1986). Semigraphical analysis in upland cotton (*Gossypium hirsutum* L.). *Crop Improvement* 13(2): 186-188.
- Karev, K. S. (1986). Parameters of photosynthetic potential and net productivity in cotton (*Gossypium hirsutum* cultivar) Chirpan 433 grown on leached cinnamon forest soil. *Rasteniyev'Dni Nauki* 23(2): 51-55.
- Karim, Z., M. S. Khan, et al. (1986). Effect of permeability on the shoot and root growth of cotton (*Gossypium hirsutum*). *Bangladesh Journal Of Agriculture* 11(3): 39-46.
- Keeley, P. E., R. J. Thullen, et al. (1986). Influence of planting date on growth of ivyleaf morning glory (*Ipomoea hederacea*) in cotton (*Gossypium hirsutum*). *Weed Science* 34(6): 906-910.
- Kennedy, C. W., W. C. J. Smith, et al. (1986). Effect of early season square removal on three leaf types of cotton (*Gossypium hirsutum*). *Crop Science* 26(1): 139-145.

- Kerbabaeva, Z. A., L. N. Ivonina, et al. (1986). Salt resistance of cultivars and lines from the genetic collection of cotton. *Izvestiya Akademii Nauk Turkmenskoi Ssr Seriya Biologicheskikh Nauk*(4): 8-13.
- Kerby, T. A., K. Hake, et al. (1986). Cotton (*Gossypium hirsutum*) fruiting modification with mepiquat chloride. *Agronomy Journal* 78(5): 907-912.
- Khan, Y. A., S. H. Hussaini, et al. (1986). Effects of seed size on crop growth and yield of cotton (*Gossypium* spp. L.). *Seed Research* 14(2): 144-155.
- Kishor, P. B. K. and A. R. Mehta (1986). The influence of applied carbohydrates on the growth and amylase activity in cultured cotton cells. *Indian Botanical Reporter* 5(1): 19-23.
- Kittock, D. L., R. A. Selley, et al. (1986). Plant population and plant height effects on pima cotton (*Gossypium barbadense*) lint yield. *Agronomy Journal* 30: 534-538.
- Kleifeld, Y., T. Blumenfeld, et al. (1986). Control of Johnsongrass (*Sorghum halepense*) in cotton (*Gossypium hirsutum*) fields. *Phytoparasitica* 14(1): 55-62.
- Klich, M. A. (1986). Presence of *Aspergillus flavus* in developing cotton (*Gossypium hirsutum*) bolls and its relation to contamination of mature seeds. *Applied And Environmental Microbiology* 52(4): 963-965.
- Kurdov, M. (1986). Prognosis of the dynamics of the population size of the bollworm *Helicoverpa armigera* Hbn. (*Heliothis armigera* Hbn.) in the Turkmen SSR (USSR). *Izvestiya Akademii Nauk Turkmenskoi Ssr Seriya Biologicheskikh Nauk*(3): 21-26.
- Kurtz, M. E. (1986). Control of *Sorghum halepense* in cotton (*Gossypium hirsutum*) with post-emergence herbicides and trifluralin. *Weed Research* 26(2): 83-88.
- Lee, J. A. (1986). Effects of boll pilosity on some traits on cultivar pima cotton (*Gossypium barbadense*). *Crop Science* 26(4): 741-743.
- Lee, S. M., N. A. Garas, et al. (1986). High-performance liquid chromatographic determination of sesquiterpenoid stress metabolites in *Verticillium dahliae* infected cotton stele. *Journal Of Agricultural And Food Chemistry* 34(3): 490-493.
- Lieth, J. H., G. F. Arkin, et al. (1986). Modeling cotton (*Gossypium hirsutum*) fruiting form abscission. *Agronomy Journal* 78(4): 730-735.
- Loper, G. M. (1986). Cotton pollen: Honeybee (*Apis mellifera*) avoidance and absence of gossypol. *Journal Of Economic Entomology* 79(1): 103-106.
- Magnuson, C. E., J. D. Goeschl, et al. (1986). Experimental tests of the Muench-Horwitz theory of phloem transport: Effects of loading rates. *Plant Cell And Environment* 9(2): 103-110.
- Marquette, J. (1986). Evaluation of the effect of a previous groundnut crop in Togo. *Agronomie Tropicale Riz Et Riziculture Et Cultures Vivrieres Tropicales* 41(3-4): 231-241.
- Marschner, H. and I. Cakmak (1986). Mechanism of phosphorus-induced zinc deficiency in cotton (*Gossypium hirsutum* cultivar Deltapine 15/21): II. Evidence for impaired shoot control of phosphorus uptake and translocation under zinc deficiency. *Physiologia Plantarum* 68(3): 491-496.
- Mavi, G. S. and A. S. Sidhu (1986). Bio-assay technique of screening cotton (*Gossypium hirsutum*) germ-plasm for jassid resistance. *Indian Journal Of Agricultural Sciences* 56(4): 286-289.
- McCall, L. L., L. M. Verhalen, et al. (1986). Multidirectional selection for fiber strength in upland cotton (*Gossypium hirsutum*). *Crop Science* 26(4): 744-750.

- McCarty, J. C. J., J. N. Jenkins, et al. (1986). Yield response of two cotton (*Gossypium hirsutum*) cultivars to tobacco budworm (*Heliothis virescens*) infestation. *Crop Science* 26(1): 136-139.
- Mellon, J. E. (1986). Some characteristics of peroxidase secreted by cotton (*Gossypium hirsutum*) ovule cultures. *Plant Cell Reports* 5(5): 338-341.
- Menzel, M. Y., C. A. Hasenkampf, et al. (1986). Characteristics of duplications and deficiencies from chromosome translocations in *Gossypium hirsutum*. *Journal Of Heredity* 77(3): 189-201.
- Menzel, M. Y. and K. L. Richmond (1986). A revised map of chromosome 15 in *Gossypium hirsutum*. *Canadian Journal Of Genetics And Cytology* 28(2): 272-277.
- Meredith, W. R. J. and R. Wells (1986). Normal vs. okra leaf yield interactions in cotton (*Gossypium hirsutum*): I. Performance of near-isogenic lines from bulk populations. *Crop Science* 26(2): 219-222.
- Mert, H. H. (1986). Effects of oxygen in air and sodium chloride in medium on photosynthesis and photorespiration in two cotton cultivars. *Biotronics* 15: 1-8.
- Meshram, M. K. and S. Raj (1986). Reaction of the germ-plasm of *Gossypium hirsutum* to bacterial blight. *Indian Journal Of Agricultural Sciences* 56(4): 236-237.
- Misra, R. K., A. R. Dexter, et al. (1986). Maximum axial and radial growth pressure of plant roots. *Plant And Soil* 95(3): 315-326.
- Misra, R. K., A. R. Dexter, et al. (1986). Penetration of soil aggregates of finite size: II. Plant roots. *Plant And Soil* 94(1): 59-86.
- Nastasi, P., R. Frans, et al. (1986). Economics and new alternatives in cotton (*Gossypium hirsutum*) weed management programs. *Weed Science* 34(4): 634-638.
- Nazirov, N. N. and S. S. Kozubaev (1986). The frequency of occurrence of early maturing large-fruited cotton forms of varying origin under the action of radioactive phosphorus. *Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni*: 12-13.
- Nightingale, H. I., K. R. Davis, et al. (1986). Trickle irrigation of cotton (*Gossypium hirsutum*): Effect on soil chemical properties. *Agricultural Water Management* 11(2): 159-168.
- Oosterhuis, D. M. and H. H. Wiebe (1986). Water stress preconditioning and cotton (*Gossypium hirsutum*) root pressure-flux relationships. *Plant And Soil* 95(1): 69-76.
- Ouedraogo, M., C. Hubac, et al. (1986). Effect of far red light on root growth and on xylem sap in cotton (*Gossypium hirsutum* cultivar Bou). *Plant And Cell Physiology* 27(1): 17-24.
- Owen, W. J. and B. Donzel (1986). Oxidative degradation of chlortoluron, propiconazole and metalaxyl in suspension cultures of various crop plants. *Pesticide Biochemistry And Physiology* 26(1): 75-89.
- Pavlovkin, J., A. Novacky, et al. (1986). Membrane potential changes during bacteria-induced hypersensitive reaction. *Physiological And Molecular Plant Pathology* 28(1): 125-136.
- Percy, R. G. (1986). Effects of environment upon ovule abortion in interspecific F-1 hybrids and single species cultivars of cotton. *Crop Science* 26(5): 938-942.
- Prakash, S. and H. C. Kapoor (1986). In vitro studies of nitrate reductase in cotton (*Gossypium hirsutum* cultivar 9-40-54) leaves: Role of light, carbohydrate and cytokinin in its regulation. *Indian Journal Of Biochemistry And Biophysics* 23(3): 143-147.
- Radin, J. W. and M. P. Eidenbock (1986). Carbon accumulation during photosynthesis in leaves of nitrogen- and phosphorus-stressed cotton (*Gossypium hirsutum*). *Plant Physiology* 82(3): 869-871.

- Radin, J. W. and D. L. Hendrix (1986). Accumulation and turnover of abscisic acid in osmotically stressed cotton (*Gossypium hirsutum* cultivar Deltapine 70) leaf tissue in relation to temperature. *Plant Science* 45(1): 37-42.
- Radin, J. W., J. R. Mauney, et al. (1986). Diurnal osmotic cycling in cotton (*Gossypium hirsutum*) leaves as an indicator of source-sink balance. *Plant Cell And Environment* 9(4): 349-352.
- Ramalho, F. S. and F. M. M. D. Jesus (1986). Chemical control of cotton boll weevil, *Anthonomus grandis* Boheman, 1843 (Coleoptera: Curculionidae). *Anais Da Sociedade Entomologica Do Brasil* 15(2): 335-342.
- Ramalho, F. S., F. M. M. D. Jesus, et al. (1986). Evaluation of acaricides against two-spotted spider mite (*Tetranychus urticae* Koch, 1836) on cotton. *Anais Da Sociedade Entomologica Do Brasil* 15(2): 247-256.
- Randhawa, L. S., G. S. Chahal, et al. (1986). Role of epistasis in the inheritance of yield and its components in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 56(7): 494-496.
- Regier, C., R. E. Dilbeck, et al. (1986). Cotton (*Gossypium hirsutum*) resistance to 2,4-D spray drift. *Crop Science* 26(2): 376-377.
- Rhyne, C. L., D. M. Rhyne, et al. (1986). Duplicate pollen-sterile locus ms -8 and linkage group V of *Gossypium hirsutum*. *Journal Of Heredity* 77(5): 332-336.
- Richa, A. S. Basra, et al. (1986). Hydrolytic enzymes in cotton ovules during early development. *Proceedings Of The Indian Academy Of Sciences Plant Sciences* 96(1): 49-54.
- Robbins, C. W. (1986). Sodic calcareous soil reclamation as affected by different amendments and crops. *Agronomy Journal* 78(5): 916-920.
- Robbins, C. W. (1986). Carbon dioxide partial pressure in lysimeter soils. *Agronomy Journal* 78(1): 151-158.
- Rodriguez, G. B. and J. R. Barrow (1986). Short-term storage of cotton (*Gossypium hirsutum*) pollen. *Plant Cell Reports* 5(5): 332-333.
- Rogers, C. B., R. Talbert, et al. (1986). Effect of cotton (*Gossypium hirsutum*) herbicide carryover on subsequent crops. *Weed Science* 34(5): 756-760.
- Rogers, C. B., R. E. Talbert, et al. (1986). Residual fluometuron levels in three Arkansas soils under continuous cotton (*Gossypium hirsutum*) production. *Weed Science* 34(1): 122-130.
- Sangwan, R. S. and J. S. Yadava (1986). Heterosis in upland cotton (*Gossypium hirsutum*) and its commercial utilization. *Indian Journal Of Agricultural Sciences* 56(11): 765-768.
- Santana, J. C. F. D., E. O. D. Santos, et al. (1986). Evaluation of herbaceous cotton (*Gossypium hirsutum* latifolium) lines and cultivars in the northeast of Brazil: I. Yield. *Pesquisa Agropecuaria Brasileira* 21(1): 43-52.
- Sarsar, S. M., R. A. Patil, et al. (1986). Heterosis and combining ability in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 56(8): 567-573.
- Savoy, S. G. (1986). Comparative resistance of Bulgarian and foreign cultivars to Verticillium wilt. *Rasteniyev'Dni Nauki* 23(5): 63-67.
- Schmidt, J. H. and R. Wells (1986). Recovery of soluble proteins from glanded cotton tissues with amines. *Analytical Biochemistry* 154(1): 244-249.
- Sengonca, C., M. S. Lababidi, et al. (1986). The effects of different cotton varieties on the carmine spider mite, *Tetranychus cinnabarinus* Bois. (Acari: Tetranychidae). *Plant Breeding* 97(4): 297-303.

- Shamina, Z. B., A. M. Turaev, et al. (1986). Callus-forming capacity in cultivated cotton anthers. *Tsitologiya I Genetika* 20(4): 276-280.
- Sheikh, A. S., V. P. Singh, et al. (1986). Transpiration suppression in young cotton plants by flurenol (EMD IT 3233). *Egyptian Journal Of Botany* 30(1-3): 265-272.
- Shepherd, R. L. (1986). Cotton (*Gossypium hirsutum*) resistance to the root knot-fusarium wilt complex: II. Relation to root-knot resistance and its implications on breeding for resistance. *Crop Science* 26(2): 233-237.
- Shepherd, R. L. and A. J. Kappelman (1986). Cotton (*Gossypium hirsutum*) resistance to the root knot-fusarium wilt complex: I. Relation to fusarium wilt resistance and its implications on breeding for resistance. *Crop Science* 26(2): 228-232.
- Shinde, Y. M., M. V. Thombre, et al. (1986). Stability for lint index and seed index in rainfed cotton. *Journal Of Maharashtra Agricultural Universities* 11(3): 318-321.
- Shoemaker, R. C., L. J. Couche, et al. (1986). Characterization of somatic embryogenesis and plant regeneration in cotton (*Gossypium hirsutum*). *Plant Cell Reports* 5(3): 178-181.
- Sidhu, A. S. and A. K. Dhawan (1986). Bollworm damage and carryover in *Gossypium hirsutum* cotton in Punjab, (India). *Journal Of Research Punjab Agricultural University* 23(3): 444-450.
- Sidhu, A. S. and A. K. Dhawan (1986). Estimation of loss due to the attack of cotton jassid (*Amrasca biguttula biguttula* (Ishida)) on *Gossypium hirsutum* cotton. *Indian Journal Of Plant Protection* 14(1): 45-50.
- Singh, B., G. S. Chahal, et al. (1986). Use of parental performance for the prediction of heterosis on cotton (*Gossypium hirsutum*). *Crop Improvement* 13(1): 29-33.
- Singh, D. (1986). Effect of low light intensity on growth and yield of rainfed cotton. *Indian Journal Of Plant Physiology* 29(3): 230-236.
- Singh, J., S. S. Sandhu, et al. (1986). Role of various mortality factors in reducing the carryover of pink bollworm, *Pectinophora gossypiella* (Saunders) of cotton through different sources in Punjab (India). *Indian Journal Of Ecology* 13(2): 291-300.
- Singh, J. P., B. R. Mor, et al. (1986). Diapausing of pink bollworm (*Pectinophora gossypiella*) in relation to the time and formation of cotton bolls (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 56(3): 198-199.
- Singh, M., V. P. Singh, et al. (1986). Improvement of yield and quality in upland cotton (*Gossypium hirsutum* cultivar Pusa 595B) through progeny bulk selection. *Indian Journal Of Agricultural Sciences* 56(8): 562-566.
- Singh, R. and R. A. Agarwal (1986). Fumigation in controlling the carry-over of pink bollworm (*Pectinophora gossypiella*) in cotton under rural conditions. *Indian Journal Of Agricultural Sciences* 56(11): 800-802.
- Singh, T. H. and S. S. Gill (1986). Genotype times environment interaction in upland cotton (*Gossypium hirsutum* L.). *Crop Improvement* 13(2): 215-217.
- Smith, C. W., J. T. Cothren, et al. (1986). Yield and fiber quality of cotton (*Gossypium hirsutum*) following application of 2-chloroethylphosphonic acid. *Agronomy Journal* 78(5): 814-818.
- Smith, G. S. and R. W. Roncadori (1986). Responses of three vesicular-arbuscular mycorrhizal fungi at four soil temperatures and their effects on cotton growth. *New Phytologist* 104(1): 89-96.
- Starr, J. L. and J. A. Veech (1986). Susceptibility to root-knot nematodes (*Meloidogyne incognita*) in cotton (*Gossypium hirsutum*) lines resistant to the fusarium wilt/root-knot complex. *Crop Science* 26(3): 543-546.

- Stoilova, A. and N. Savova (1986). Growth and development study of some wild cotton species and possibilities for their hybridization with the cultivated species *Gossypium hirsutum*. *Rasteniev'Dni Nauki* 23(5): 5-9.
- Sun, C. W. and Z. I. Liang (1986). Esterase isozyme analyses of the species and their varieties of the genus *Gossypium*. *Acta Botanica Sinica* 28(3): 263-269.
- Suttle, J. C. (1986). Cytokinin-induced ethylene biosynthesis in nonsenescing cotton leaves. *Plant Physiology* 82(4): 930-935.
- Suttle, J. C. and E. R. Mansager (1986). The physiological significance of phenylacetic acid in abscising cotton (*Gossypium hirsutum* cultivar LG 102) cotyledons. *Plant Physiology* 81(2): 434-438.
- Taneja, S. L. and A. P. Jayaswal (1986). Population dynamics of pink bollworm, *Pectinophora gossypiella* on upland cotton (*Gossypium hirsutum*). *Insect Science And Its Application* 7(4): 569-574.
- Taylor, G. E. J. and C. A. Gunderson (1986). The response of foliar gas exchange to exogenously applied ethylene. *Plant Physiology* 82(3): 653-657.
- Temple, P. J. (1986). Stomatal conductance and transpirational responses of field-grown cotton (*Gossypium hirsutum*) to ozone. *Plant Cell And Environment* 9(4): 315-322.
- Thaker, V. S., S. Saroop, et al. (1986). Physiological and biochemical changes associated with cotton (*Gossypium hirsutum* cultivar SRT I) fiber development: III. Indolyl-3-acetaldehyde dehydrogenase. *Biochemie Und Physiologie Der Pflanzen* 181(5): 339-345.
- Thaker, V. S., S. Saroop, et al. (1986). Role of peroxidase and esterase activities during cotton (*Gossypium hirsutum* cultivar SRT-1) fiber development. *Journal Of Plant Growth Regulation* 5(1): 17-28.
- Thengane, S., S. V. Paranjpe, et al. (1986). Hybridization of *Gossypium* species through in ovulo embryo culture. *Plant Cell Tissue And Organ Culture* 6(3): 209-220.
- Timpa, J. D., J. J. Burke, et al. (1986). Effects of water stress on the organic acid and carbohydrate compositions of cotton (*Gossypium hirsutum*) plants. *Plant Physiology* 82(3): 724-728.
- Touchton, J. T., D. H. Rickerl, et al. (1986). Starter fertilizer combinations and placement for conventional and no-tillage cotton (*Gossypium hirsutum*). *Journal Of Fertilizer Issues* 3(3): 91-98.
- Turaev, A. M. and Z. B. Shamina (1986). Optimization of the medium for the cultivation of cotton (*Gossypium hirsutum*) anthers in vitro. *Fiziologiya Rastenii* 33(3): 565-571.
- Turcotte, E. L. (1986). Round leaf-3 mutant in American pima cotton (*Gossypium barbadense*). *Journal Of Heredity* 77(5): 364-366.
- Turner, N. C., A. B. Hearn, et al. (1986). Cotton (*Gossypium hirsutum*): Physiological and morphological responses to water deficits and their relationship to yield. *Field Crops Research* 14(2): 153-170.
- Tyagi, A. P. (1986). Association analysis for yield components in upland cotton (*Gossypium hirsutum* L.). *East African Agricultural And Forestry Journal* 52(1): 37-40.
- Tyagi, A. P. (1986). Variability and heritability for yield and its components in upland cotton (*Gossypium hirsutum*). *East African Agricultural And Forestry Journal* 52(1): 41-46.
- Vesmanova, O. Y., O. I. Otakhanov, et al. (1986). Cytogenetic consequences of using the defoliant butiphos in cotton over many years. *Tsitologiya I Genetika* 20(4): 267-269.
- Vlasova, N. A. and V. A. Krakhmalev (1986). Features of growth of the apical part of cottonseed fibers. *Doklady Akademii Nauk Sssr* 288(4): 980-983.

- Wallace, A. and G. A. Wallace (1986). Effects of soil conditioners on emergence and growth of tomato (*Lycopersicon esculentum* cultivar Tropic), cotton (*Gossypium hirsutum* cultivar Acala), and lettuce (*Lactuca sativa* cultivar Clima x) seedlings. *Soil Science* 141(5): 313-316.
- Wanjura, D. F. and J. D. J. Bilbro (1986). Ground cover and weathering effects on reflectances of three crop residues. *Agronomy Journal* 78(4): 694-698.
- Wells, R. and W. R. J. Meredith (1986). Normal vs. okra leaf yield interactions in cotton (*Gossypium hirsutum*): II. Analysis of vegetative and reproductive growth. *Crop Science* 26(2): 223-228.
- Wells, R., W. R. J. Meredith, et al. (1986). Canopy photosynthesis and its relationship to plant productivity in near-isogenic cotton (*Gossypium hirsutum*) lines differing in leaf morphology. *Plant Physiology* 82(3): 635-640.
- Wiese, A. F. and E. W. Chenault (1986). Incorporating herbicides on sandy soil for dryland cotton (*Gossypium hirsutum*). *Agronomy Journal* 78(5): 897-900.
- Willis, G. H., L. L. McDowell, et al. (1986). Permethrin washoff from cotton (*Gossypium hirsutum*) plants by simulated rainfall. *Journal Of Environmental Quality* 15(2): 116-120.
- Wilson, F. D. (1986). Inheritance of ephemeral leaf mutant of upland cotton (*Gossypium hirsutum*). *Crop Science* 26(3): 498-500.
- Wilson, F. D. and B. W. George (1986). Smoothleaf and hirsute cottons (*Gossypium hirsutum*): Response to insect pests and yield in Arizona (USA). *Journal Of Economic Entomology* 79(1): 229-232.
- Wong, S. C. and K. C. Woo (1986). Simultaneous measurements of steady state chlorophyll a fluorescence and carbon dioxide assimilation in leaves: The relationship between fluorescence and photosynthesis in 3-carbon photosynthetic pathway and 4-carbon photosynthetic pathway plants. *Plant Physiology* 80(4): 877-883.
- Wullschlegel, S. D. and D. M. Oosterhuis (1986). A rapid leaf-disc sampler for psychrometric water potential measurements. *Plant Physiology* 81(2): 684-685.
- Yatsu, L., T. J. Jacks, et al. (1986). Chemical and microscopic studies of the matrix substance in pigment glands of cotton (*Gossypium hirsutum*) seeds. *Journal Of The American Oil Chemists' Society* 63(4): 534-537.
- Yunuskhonov, S. H. and B. D. Dzhililov (1986). Study of water-soluble proteins from the seeds of cotton *Gossypium hirsutum* and *Gossypium barbadense*. *Khimiya Prirodnikh Soedinenii*(4): 488-491.
- Ackerson, R. C. (1985). Osmoregulation in cotton (*Gossypium hirsutum*) in response to water stress: 3. Effects of phosphorus fertility. *Plant Physiology* 77(2): 309-312.
- Armbrust, D. V. and L. Lyles (1985). Equivalent wind-erosion protection from selected growing crops. *Agronomy Journal* 77(5): 703-707.
- Azevedo, D. M. D., L. B. D. Nobrega, et al. (1985). Effect of height of pruning on herbaceous cotton ratoon production in the humid valleys in northeast Brazil. *Pesquisa Agropecuaria Brasileira* 20(2): 207-214.
- Azevedo, D. M. P. D., N. E. D. Beltrao, et al. (1985). Effect of corn (*Zea mays*) and cowpea (*Vigna unguiculata*) planting date on perennial cotton (*Gossypium hirsutum*) association. *Pesquisa Agropecuaria Brasileira* 20(7): 821-830.
- Bajaj, Y. P. S. and M. S. Gill (1985). In vitro induction of genetic variability in cotton (*Gossypium* spp.). *Theoretical And Applied Genetics* 70(4): 363-368.

Balasubramanian, G., S. Chelliah, et al. (1985). Consumption of food and its utilization by *Spodoptera litura* fed on 8 different host plants. *Indian Journal Of Agricultural Sciences* 55(3): 193-200.

Barker, G. L., J. L. Hatfield, et al. (1985). Cotton phenology parameters affected by wind. *Field Crops Research* 12(1): 33-48.

Bayles, M. B. and W. M. Johnson (1985). Growth of homologous and heterologous phytopathogenic bacteria in cotton lines susceptible, resistant or immune to blight. *Phytopathology* 75(5): 515-519.

Beach, R. M., J. W. Todd, et al. (1985). Nectaried and nectariless cotton (*Gossypium hirsutum*) cultivars as nectar sources for the adult soybean looper (*Pseudoplusia includens*). *Journal Of Entomological Science* 20(2): 233-236.

Bell, M. R. and C. L. Romine (1985). Microbial control of *Heliothis* spp. (Lepidoptera: Noctuidae) in cotton (*Gossypium hirsutum*): Dosage and management trials. *Journal Of Entomological Science* 20(2): 146-151.

Bharadwaj, D. N. and N. G. Simongulyan (1985). Genetic analysis of harvest index in upland cotton (*Gossypium hirsutum*) and their "intensive" types. *Indian Journal Of Agricultural Sciences* 55(12): 735-738.

Bhardwaj, S. N. and G. P. Verma (1985). Correlation between endogenous hormone level and fiber elongation in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Plant Physiology* 28(3): 249-258.

Bhattacharyya, D. K., M. M. Chakrabarty, et al. (1985). Studies on nutritional quality of interesterified fat products. *Fette Seifen Anstrichmittel* 87(12): 486-489.

Bilapate, G. G., A. K. Raodeo, et al. (1985). Investigations on *Heliothis armigera* in Marathwada (India): 10. Larval instars and progression factors on cotton (*Gossypium hirsutum*). *Journal Of Maharashtra Agricultural Universities* 10(1): 12-13.

Bindra, O. S. (1985). Relation of cotton cultivars to the cotton-pest problem in the Sudan Gezira. *Euphytica* 34(3): 849-856.

Boghara, D. G., N. P. Mehta, et al. (1985). Genetic studies on oil content in upland cotton (*Gossypium hirsutum*). *Gujarat Agricultural University Research Journal* 10(2): 1-4.

Bourland, F. M. and L. S. Bird (1985). A diallel analysis of seedling growth in cotton (*Gossypium hirsutum*). *Field Crops Research* 10(2): 197-204.

Brooks, D. W., H. S. Bevinakatti, et al. (1985). Practical total synthesis of racemic strigol. *Journal Of Organic Chemistry* 50(5): 628-632.

Brooks, D. W., H. S. Bevinakatti, et al. (1985). The absolute structure of dextro-strigol. *Journal Of Organic Chemistry* 50(20): 3779-3781.

Brown, S. M. and T. Whitwell (1985). Weed control programs for minimum-tillage cotton (*Gossypium hirsutum*). *Weed Science* 33(6): 843-847.

Brown, S. M., T. Whitwell, et al. (1985). Common bermudagrass (*Cynodon dactylon*) competition in cotton (*Gossypium hirsutum* cultivar Stoneville 213). *Weed Science* 33(4): 503-506.

Bucheli, P., M. Durr, et al. (1985). Beta-glucanases in developing cotton (*Gossypium hirsutum*) fibers. *Planta* 166(4): 530-536.

Burke, J. J., P. E. Gamble, et al. (1985). Plant morphological and biochemical responses to field water deficits: I. Responses of glutathione reductase activity and paraquat sensitivity. *Plant Physiology* 79(2): 415-419.

- Burke, J. J., J. L. Hatfield, et al. (1985). Accumulation of heat shock proteins in field-grown cotton (*Gossypium hirsutum*). *Plant Physiology* 78(2): 394-398.
- Butler, G. D. J. (1985). Populations of several insects on cotton (*Gossypium hirsutum* cultivar Deltapine 61) in open-top carbon dioxide enrichment chambers. *Southwestern Entomologist* 10(4): 264-267.
- Cathey, G. W. (1985). Conditioning cotton (*Gossypium hirsutum*) for increased response to defoliant chemicals. *Field Crops Research* 10(4): 347-354.
- Chakravarthy, A. K. and P. K. A. Rao (1985). Dispersion patterns, sample unit -sizes and techniques for sampling cotton jassid (*Amrasca biguttula biguttula*) and whitefly (*Bemisia tabaci*). *Insect Science And Its Application* 6(6): 661-666.
- Chiavegato, E. J., P. I. L. Gridi, et al. (1985). Interspecific hybridization between *Gossypium hirsutum* var. *latifolium* and *Gossypium herbaceum* var. *africanum*. *Bragantia* 44(2): 629-644.
- Chivinge, O. A. (1985). Allelopathic effects of purple nutsedge (*Cyperus rotundus*) on the growth and development of cotton (*Gossypium hirsutum*), maize (*Zea mays*) and soybeans (*Glycine max*). *Zimbabwe Agricultural Journal* 82(5): 151-152.
- Chopart, J. L. and D. Kone (1985). Influence of various tillage methods on the water supply of maize (*Zea mays*) and cotton (*Gossypium hirsutum*) crops in the Ivory Coast. *Agronomie Tropicale Riz Et Riziculture Et Cultures Vivrieres Tropicales* 40(3): 223-229.
- Coffey, B. N. and D. D. Davis (1985). Short-branch and cluster-fruiting habit inheritance in crosses of 8 cotton lines. *Crop Science* 25(5): 729-731.
- Coudriet, D. L., N. Prabhaker, et al. (1985). Sweet potato whitefly (*Bemisia tabaci*) (Homoptera: Aleyrodidae): Effects of neem-seed extract on oviposition and immature stages. *Environmental Entomology* 14(6): 776-779.
- Cramer, G. R., A. Lauchli, et al. (1985). Displacement of calcium by sodium from the plasmalemma of root cells: A primary response to salt stress? *Plant Physiology* 79(1): 207-211.
- Doman, D. C. and R. N. Trelease (1985). Protein A-gold immunocytochemistry of isocitrate lyase in cotton (*Gossypium hirsutum*) seeds. *Protoplasma* 124(3): 157-167.
- Elissalde, M. H., R. D. Stipanovic, et al. (1985). Histamine release from mast cells by terpenoid aldehydes isolated from glanded varieties of cotton. *American Industrial Hygiene Association Journal* 46(7): 396-401.
- Gerik, T. J., C. M. Rush, et al. (1985). Optimizing plot size for field studies of phymatotrichum root rot of cotton (*Gossypium hirsutum*). *Phytopathology* 75(2): 240-243.
- Glaubig, B. A. and F. T. Bingham (1985). Boron toxicity characteristics of 4 northern California (USA) endemic tree species. *Journal Of Environmental Quality* 14(1): 72-77.
- Guinn, G. (1985). Fruiting of cotton (*Gossypium hirsutum*): III. Nutritional stress and cutout. *Crop Science* 25(6): 981-985.
- Guinn, G. (1985). Abscisic acid and cutout in cotton (*Gossypium hirsutum*). *Plant Physiology* 77(1): 16-20.
- Gulbransen, B. (1985). Survival feeding of cattle with molasses: 2. Feeding steers with molasses/urea plus either sorghum grain (*Sorghum vulgare*) or cottonseed meal (*Gossypium hirsutum*). *Australian Journal Of Experimental Agriculture* 25(1): 4-8.
- Hallman, G. J. (1985). Comparison and contrast of cotton (*Gossypium hirsutum*) and *Desmodium tortuosum* as hosts of *Heliothis virescens*. *Insect Science And Its Application* 6(5): 573-578.

- Hearn, A. B. and R. G. D. Da (1985). A simple model for crop management applications for cotton (*Gossypium hirsutum*). *Field Crops Research* 12(1): 49-70.
- Heilman, M. D. (1985). Effect of mepiquat chloride and nitrogen levels on yield, growth characteristics and elemental composition of cotton (*Gossypium hirsutum*). *Journal Of Plant Growth Regulation* 4(1): 41-48.
- Hiltbold, A. E., R. M. Patterson, et al. (1985). Soil populations of *Rhizobium japonicum* in a cotton (*Gossypium hirsutum*), corn (*Zea mays*), soybean (*Glycine max*) rotation. *Soil Science Society Of America Journal* 49(2): 343-348.
- Hubac, C., E. Boutelier, et al. (1985). Drought resistance in cotton (*Gossypium hirsutum* cultivar Bou) plants: Some instances of intraindividual variations of metabolic parameters. *Bulletin De La Societe Botanique De France Actualites Botaniques* 132(2): 111-118.
- Huete, A. R., R. D. Jackson, et al. (1985). Spectral response of a plant canopy with different soil backgrounds. *Remote Sensing Of Environment* 17(1): 37-54.
- Imamaliyev, A. I. and I. E. Lavygina (1985). Dynamics of monosaccharides and cellulose in the fibers of intracultivar cotton hybrids. *Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni*: 11-13.
- Ismail, A. M. A. (1985). A comparative study of the effects of clumping, substrate, light, depth of burial and soil moisture upon the germination of seeds of weeds and crops grown in the Sudan Gezira. *Qatar University Science Bulletin* 5: 153-168.
- Johnson, L. F. and G. K. Palmer (1985). Symptom variability and selection for reduced severity of cotton (*Gossypium hirsutum*) seedling disease caused by *Pythium ultimum*. *Plant Disease* 69(4): 298-300.
- Jones, W. A. J., H. E. Walker, et al. (1985). Biology of *Niesthrea louisianica* (Hemiptera, Heteroptera: Rhopalidae) on selected plants, and its potential for biocontrol of velvetleaf, *Abutilon theophrasti* (Malvaceae). *Annals Of The Entomological Society Of America* 78(3): 326-330.
- Joshi, P. A., J. M. Stewart, et al. (1985). Localization of beta-glycerophosphatase activity in cotton (*Gossypium hirsutum*) fiber during differentiation. *Protoplasma* 125(1-2): 75-85.
- Kalsy, H. S., B. Singh, et al. (1985). Comparative incidence of pink bollworm (*Pectinophora gossypiella*) on some cultivars of *Gossypium hirsutum* cotton. *Crop Improvement* 12(2): 195-196.
- Kapse, S. S. and Y. S. Nerkar (1985). Polyacrylamide gel electrophoresis of soluble seed proteins in relation to cultivar identification in cotton. *Seed Science And Technology* 13(3): 847-852.
- Karve, A. D. and V. A. Deshmukh (1985). Photomorphogenic effect of high population density of reproductive development of crop plants. *Indian Botanical Reporter* 4(1): 1-4.
- Keim, K. R., J. E. Quisenberry, et al. (1985). Heritability of leaf characteristics in upland cotton (*Gossypium hirsutum*). *Crop Science* 25(2): 291-293.
- Kent, L. M. and A. Lauchli (1985). Germination and seedling growth of cotton (*Gossypium hirsutum*): Salinity-calcium interactions. *Plant Cell And Environment* 8(2): 155-160.
- Kerby, T. A. (1985). Cotton (*Gossypium hirsutum*) response to mepiquat chloride. *Agronomy Journal* 77(4): 515-518.
- Khan, M. A., S. M. Rao, et al. (1985). *Gossypium hirsutum* times *Gossypium barbadense* derivatives and their field performance under Faisalabad (Pakistan) conditions. *Pakistan Journal Of Agricultural Research* 6(3): 209-212.

- Khorgade, P. W. and A. P. Ekbote (1985). Biometrical relationships of certain characters in upland cotton (*Gossypium hirsutum*). P.K.: 1-4.
- Khushk, M. T. and J. G. Vaughan (1985). Seed structure in relation to the taxonomy of the Hibisceae (*Gossypium*, *Lebronnecia* and *Thespesia*). *Pakistan Journal Of Botany* 17(1): 119-130.
- Kishor, P. B. K. and P. Satyanarayana (1985). Effect of growth regulators on growth and phenolic production in cotton (*Gossypium hirsutum* cultivar Sankar-5) callus cultures. *Indian Botanical Reporter* 4(2): 111-114.
- Kohel, R. J. (1985). Genetic analysis of fiber color variants in cotton (*Gossypium hirsutum*). *Crop Science* 25(5): 792-797.
- Kohel, R. J., J. Glueck, et al. (1985). Comparison of cotton (*Gossypium hirsutum*) germplasm collections for seed-protein content. *Crop Science* 25(6): 961-963.
- Koskinen, W. C., H. R. Leffler, et al. (1985). Effect of trifluralin soil metabolites on cotton (*Gossypium hirsutum* cultivar Stoneville 213) boll components and fiber and seed properties. *Journal Of Agricultural And Food Chemistry* 33(5): 958-961.
- Krishnamurthi, M. S. (1985). Respiratory changes in bacterial blight of cotton (*Gossypium hirsutum* cultivar Lakshmi). *Indian Journal Of Plant Pathology* 3(2): 263-266.
- Lee, J. A. (1985). Inheritance of two new mutations in cotton. *Journal Of Heredity* 76(6): 479-480.
- Leffler, H. R. and J. H. Hunter (1985). Reproductive development and seed quality of cotton (*Gossypium hirsutum*) cultivars as affected by nitrogen fertilization. *Field Crops Research* 10(3): 219-228.
- Loper, G. M. and D. D. Davis (1985). Disparity of cotton pollen dispersal by honey bees (*Apis mellifera*) visiting upland (*Gossypium hirsutum*) and pima (*Gossypium barbadense*) pollen parents. *Crop Science* 25(4): 585-589.
- Lordello, R. R. A. and N. P. Sabino (1985). Effect of nematode control on fiber properties of upland cotton. *Bragantia* 44(2): 699-700.
- Mangat, N. S. (1985). Impact of weather and technology on cotton production in Punjab (India). *Indian Journal Of Agricultural Sciences* 55(5): 328-332.
- Marani, A., D. N. Baker, et al. (1985). Effect of water stress on canopy senescence and carbon exchange rates in cotton (*Gossypium hirsutum*). *Crop Science* 25(5): 798-802.
- Marani, A. and J. Ephrath (1985). Penetration of radiation into cotton (*Gossypium hirsutum* cultivar Acala SJ 2) crop canopies. *Crop Science* 25(2): 309-313.
- Mary, T. N. and K. V. K. Rao (1985). Cytomorphological studies in a haploid of *Gossypium hirsutum*. *Indian Journal Of Botany* 8(2): 115-121.
- McMichael, B. L., J. J. Burke, et al. (1985). Root vascular bundle arrangements among cotton (*Gossypium hirsutum*) strains and cultivars. *Environmental And Experimental Botany* 25(1): 23-30.
- Mellon, J. E. and L. S. Lee (1985). Elicitation of cotton isoperoxidases by *Aspergillus flavus* and other fungi pathogenic to cotton (*Gossypium hirsutum*). *Physiological Plant Pathology* 27(3): 281-288.
- Menges, R. M., P. R. Nixon, et al. (1985). Light reflectance and remote sensing of weeds in agronomic and horticultural crops. *Weed Science* 33(4): 569-581.
- Menzel, M. Y., K. L. Richmond, et al. (1985). A chromosome translocation breakpoint map of the *Gossypium hirsutum* genome. *Journal Of Heredity* 76(6): 406-414.

- Meredith, W. R. J. (1985). Lint yield genotype times environment interaction in upland cotton (*Gossypium hirsutum*) as influenced by leaf canopy isolines. *Crop Science* 25(3): 509-512.
- Meyer, W. S. and D. C. Reicosky (1985). Enclosing leaves for water potential measurement and its effect on interpreting soil-induced water stress. *Agricultural And Forest Meteorology* 35(1): 187-192.
- Meyer, W. S., D. C. Reicosky, et al. (1985). Errors in field measurement of leaf diffusive conductance associated with leaf temperature. *Agricultural And Forest Meteorology* 36(1): 55-64.
- Mollet, J. A. and V. Sevacherian (1985). Scirtothrips citri (Thysanoptera: Thripidae) development on cotton leaves and/or eggs of Tetranychus cinnabarinus (Acari: Tetranychidae). *Journal Of The Kansas Entomological Society* 58(3): 532-533.
- Nandal, A. S., J. P. Singh, et al. (1985). Efficacy and economics of spray schedules with synthetic pyrethroids against bollworms of upland cotton (*Gossypium hirsutum* cultivar H 777). *Indian Journal Of Agricultural Sciences* 55(3): 185-189.
- Netzer, D., Y. Tal, et al. (1985). Resistance of interspecific cotton hybrids (*Gossypium hirsutum* times *Gossypium barbadense* containing *Gossypium harknessii* cytoplasm) to Fusarium wilt. *Plant Disease* 69(4): 312-313.
- Nie, R. Z. and M. X. Li (1985). Studies of karyotypes in 3 wild and 4 cultivated species of *Gossypium*. *Acta Botanica Sinica* 27(2): 113-121.
- Osman, A. A., K. S. M. Abo, et al. (1985). Toxicity of some new pesticides to mites on cotton. *Indian Journal Of Agricultural Sciences* 55(8): 533-536.
- Osman, A. A., K. S. M. Abo, et al. (1985). Field efficacy of some granular pesticides against early-season pests of cotton (*Gossypium hirsutum*) in Egypt. *Indian Journal Of Agricultural Sciences* 55(8): 531-532.
- Patil, R. A. and P. R. Chopde (1985). Heterosis for yield and its traits in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 55(6): 402-406.
- Patil, V. N. and C. H. Andrews (1985). Development and release of hardseeded to dormancy in cotton (*Gossypium hirsutum*). *Seed Science And Technology* 13(3): 691-698.
- Percy, R. G. and L. S. Bird (1985). Rust resistance expression in cotyledons, petioles and stems of *Gossypium hirsutum*. *Journal Of Heredity* 76(3): 202-204.
- Pessaraki, M. and T. C. Tucker (1985). Uptake of nitrogen-15 by cotton (*Gossypium hirsutum*) under salt stress. *Soil Science Society Of America Journal* 49(1): 149-152.
- Pham, T. A. T., F. C. Borrel, et al. (1985). Effects of water stress on lipid metabolism in cotton (*Gossypium hirsutum*) leaves. *Phytochemistry* 24(4): 723-728.
- Price, C. E. and N. H. Anderson (1985). Uptake of chemicals from foliar deposits: Effects of plant species and molecular structure. *Pesticide Science* 16(4): 369-377.
- Quimby, P. C. J. and K. E. Frick (1985). Evaluation of herbicide-coated larvae of *Bactra verutana* (Lepidoptera: Tortricidae) to control nutsedges (*Cyperus rotundus* and *Cyperus esculentus*). *Entomophaga* 30(3): 287-292.
- Quisenberry, J. E., C. W. Wendt, et al. (1985). Potential for using leaf turgidity to select drought tolerance in cotton (*Gossypium hirsutum*). *Crop Science* 25(2): 294-299.
- Radin, J. W., J. R. Mauney, et al. (1985). Effects of nitrogen fertility on plant water relations and stomatal responses to water stress in irrigated cotton (*Gossypium hirsutum*). *Crop Science* 25(1): 110-115.

- Rakhimov, D. A., N. P. Yuldasheva, et al. (1985). Polysaccharides from the waste products of some fruit, berry, vegetable and industrial crops. *Khimiya Prirodnikh Soedinenii*(1): 21-24.
- Rao, V. S. and J. A. Inamdar (1985). Leaf architecture in cultivars of cotton. *Phyton* 25(1): 65-72.
- Reicosky, D. C., W. S. Meyer, et al. (1985). Cotton (*Gossypium hirsutum* cultivar Deltapine) response to short-term waterlogging imposed with a water-table gradient facility. *Agricultural Water Management* 10(2): 127-144.
- Reicosky, D. C., R. C. G. Smith, et al. (1985). Foliage temperature as a means of detecting stress of cotton subjected to a short-term water-table gradient. *Agricultural And Forest Meteorology* 35(1): 193-204.
- Rikin, A., E. Chalutz, et al. (1985). Rhythmicity in cotton (*Gossypium hirsutum*) seedlings: Rhythmic ethylene production as affected by silver ions as related to other rhythmic processes. *Planta* 163(2): 227-231.
- Rushing, D. W., D. S. Murray, et al. (1985). Weed interference with cotton (*Gossypium hirsutum*): I. Buffalo bur (*Solanum rostratum*). *Weed Science* 33(6): 810-814.
- Rushing, D. W., D. S. Murray, et al. (1985). Weed interference with cotton (*Gossypium hirsutum*): II. Tumble pigweed (*Amaranthus albus*). *Weed Science* 33(6): 815-818.
- Ryser, U. and P. J. Holloway (1985). Ultrastructure and chemistry of soluble and polymeric lipids in cell walls from seed coats and fibers of *Gossypium* species. *Planta* 163(2): 151-163.
- Sagar, P. and V. O. Reddy (1985). Host range and preference of mentha leaf webber, *Syngamia abruptalis* (Lepidoptera: Pyralidae) in Punjab (India). *Journal Of Research Punjab Agricultural University* 22(3): 488-492.
- Sanyasi, I. P. S., B. R. Mor, et al. (1985). Genetics of growth characters in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Research* 19(3): 143-149.
- Sasek, T. W., E. H. Delucia, et al (1985). Reversibility of photosynthetic inhibition in cotton after long-term exposure to elevated carbon dioxide concentrations. *Plant Physiology* 78(3): 619-622.
- Sharma, R. K. and Y. S. Murty (1985). The interrelationship of growth substances and stomatal characters in *Gossypium hirsutum* cultivar H-777. *Proceedings Of The Indian Academy Of Sciences Plant Sciences* 95(1): 21-28.
- Shin, I. V., L. V. Chandylova, et al. (1985). Effect of bicarbonate on the Hill reaction and photophosphorylation in chloroplasts of peas and cotton. *Fiziologiya Rastenii* 32(5): 837-842.
- Shivkumar, P. and H. G. Singh (1985). Genetic analysis of protein content in cotton seed. *Indian Journal Of Genetics & Plant Breeding* 45(2): 385-388.
- Sidhu, A. S. and A. K. Dhawan (1985). Effect of different spacings on incidence of cotton jassid (*Amrasca biguttula biguttula*) and pink bollworm (*Pectinophora gossypiella*) on F 414 variety of *Gossypium hirsutum* cotton. *Journal Of Research Punjab Agricultural University* 22(3): 483-487.
- Singh, M., T. H. Singh, et al. (1985). Genetic analysis of some seed quality characters in upland cotton (*Gossypium hirsutum*). *Theoretical And Applied Genetics* 71(1): 126-128.
- Singh, M., V. P. Singh, et al. (1985). Selection for yield and quality of *Gossypium hirsutum*. *Indian Journal Of Agricultural Sciences* 55(8): 521-525.
- Singh, V. V. and A. K. Singh (1985). Variability for seed yield, seed index and oil content in the germ-plasm of upland cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Sciences* 55(5): 321-323.

- Smith, C. W. and J. J. Varvil (1985). Fungicide and temperature affect leachate-predicted germination of cotton (*Gossypium hirsutum*). *Agronomy Journal* 77(1): 9-12.
- Stone, T. B., A. C. Thompson, et al. (1985). Analysis of lipids in cotton (*Gossypium hirsutum*) extrafloral nectar. *Journal Of Entomological Science* 20(4): 422-428.
- Street, J. E., C. E. Snipes, et al. (1985). Competition of a binary weed system with cotton (*Gossypium hirsutum*). *Weed Science* 33(6): 807-809.
- Suttle, J. C. (1985). Involvement of ethylene in the action of the cotton (*Gossypium hirsutum* cultivar Stoneville 519) defoliant thidiazuron. *Plant Physiology* 78(2): 272-276.
- Tabor, J. A., A. W. Warrick, et al. (1985). Spatial variability of nitrate in irrigated cotton (*Gossypium hirsutum*): 2. Soil nitrate and correlated variables. *Soil Science Society Of America Journal* 49(2): 390-394.
- Temple, P. J., O. C. Taylor, et al. (1985). Cotton yield responses to ozone as mediated by soil moisture and evapotranspiration. *Journal Of Environmental Quality* 14(1): 55-60.
- Thead, L. G., H. N. Pitre, et al. (1985). Feeding behavior of adult *Geocoris punctipes* (Hemiptera: Heteroptera: Lygaeidae) on nectaried (cultivar Stoneville-213) and nectariless (cultivar Stoneville 7A) cotton (*Gossypium hirsutum*). *Environmental Entomology* 14(2): 134-137.
- Trufem, S. F. B. and V. L. Bononi (1985). Vesicular-arbuscular mycorrhizae in crop plants from "cerrado" areas. *Rickia* 12: 165-187.
- Tzeng, D. D. and J. E. Devay (1985). Physiological responses of *Gossypium hirsutum* cultivar Acala SJ-2 to infection by defoliating and nondefoliating pathotypes of *Verticillium dahliae*. *Physiological Plant Pathology* 26(1): 57-72.
- Tzeng, D. D., R. J. Wakeman, et al. (1985). Relationships among verticillium wilt development, leaf water potential, phenology and lint yield in cotton (*Gossypium hirsutum*). *Physiological Plant Pathology* 26(1): 73-82.
- Vigil, E. L., R. L. Steere, et al. (1985). Structure of plasma membrane in radicles from cotton seeds (*Gossypium hirsutum* cultivar M-8). *Protoplasma* 129(2-3): 168-177.
- Villacorta, A., A. P. Gutierrez, et al. (1985). An analysis of growth and development of cotton (*Gossypium hirsutum* cultivar IAC-17) in Parana state (Brazil): A simulation model for the variety IAC-17. *Pesquisa Agropecuaria Brasileira* 20(1): 115-128.
- Virk, J. S., R. Singh, et al. (1985). Effect of planting pattern and spacing on the growth and yield of upland cotton (*Gossypium hirsutum*). *Journal Of Research Punjab Agricultural University* 22(2): 220-228.
- Voss, M. and N. Sidiras (1985). Nodulation of soybean (*Glycine max*) under no-tillage in comparison to conventional tillage. *Pesquisa Agropecuaria Brasileira* 20(7): 775-782.
- Waller, G. D., J. O. Moffet, et al. (1985). An evaluation of honey bee foraging activity and pollination efficacy for male-sterile cotton. *Crop Science* 25(2): 211-214.
- Waller, G. D., B. E. Vaissiere, et al. (1985). Comparison of carpenter bees (*Xylocopa varipuncta*) (Hymenoptera: Anthophoridae) and honeybees (*Apis mellifera*) (Hymenoptera: Apidae) as pollinators of male-sterile cotton in cages. *Journal Of Economic Entomology* 78(3): 558-561.
- Watson, J. E., I. L. Pepper, et al. (1985). Yields and leaf elemental composition of cotton (*Gossypium hirsutum*) grown on sludge-amended soil. *Journal Of Environmental Quality* 14(2): 174-177.
- Wilhelm, S., J. E. Sagen, et al. (1985). Phenotype modification in cotton for control of verticillium wilt through dense plant population culture. *Plant Disease* 69(4): 283-288.

- Willis, G. H., L. L. McDowell, et al. (1985). Toxaphene, methyl parathion and fenvalerate disappearance from cotton (*Gossypium hirsutum*) foliage in the Mid-South (USA). *Journal Of Environmental Quality* 14(3): 446-450.
- Wilson, F. D. (1985). A genetic study of number of involucre bract teeth in cotton (*Gossypium hirsutum*). *Crop Science* 25(1): 32-34.
- Wong, S. C., I. R. Cowan, et al. (1985). Leaf conductance in relation to rate of carbon dioxide assimilation: I. Influence of nitrogen nutrition, phosphorus nutrition, photon flux density, and ambient partial pressure of carbon dioxide during ontogeny. *Plant Physiology* 78(4): 821-825.
- Woodstock, L. W., K. Furman, et al. (1985). Relationship between weathering deterioration and germination, respiratory metabolism and mineral leaching from cottonseeds (*Gossypium hirsutum*). *Crop Science* 25(3): 459-466.
- Wu, Z., D. Liu, et al. (1985). Genetic analysis of quantitative characters in upland cotton (*Gossypium hirsutum*): I. Estimation of genic effects on 17 agronomic characters from generation means. *Acta Genetica Sinica* 12(5): 344-349.
- York, A. C. and M. R. Tucker (1985). Nitrification inhibitor evaluation on cotton (*Gossypium hirsutum*): 2. Inhibitor incorporation depth and nitrogen placement. *Agronomy Journal* 77(3): 407-411.
- York, A. C. and M. R. Tucker (1985). Nitrification inhibitor evaluation on cotton (*Gossypium hirsutum*): 1. Rate and time of nitrogen application. *Agronomy Journal* 77(3): 403-406.
- Aliev, K. A., Z. N. Abdurakmanova, et al. (1984). Biosynthesis of ribulose bisphosphate carboxylase (EC 4.1.1.39) and development of the photosynthetic apparatus activity in cotton (*Gossypium hirsutum*) leaves. *Fiziologiya I Biokhimiya Kul'Turnykh Rastenii* 16(4): 324-329.
- Azevedo, D. M. P. D. and N. E. D. M. Beltrao (1984). Weed control herbicide selectivity in perennial cotton (*Gossypium hirsutum* var. *marie galante*) associated with corn (*Zea mays*) and cowpea (*Vigna unguiculata*). *Pesquisa Agropecuaria Brasileira* 19(5): 583-590.
- Bailey, J. C. (1984). Influence of nectaried and nectariless cottons, *Gossypium hirsutum*, on field populations of spider mites, *Tetranychus urticae*. *Journal Of The Georgia Entomological Society* 19(4): 437-439.
- Barbosa, L. M. and P. R. C. Castro (1984). Some effects of plant growth regulators on the morphology of cotton *Gossypium hirsutum* cultivar IAC -17. *Hoehnea* 11: 59-66.
- Basu, A. K. and M. G. Bhat (1984). Inheritance of certain qualitative characters in cotton (*Gossypium hirsutum*). *Indian Journal Of Genetics & Plant Breeding* 44(1): 153-158.
- Beltrao, N. E. D. M., D. M. P. D. Azevedo, et al. (1984). The optimum planting date of sorghum (*Sorghum bicolor*) in association with perennial cotton (*Gossypium hirsutum* cultivar *marie-galante*). *Pesquisa Agropecuaria Brasileira* 19(6): 709-718.
- Bharambe, P. R. and S. B. Varade (1984). Effects of plant water stress on some biochemical changes in cotton. *Journal Of Maharashtra Agricultural Universities* 9(1): 47-50.
- Bhardwaj, H. J. and J. B. J. Weaver (1984). Combining-ability analysis in cotton (*Gossypium hirsutum*) for agronomic characters, fruiting efficiency, photosynthesis and bollworm resistance. *Journal Of Agricultural Science* 103(3): 511-518.
- Bhatt, J. G., C. V. Raman, et al. (1984). The effect of removing bracts on boll growth, oil and protein content of seeds and fiber characters of lint in cotton. *Indian Journal Of Plant Physiology* 27(1): 8-14.

- Bielorai, H., I. Vaisman, et al. (1984). Drip irrigation of cotton (*Gossypium hirsutum* cultivar Acala Sj-2) with treated municipal effluents: 1. Yield response. *Journal Of Environmental Quality* 13(2): 231-234.
- Bilapate, G. G., A. K. Raodeo, et al. (1984). Investigations on *Heliothis armigera* in Marathwada (India): 6. Life tables studies on cotton (*Gossypium hirsutum*) squares. *Journal Of Maharashtra Agricultural Universities* 9(3): 261-262.
- Brown, J. K. and M. R. Nekon (1984). Geminate particles associated with cotton leaf crumple disease in Arizona (USA). *Phytopathology* 74(8): 987-990.
- Carvalho, L. P. D., F. B. Cavalcanti, et al. (1984). Influence of ramulosis on fiber characteristics and yield of cotton (*Gossypium hirsutum* f. *latifolium*). *Fitopatologia Brasileira* 9(3): 593-598.
- Castrejon, S. A., D. M. L. Ortego, et al. (1984). Detection of some hydrolytic and oxidative enzymes in the tissue of cotton (*Gossypium hirsutum* cultivar Coker 310) parasitized by *Phymatotrichum omnivorum*. *Agrociencia*(56): 195-208.
- Charyulu, N. R. and P. A. Rao (1984). Combining ability analysis for seed cotton yield in interspecific cotton hybrids. *Indian Journal Of Genetics & Plant Breeding* 44(3): 469-471.
- Combrink, N. J. J. (1984). Physiological leaf abnormalities and boll yield of cotton (*Gossypium hirsutum*) as affected by potassium, calcium and boron nutrition. *South African Journal Of Plant And Soil* 1(4): 106-110.
- Cope, J. T., J. G. Starling, et al. (1984). Response of peanuts (*Arachis hypogaea*) and other crops to fertilizers and lime in 2 long-term experiments. *Peanut Science* 11(2): 91-94.
- Ekbote, A. P. and P. W. Khorgade (1984). Genetic divergence in upland cotton (*Gossypium hirsutum*). P.K.: 1-4.
- Feigin, A., H. Bielorai, et al. (1984). Drip irrigation of cotton (*Gossypium hirsutum*) with treated municipal effluents: 2. Nutrient availability in soil. *Journal Of Environmental Quality* 13(2): 234-238.
- Ferrari, I. R., A. T. P. Thi, et al. (1984). Effect of water stress on the lipid and fatty acid composition of cotton (*Gossypium hirsutum* cultivar Reba) chloroplasts. *Physiologia Plantarum* 62(2): 219-224.
- Gausman, H. W., J. E. Quisenberry, et al. (1984). Leaf spectral measurements to screen cotton (*Gossypium hirsutum*) strains for characters affected by stress. *Field Crops Research* 9(5): 373-382.
- Gerard, C. J., P. D. Sexton, et al. (1984). Effect of furrow diking, subsoiling and slope position on crop yields. *Agronomy Journal* 76(6): 945-950.
- Goeschl, J. D., C. E. Magnuson, et al. (1984). Spontaneous and induced blocking and unblocking of phloem transport. *Plant Cell And Environment* 7(8): 607-614.
- Gross, H. R. J., W. J. Lewis, et al. (1984). *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae): Effects of augmented densities and distributions of *Heliothis zea* (Lepidoptera: Noctuidae) host eggs and kairomones on field performance. *Environmental Entomology* 13(4): 981-985.
- Grossmann, K., W. Rademacher, et al. (1984). Comparative potency of different plant growth retardants in cell cultures and intact plants. *Journal Of Plant Growth Regulation* 3(4): 197-206.
- Gupta, G. P., N. P. Agnihotri, et al. (1984). Efficacy and residues of synthetic pyrethroids in cotton (*Gossypium hirsutum* cultivar Bikaneri Nerma). *Journal Of Entomological Research* 8(1): 73-77.
- Gupta, S. N., D. S. Jatasra, et al. (1984). Heterosis for nitrogen, phosphorus and potassium in leaves and their relationship with heterosis for yield in upland cotton (*Gossypium hirsutum*). *Indian Journal Of Genetics & Plant Breeding* 44(1): 134-139.

- Gupta, S. P. and T. H. Singh (1984). Genetic analysis of some fiber and seed characters in upland cotton (*Gossypium hirsutum*). *Crop Improvement* 11(2): 132-134.
- Gutierrez, A. P., M. A. Pizzamiglio, et al. (1984). A general distributed delay time varying life table plant population model: Cotton (*Gossypium hirsutum*) growth and development as an example. *Ecological Modelling* 26(3-4): 231-250.
- Hendrix, D. L. and J. W. Radin (1984). Seed development in cotton (*Gossypium hirsutum*): Feasibility of a hormonal role for abscisic acid in controlling vivipary. *Journal Of Plant Physiology* 117(3): 211-222.
- Herrera, R. L. A., L. F. Valencia, et al. (1984). Relationships between some physical parameters and humidity constants in determining water levels for irrigation. *Acta Agronomica* 34(2): 26-41.
- Hill, R. L., B. Dalili, et al. (1984). Effect of seed-treatment methods and soil water content on early growth and acephate uptake by corn (*Zea mays*) and cotton (*Gossypium hirsutum*) plants. *Journal Of Economic Entomology* 77(4): 880-884.
- Hillocks, R. J. (1984). Production of cotton varieties with resistance to *Fusarium* wilt with special reference to Tanzania. *Tropical Pest Management* 30(3): 234-246.
- Hoo, C. R. S., D. L. Coudriet, et al. (1984). *Trichoplusia ni* (Lepidoptera: Noctuidae) larval development on wild and cultivated plants. *Environmental Entomology* 13(3): 843-846.
- Howell, T. A., K. R. Davis, et al. (1984). Water use efficiency of narrow row cotton (*Gossypium hirsutum* cultivar Acala). *Irrigation Science* 5(3): 195-214.
- Howell, T. A., J. L. Hatfield, et al. (1984). Response of cotton (*Gossypium hirsutum* cultivar Acala-ST-Z) water stress indicators to soil salinity. *Irrigation Science* 5(1): 25-36.
- Hubac, C., M. Ouedraogo, et al. (1984). Effect of photoperiod on the adaptation to drought of cotton (*Gossypium hirsutum*) plants. *Bulletin De La Societe Botanique De France Actualites Botaniques* 131(1): 79-88.
- Hussain, T. (1984). Prevalence and distribution of *Xanthomonas campestris* pathovar *malvacearum* races in Pakistan and their reaction to different cotton (*Gossypium hirsutum*) lines. *Tropical Pest Management* 30(2): 159-162.
- Idso, S. B., P. J. J. Pinter, et al. (1984). Stomatal conductance and photosynthesis in water hyacinth (*Eichhornia crassipes*): Effects of removing water from roots as quantified by a foliage-temperature-based plant water stress index. *Agricultural And Forest Meteorology* 32(3-4): 249-256.
- Kalsy, H. S. and H. R. Garg (1984). Stability of yield in upland cotton (*Gossypium hirsutum*) in Punjab (India). *Indian Journal Of Agricultural Sciences* 54(11): 967-969.
- Kaushik, L. S., D. P. Singh, et al. (1984). Line times tester analysis for fixed effect model in cotton (*Gossypium hirsutum*). *Theoretical And Applied Genetics* 68(6): 487-491.
- Kerbabaeva, Z. A., A. S. Konoplya, et al. (1984). Evaluation of the homeostaticity of cotton (*Gossypium hirsutum*) based on yield response to increased stand density. *Izvestiya Akademii Nauk Turkmenskoi Ssr Seriya Biologicheskikh Nauk*(5): 51-55.
- Kiristaeva, N. M., M. V. Satskaya, et al. (1984). The role of hypocotyls in the process of ascending distant transport of amino acids and amides in cotton (*Gossypium hirsutum*). *Izvestiya Akademii Nauk Sssr Seriya Biologicheskaya*(6): 933-938.
- Kiristaeva, N. M., M. V. Satskaya, et al. (1984). The role of hypocotyls in the process of the descent of amino acids and amides in cotton (*Gossypium hirsutum*). *Izvestiya Akademii Nauk Sssr Seriya Biologicheskaya*(5): 771-777.

- Kolte, T. B. and M. V. Thombre (1984). Heterosis and combining ability studies in *Gossypium* spp. *Journal Of Maharashtra Agricultural Universities* 9(3): 252-254.
- Kunce, C. M., R. N. Trelease, et al. (1984). Ontogeny of glyoxysomes in maturing and germinated cottonseeds (*Gossypium hirsutum*): A morphometric analysis. *Planta* 161(2): 156-164.
- Lee, J. A. (1984). 2 new alleles at the *sm-1* locus in cotton (*Gossypium hirsutum*). *Crop Science* 24(5): 945-947.
- Lee, J. A. (1984). Effects of plant smoothness on agronomic traits of upland cotton (*Gossypium hirsutum*): Fiber properties. *Crop Science* 24(4): 716-720.
- Lima, E. F., R. M. Vieira, et al. (1984). Influence of *Rhizopus* sp., *Aspergillus niger* and *Aspergillus flavus* on cotton, (*Gossypium hirsutum*) seed deterioration under storage. *Fitopatologia Brasileira* 9(3): 555-560.
- Mangat, N. S. (1984). Optimum size and shape of plot and block in cotton (*Gossypium hirsutum*) field experiments. *Journal Of Research Punjab Agricultural University* 21(4): 581-589.
- Manohar, N., T. Guruswamy, et al. (1984). Estimation of leaf area of a few local varieties of different species of cotton. *Mysore Journal Of Agricultural Sciences* 18(1): 39-42.
- Matveev, G. G. (1984). Fusarium wilt of upland cotton (*Gossypium hirsutum*). *Mikologiya I Fitopatologiya* 18(5): 412-416.
- Matveev, G. G. and K. Zhuraev (1984). Resistance factors of various cotton cultivars to *Verticillium* wilt. *Mikologiya I Fitopatologiya* 18(3): 239-245.
- McMichael, B. L., W. R. Jordan, et al. (1984). Leaf production and growth rates of exotic cottons (*Gossypium hirsutum*). *Agronomy Journal* 76(6): 901-905.
- Menzel, M. Y., S. Naqi, et al. (1984). Incipient genome differentiation in *Gossypium*: 4. The genome of *Gossypium laxum*. *Journal Of Heredity* 75(5): 389-391.
- Meredith, W. R. J. (1984). Influence of leaf morphology on lint yield of cotton (*Gossypium hirsutum*): Enhancement by the sub okra trait. *Crop Science* 24(5): 855-857.
- Morison, J. I. L. and R. M. Gifford (1984). Plant growth and water use with limited water supply in high carbon dioxide concentrations: 1. Leaf area, water use and transpiration. *Australian Journal Of Plant Physiology* 11(5): 361-374.
- Morison, J. I. L. and R. M. Gifford (1984). Plant growth and water use with limited water supply in high carbon dioxide concentrations: 2. Plant dry weight, partitioning and water use efficiency. *Australian Journal Of Plant Physiology* 11(5): 375-384.
- Murty, M. G. (1984). Phyllosphere of cotton as a habitat for diazotrophic microorganisms. *Applied And Environmental Microbiology* 48(4): 713-718.
- Mutsaers, H. J. W. (1984). KUTUN: A morphogenetic model for cotton (*Gossypium hirsutum*). *Agricultural Systems* 14(4): 229-258.
- Narsale, R. N., B. B. More, et al. (1984). Effects of *Azotobacter* with a fungicide and insecticides on cotton (*Gossypium hirsutum*). *Journal Of Maharashtra Agricultural Universities* 9(1): 79-80.
- Ogunlela, V. B., V. Kumar, et al. (1984). Effect of sowing date on the performance of 3 cotton varieties in the southern cotton growing zone of Nigeria. *Samaru Journal Of Agricultural Research* 2(1-2): 3-12.

- Omel'Chenko, V. S. (1984). Spontaneous hybridization and population structure of cotton (*Gossypium hirsutum*) cultivars with respect to heteromorphism of flowering. Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni: 15-16.
- Pavlovskaya, N. E. (1984). Distribution of multiple oxide reductase forms in various organs of cotton seedlings. Doklady Vsesoyuznoi Ordena Lenina I Ordena Trudovogo Krasnogo Znameni Akademii Sel'Skokhozyaistvennykh Nauk Imeni: 19-21.
- Pham, T. A. T. (1984). Effects of water stress on the polar lipids of cotton (*Gossypium hirsutum*) leaves. Bulletin De La Societe Botanique De France Actualites Botaniques 131(1): 89-98.
- Radin, J. W. (1984). Stomatal responses to water stress and to abscisic acid in phosphorus-deficient cotton (*Gossypium hirsutum*) plants. Plant Physiology 76(2): 392-394.
- Ramalho, F. S., J. C. J. McCarty, et al. (1984). Distribution of tobacco budworm (*Heliothis virescens*) (Lepidoptera: Noctuidae) larvae within cotton plants (*Gossypium hirsutum*). Journal Of Economic Entomology 77(3): 591-594.
- Ray, D. T. (1984). Metaphase I configurations of the reciprocal translocations in a cytogenetic tester set of *Gossypium hirsutum*. Journal Of Heredity 75(5): 371-377.
- Refaat, M., L. Rossignol, et al. (1984). Interspecific hybrid *Gossypium hirsutum* & *X Gossypium barbadense* via in vitro fertilization and ovule culture. Zeitschrift Fuer Pflanzenzuechtung 93(2): 137-146.
- Rikin, A., J. J. B. St, et al. (1984). Rhythmical changes in the sensitivity of cotton (*Gossypium hirsutum*) seedlings to herbicides. Plant Physiology 76(2): 297-300.
- Sadykhova, L. D. and T. K. Makhmudov (1984). Some cytoanatomical characters of the cotton hybrid, *Gossypium hirsutum* times *Gossypium stocksii*. Sel'Skokhozyaistvennaya Biologiya(4): 52-55.
- Saleh, H. and R. A. Sikora (1984). Relationship between *Glomus fasciculatum* root colonization of cotton (*Gossypium hirsutum* cultivar Coker 201) and its effect on *Meloidogyne incognita*. Nematologica 30(2): 230-237.
- Santana, J. C. F. D., S. E. O. Dos, et al. (1984). Evaluation of cotton (*Gossypium hirsutum* var. *latifolium*) genotypes in the northeast of Brazil. Pesquisa Agropecuaria Brasileira 19(6): 679-688.
- Sharma, R. K., Vidhu, et al. (1984). Influence of different growth substances on seed germination and seedling growth of *Gossypium hirsutum* cultivar H-777. Indian Botanical Reporter 3(1): 60-73.
- Singh, D., J. S. Virk, et al. (1984). Response of promising strains and varieties of *Gossypium arboreum* and *Gossypium hirsutum* cotton to nitrogen fertilization. Journal Of Research Punjab Agricultural University 21(4): 501-508.
- Singh, M., T. H. Singh, et al. (1984). Combining ability and heterosis for some seed quality traits in upland cotton (*Gossypium hirsutum*). Crop Improvement 11(1): 30-34.
- Singh, M., V. P. Singh, et al. (1984). Breeding for desirable leaf types in cotton (*Gossypium hirsutum*). Indian Journal Of Plant Physiology 27(3): 247-254.
- Singh, N., P. S. Randhawa, et al. (1984). Quantitative and qualitative bacterial flora of 2 cotton cultivars in relation to age of the crop and environmental conditions. Indian Journal Of Ecology 11(1): 127-133.
- Singh, T. H. and S. S. Gill (1984). Genetic diversity in upland cotton (*Gossypium hirsutum*) under different environments. Indian Journal Of Genetics & Plant Breeding 44(3): 506-513.
- Stoilova, A. (1984). Irradiation effect on F-2 segregation for earliness, productiveness and fiber length in *Gossypium hirsutum* times *Gossypium barbadense* hybridization. Genetika I Seleksiya 17(3): 170-178.

- Sundararajan, K. S. (1984). Phase relations between the circadian leaf movement rhythm and its Zeitgeber within the range of entrainment in the cotton plant, *Gossypium hirsutum*. *Journal Of Interdisciplinary Cycle Research* 15(2): 155-162.
- Thomas, J. C. and F. R. H. Katterman (1984). The control of spontaneous lysis of protoplasts from *Gossypium hirsutum* anther callus. *Plant Science Letters* 36(2): 149-154.
- Touchton, J. T., D. H. Rickerl, et al. (1984). Winter legumes as a nitrogen source for no-tillage cotton (*Gossypium hirsutum*). *Soil & Tillage Research* 4(4): 391-402.
- Turner, N. C., E. D. Schulze, et al. (1984). The responses of stomata and leaf gas exchange to vapor pressure deficits and soil water content: 1. Species comparisons at high soil water contents. *Oecologia* 63(3): 338-342.
- Vieira, D. S. J. (1984). Application of fundamental research to the breeding of drought resistant plants. *Bulletin De La Societe Botanique De France Actualites Botaniques* 131(1): 51-58.
- Virk, J. S., S. R. Sharma, et al. (1984). Effect of Cycocel on yield and other characters of cotton (*Gossypium hirsutum*) cultivar F-414. *Journal Of Research Punjab Agricultural University* 21(4): 483-488.
- Virk, P. S., H. S. Kalsy, et al. (1984). Comparative estimation of genetic components of continuous variation in upland cotton (*Gossypium hirsutum*) using 2 approaches of diallel analysis. *Crop Improvement* 11(2): 111-114.
- Weaver, J. B. J., M. A. M. El, et al. (1984). Yield, fiber and spinning performance of interspecific cotton hybrids having a common parent. *Crop Science* 24(4): 637-640.
- Wells, R. and W. R. J. Meredith (1984). Comparative growth of obsolete and modern cotton (*Gossypium hirsutum*) cultivars: 3. Relationship of yield to observed growth characteristics. *Crop Science* 24(5): 868-872.
- Wells, R. and W. R. J. Meredith (1984). Comparative growth of obsolete and modern cotton (*Gossypium hirsutum*) cultivars: 1. Vegetative dry matter partitioning. *Crop Science* 24(5): 858-862.
- Wells, R. and W. R. J. Meredith (1984). Comparative growth of obsolete and modern cotton (*Gossypium hirsutum*) cultivars: 2. Reproductive dry matter partitioning. *Crop Science* 24(5): 863-868.
- Wilson, F. D. and B. W. George (1984). Pink bollworm (*Pectinophora gossypiella*) (Lepidoptera: Gelechiidae): Selecting for antibiosis in artificially and naturally infested cotton plants (*Gossypium hirsutum*). *Journal Of Economic Entomology* 77(3): 720-724.
- Wilson, F. D., B. W. George, et al. (1984). Pink bollworm (*Pectinophora gossypiella*) (Lepidoptera: Gelechiidae): Oviposition and larval success on resistant and susceptible cotton plants (*Gossypium hirsutum*). *Journal Of Economic Entomology* 77(3): 709-714.
- Yadava, B. D., S. S. Pahuja, et al. (1984). Cultural and chemical control of weeds in cotton (*Gossypium hirsutum*). *Indian Journal Of Agricultural Research* 18(4): 215-220.
- Yazkulyev, A. and A. Durdyev (1984). Cell thermoresistance and intracellular osmotic pressure in diploid and tetraploid species of the genus *Gossypium* (Malvaceae). *Izvestiya Akademii Nauk Turkmenskoi Ssr Seriya Biologicheskikh Nauk*(6): 3-8.
- Yunus Khanov, S. H. and A. P. Ibragimov (1984). Identification of protein markers of the seeds of *Gossypium hirsutum*, *Gossypium barbadense* and study of the relationship of their inheritance with some characters. *Genetika* 20(6): 989-997.

Zakharova, O. Y. and A. K. Ergashev (1984). Cytological study of the mutagenic action of the defoliant butylcaptopax in the cells of cotton of the species *Gossypium hirsutum* and *Gossypium herbaceum*. *Tsitologiya I Genetika* 18(4): 279-283.

Zeringue, H. J. J. (1984). The accumulation of 5 fluorescent compounds in the cotton leaf induced by cell-free extracts of *Aspergillus flavus*. *Phytochemistry* 23(11): 2501-2504.

Zummo, G. R., J. H. Benedict, et al. (1984). Effect of the plant growth regulator mepiquat chloride on host plant resistance in cotton (*Gossypium hirsutum*) to bollworm (*Heliothis zea*) (Lepidoptera: Noctuidae). *Journal Of Economic Entomology* 77(4): 922-924.

Barbosa, L. M. and P. R. C. Costo (1983). Comparison between concentrations and application time of mepiquat chloride, chlorocholine chloride and ethephon in cotton (*Gossypium hirsutum* cultivar IAC-17). *Planta Daninha* 6(1): 1-10.

Beltrao, N. E. D. M., J. F. D. Silva, et al. (1983). Resistance of cotton species and cultivars to the herbicide diuron. *Planta Daninha* 6(1): 72-78.

Beltrao, N. E. D. M., J. F. D. Silva, et al. (1983). The behavior of upland-type cotton (*Gossypium hirsutum* latifolium cultivar IAC-17) and the control of weeds after the use of diuron and sethoxydim herbicides. *Planta Daninha* 6(1): 58-71.

Cabangbang, R. P., E. C. Rinen, et al. (1983). Some observations on the morphological variation of cotton (*Gossypium hirsutum* cultivar Deltapine 16). *Philippine Journal Of Crop Science* 8(1): 27-32.

Morozova, A. V., S. I. Krasil'Nikova, et al. (1983). Fiber lengthening in cotton *Gossypium hirsutum* and *Gossypium barbadense*. *Izvestiya Akademii Nauk Turkmenskoi Ssr Seriya Biologicheskikh Nauk*(4): 11-14.

Wilson, L. T., D. R. Booth, et al. (1983). The behavioral activity and vertical distribution of the cotton harlequin bug *Tectocoris diophthalmus* (Heteroptera: Scutelleridae) on cotton plants in a glasshouse. *Journal Of The Australian Entomological Society* 22(4): 311-318.

Ismail, F. M. and A. M. Esmail (1982). Response of upland cotton variety McNair 220 (*Gossypium hirsutum* L.) to nitrogen and phosphorus fertilizations under the conditions of Middle Egypt. *Egyptian Journal Of Agronomy* 7(2): 101-110.

Riffle, M. S., G. R. Waller, et al. (1927). Devil's-claw (*Proboscidea louisianica*), essential oil and its components: Potential allelochemical agents on cotton and wheat. *Journal Of Chemical Ecology* 16(6): 1927-1940.

Mozafar, A. and J. J. Oertli (1893). Effect of temperature on ion efflux properties of leaf cell compartments in several chilling-sensitive and chilling-insensitive crop plants. *Canadian Journal Of Botany* 65(9): 1893-1900.

Lodhi, M. A. K., R. Bilal, et al. (1881). Allelopathy in agroecosystems: Wheat phytotoxicity and its possible roles in crop rotation. *Journal Of Chemical Ecology* 13(8): 1881-1892.

Archer, T. L., E. D. J. Bynum, et al. (1879). Chemical coverage on soybean and cotton plants sprayed with lithium sulfate using an airplane, chemigation, or a multifunction irrigation-pesticide application system. *Journal Of Economic Entomology* 84(6): 1879-1888.

Oosterhuis, D. M. and S. D. Wullschleger (1866). Water flow through cotton roots in relation to xylem anatomy. *Journal Of Experimental Botany* 38(196): 1866-1874.

Bland, W. L. and W. A. Dugas (1850). Cotton root growth and soil water extraction. *Soil Science Society Of America Journal* 53(6): 1850-1855.

- Mitchell, E. R., F. C. Tingle, et al. (1817). Ovipositional response of three *Heliothis* spp. (Lepidoptera: Noctuidae) to allelochemicals from cultivated and wild host plants. *Journal Of Chemical Ecology* 16(6): 1817-1828.
- Simongulyan, N. G. and V. L. Kim (1815). Variability of quantitative characters in the irradiated cotton population. *Genetika* 26(10): 1815-1821.
- Hassidim, M., Y. Braun, et al. (1795). Sodium ion-proton and potassium ion-proton antiport in root membrane vesicles isolated from the halophyte *Atriplex* and the glycophyte cotton. *Plant Physiology* 94(4): 1795-1801.
- Terry, L. I. and B. B. Barstow (1785). Susceptibility of early season cotton floral bud types to thrips (Thysanoptera: Thripidae) damage. *Journal Of Economic Entomology* 81(6): 1785-1791.
- Goldberg, N. P., M. C. Hawes, et al. (1760). Specific attraction to and infection of cotton root cap cells by zoospores of *Pythium dissotocum*. *Canadian Journal Of Botany* 67(6): 1760-1767.
- Dickens, J. C. (1759). Olfaction in the boll weevil, *Anthonomus grandis* (Coleoptera: Curculionidae): Electroantennogram studies. *Journal Of Chemical Ecology* 10(12): 1759-1786.
- Boylston, E. K. (1739). Presence of silicon in developing cotton fibers. *Journal Of Plant Nutrition* 11(12): 1739-1748.
- Chandler, L. D. and J. E. Wright (1699). Longevity of the boll weevil (Coleoptera: Curculionidae) on cotton and alternate feeding sources under various temperature regimes in the laboratory. *Journal Of Economic Entomology* 84(6): 1699-1704.
- Li, J., M. Zhou, et al. (1689). Cotton response to zinc fertilizer. *Communications In Soil Science And Plant Analysis* 22(15-16): 1689-1700.
- Wilson, F. D., B. W. George, et al. (1671). Pink bollworm (Lepidoptera: Gelechiidae) egg hatch, larval success, and pupal and adult survival on okra- and normal-leaf cotton. *Journal Of Economic Entomology* 79(6): 1671-1675.
- Mauk, P. A. and R. B. Hine (1662). Infection, colonization of *Gossypium hirsutum* and *Gossypium barbadense*, and development of black root rot caused by *Thielaviopsis basicola*. *Phytopathology* 78(12 Part 2): 1662-1667.
- Koch, D. O., M. J. Jeger, et al. (1657). Effects of plant density on progress of *Phymatotrichum* root rot in cotton. *Phytopathology* 77(12): 1657-1662.
- Jeger, M. J., C. M. Kenerley, et al. (1647). Spatial dynamics of *Phymatotrichum* root rot in row crops in the Blackland Region of north central Texas (USA). *Phytopathology* 77(12): 1647-1656.
- Goldstein, C. S. and J. M. Widholm (1641). Photosynthetic characterization of photoautotrophic cells cultured in a minimal medium. *Plant Physiology* 94(4): 1641-1646.
- Derrick, M. E. and J. A. Durant (1627). Rate response of cotton to three pyrethroids. *Journal Of Economic Entomology* 79(6): 1627-1631.
- Jones, R. W., J. R. Cate, et al. (1626). Phenology and ecology of wild cotton (Malvales: Malvaceae) and the boll weevil (Coleoptera: Curculionidae) in Tamaulipas, Mexico. *Journal Of Economic Entomology* 82(6): 1626-1632.
- Morgan, P. W., C. J. He, et al. (1616). Does water deficit stress promote ethylene synthesis by intact plants? *Plant Physiology* 94(4): 1616-1624.

- Trichilo, P. J., L. T. Wilson, et al. (1604). Relative abundance of three species of spider mites (Acari: Tetranychidae) on cotton, as influenced by pesticides and time of establishment. *Journal Of Economic Entomology* 83(4): 1604-1611.
- Davis, D. A., D. Tsao, et al. (1603). Enhancement of phytoalexin accumulation in cultured plant cells by oxalate. *Phytochemistry* 31(5): 1603-1607.
- Bland, W. L. (1595). Estimating root length density by the core-break method. *Soil Science Society Of America Journal* 53(5): 1595-1597.
- Bradow, J. M. (1595). Cotton cultivar responses to suboptimal postemergent temperatures. *Crop Science* 31(6): 1595-1599.
- Bradow, J. M. (1585). Chilling sensitivity of photosynthetic oil-seedlings: I. Cotton and sunflower. *Journal Of Experimental Botany* 41(233): 1585-1594.
- Durant, J. A. (1577). Effect of treatment regimen on control of *Heliothis virescens* and *Helicoverpa zea* (Lepidoptera: Noctuidae) on cotton. *Journal Of Economic Entomology* 84(5): 1577-1584.
- McGarry, D. and I. G. Daniells (1575). Shrinkage curve indices to quantify cultivation effects on soil structure of a vertisol. *Soil Science Society Of America Journal* 51(6): 1575-1580.
- Nadzhimov, U. K., A. S. Almatov, et al. (1556). Genetic and physiological characteristics of a new tall mutant in cotton *Gossypium hirsutum* L. *Genetika* 27(9): 1556-1560.
- Russo, D. and D. Bakker (1554). Crop-water production functions for sweet corn and cotton irrigated with saline waters. *Soil Science Society Of America Journal* 51(6): 1554-1562.
- Youngman, R. R., T. F. Leigh, et al. (1549). Pesticides and cotton: Effect on photosynthesis, growth, and fruiting. *Journal Of Economic Entomology* 83(4): 1549-1557.
- Chapman, K. D. and R. N. Trelease (1525). Inhibition of cottonseed choline phosphotransferase and ethanolamine phosphotransferase by calcium during postgerminative growth. *Plant Physiology* 93(4): 1525-1529.
- Bhatti, A. U., D. J. Mulla, et al. (1523). Identifying and removing spatial correlation from yield experiments. *Soil Science Society Of America Journal* 55(6): 1523-1528.
- Saxena, K. N. and H. Kumar (1521). Interference of sonic communication and mating in leafhopper *Amrasca devastans* by certain volatiles. *Journal Of Chemical Ecology* 10(10): 1521-1532.
- Wilson, F. D., H. M. Flint, et al. (1516). Resistance of cotton lines containing a *Bacillus thuringiensis* toxin to pink bollworm (Lepidoptera: Gelechiidae) and other insects. *Journal Of Economic Entomology* 85(4): 1516-1521.
- Gulin, V. V., Y. E. Dubrova, et al. (1515). Genetic differentiation of some cotton cultivars (*Gossypium hirsutum* L.). *Genetika* 26(8): 1515-1518.
- Roeske, C. A., J. M. Widholm, et al. (1512). Photosynthetic carbon metabolism in photoautotrophic cell suspension cultures grown at low and high carbon dioxide. *Plant Physiology* 91(4): 1512-1519.
- Wiegand, C. L., J. H. Everitt, et al. (1511). Comparison of multispectral video and SPOT-1 HRV observations for cotton affected by soil salinity. *International Journal Of Remote Sensing* 13(8): 1511-1525.
- Butler, G. D. J., F. D. Wilson, et al. (1500). Cotton leaf crumple virus disease in okra -leaf and normal-leaf cottons. *Journal Of Economic Entomology* 78(6): 1500-1502.

- Bailey, J. C. and G. W. Cathey (1485). Effect of chlordimeform on tarnished plant bug (*Lygus lineolaris*) (Heteroptera: Miridae) nymph emergence. *Journal Of Economic Entomology* 78(6): 1485-1487.
- Baker, R. S., M. L. Laster, et al. (1481). Effects of the herbicide monosodium methanearsonate on insect and spider populations in cotton fields. *Journal Of Economic Entomology* 78(6): 1481-1484.
- Silva, E. C. D., V. Q. Ribeiro, et al. (1455). An inverse quadratic model to determine the size and shape of plots for maize (*Zea mays*) and cotton (*Gossypium hirsutum*) intercropped. *Pesquisa Agropecuaria Brasileira* 20(12): 1455-1460.
- Altman, D. W., R. D. Stipanovic, et al. (1451). Terpenoid aldehydes in upland cottons: II. Genotype-environment interactions. *Crop Science* 29(6): 1451-1456.
- Cakmak, I. and H. Marschner (1449). Enhanced superoxide radical production in roots of zinc-deficient plants. *Journal Of Experimental Botany* 39(207): 1449-1460.
- Goldschmidt, E. E. and S. G. Huber (1443). Regulation of photosynthesis by end-product accumulation in leaves of plants storing starch, sucrose, and hexose sugars. *Plant Physiology* 99(4): 1443-1448.
- Loreto, F., P. C. Harley, et al. (1437). Estimation of mesophyll conductance to carbon dioxide flux by three different methods. *Plant Physiology* 98(4): 1437-1443.
- Wood, C. W., P. W. Tracy, et al. (1435). Determination of cotton nitrogen status with a hand-held chlorophyll meter. *Journal Of Plant Nutrition* 15(9): 1435-1448.
- Flint, H. M., J. R. Merkle, et al. (1431). Pink bollworm (*Pectinophora gossypiella*) (Lepidoptera: Gelechiidae): Field testing a new polyethylene tube dispenser for gossypure. *Journal Of Economic Entomology* 78(6): 1431-1436.
- Gulin, V. V., K. Y. Saidov, et al. (1431). Genetic control of benzidinase in cotton *Gossypium hirsutum* L. *Genetika* 25(8): 1431-1435.
- Fialho, E. T., L. F. T. Albino, et al. (1419). Chemical composition and energetic values of some feedstuffs for swine. *Pesquisa Agropecuaria Brasileira* 20(12): 1419-1432.
- Brugnoli, E., K. T. Hubick, et al. (1418). Correlation between the carbon isotope discrimination in leaf starch and sugars of C-3 plants and the ratio of intercellular and atmospheric partial pressures of carbon dioxide. *Plant Physiology* 88(4): 1418-1424.
- Rosolem, C. A. and D. S. Mikkelsen (1417). Nitrogen source-sink relationship in cotton. *Journal Of Plant Nutrition* 12(12): 1417-1434.
- Bailey, J. C. (1410). Infesting cotton with tarnished plant bug (*Lygus lineolaris*) (Heteroptera: Miridae) nymphs reared by improved laboratory rearing methods. *Journal Of Economic Entomology* 79(5): 1410-1412.
- Terry, L. I. (1402). Effect of early season insecticide use and square removal on fruiting patterns and fiber quality of cotton. *Journal Of Economic Entomology* 85(4): 1402-1412.
- Kasperbauer, M. J. and W. J. Busscher (1376). Genotypic differences in cotton root penetration of a compacted subsoil layer. *Crop Science* 31(5): 1376-1378.
- Seagull, R. W. (1373). Changes in microtubule organization and wall microfibril orientation during in vitro cotton (*Gossypium hirsutum* cultivar Acala SJ-2) fiber development: An immunofluorescent study. *Canadian Journal Of Botany* 64(7): 1373-1381.

- Lidell, M. C., G. A. Niles, et al. (1372). Response of nectariless cotton (*Gossypium hirsutum*) genotypes to cotton fleahopper (*Pseudatomoscelis seriatus*) (Heteroptera: Miridae) infestation. *Journal Of Economic Entomology* 79(5): 1372-1376.
- Brook, K. D., A. B. Hearn, et al. (1368). Response of cotton, *Gossypium hirsutum* L., to damage by insect pests in Australia: Manual simulation of damage. *Journal Of Economic Entomology* 85(4): 1368-1377.
- Zeringue, H. J. J. (1357). Changes in cotton leaf chemistry induced by volatile elicitors. *Phytochemistry* 26(5): 1357-1360.
- Brook, K. D., A. B. Hearn, et al. (1356). Response of cotton to damage by insect pests in Australia: Pest management trials. *Journal Of Economic Entomology* 85(4): 1356-1367.
- Reeves, D. W. and J. T. Touchton (1353). Relative phytotoxicity of dicyandiamide and availability of its nitrogen to cotton (*Gossypium hirsutum*), corn (*Zea mays*), and grain sorghum (*Sorghum bicolor*). *Soil Science Society Of America Journal* 50(5): 1353-1357.
- Turley, R. B. and R. N. Trelease (1350). Cottonseed malate synthase: Biogenesis in maturing and germinated seeds. *Plant Physiology* 84(4): 1350-1356.
- Zhushbaktova, Z. A., V. N. Syrov, et al. (1348). Study of the hypolipidemic properties of polysaccharides isolated from Central Asian flora. *Khimiko Farmatsevticheskii Zhurnal* 21(11): 1348-1352.
- French, N. M., II, S. B. Ramaswamy, et al. (1347). Effect of three adjuvants on coverage, persistence and efficacy of ground-applied chlorpyrifos for suppression of cotton aphid (Homoptera: Aphididae). *Journal Of Economic Entomology* 85(4): 1347-1355.
- Trelease, R. N., C. A. Hermerath, et al. (1343). Cottonseed malate synthase: Purification and immunochemical characterization. *Plant Physiology* 84(4): 1343-1349.
- Ramalho, F. S. and F. M. M. Jesus (1339). Abamectin: Efficiency against cotton aphid and effects on *Cycloneda sanguinea* and *Pseudodoros clavatus*. *Pesquisa Agropecuaria Brasileira* 23(12): 1339-1342.
- Kennedy, C. W. and J. E. Jones (1331). Evaluating quantitative screening methods for manganese toxicity in cotton genotypes. *Journal Of Plant Nutrition* 14(12): 1331-1340.
- Souza, J. G. D., N. E. D. M. Beltrao, et al. (1327). Effect of flower bud removal on assimilation, growth and mineral nutrition of cotton (*Gossypium hirsutum latifolium*). *Pesquisa Agropecuaria Brasileira* 19(11): 1327-1334.
- Bykova, E. V. and S. V. Lev (1317). The genotypic peculiarities of the callusogenesis process in cotton. *Genetika* 24(6): 1317-1320.
- Adams, F. and P. J. Hathcock (1305). Aluminum toxicity and calcium deficiency in acid subsoil horizons of 2 coastal plains soil series. *Soil Science Society Of America Journal* 48(6): 1305-1309.
- Hodges, H. F., V. R. Reddy, et al. (1302). Mepiquat chloride and temperature effects on photosynthesis and respiration of fruiting cotton. *Crop Science* 31(5): 1302-1308.
- Xu, C., L. C. Blair, et al. (1297). Characteristics of five new photoautotrophic suspension cultures including two *Amaranthus* spp. and a cotton strain growing on ambient carbon dioxide levels. *Plant Physiology* 88(4): 1297-1302.
- Bhargava, R. and R. C. Sachar (1293). Induction of acid phosphatase in cotton seedlings: Enzyme purification, subunit structure and kinetic properties. *Phytochemistry* 26(5): 1293-1298.
- Rickerl, D. H., W. B. Gordon, et al. (1281). The effect of starter fertilizers on cotton production in legume residues. *Communications In Soil Science And Plant Analysis* 18(11): 1281-1302.

Dugger, W. M. and R. L. Palmer (1270). Changes in the carbon-14 labeled cell wall components with chase time after incorporation of UDP carbon-14 glucose by intact cotton fibers. *Plant Physiology* 86(4): 1270-1275.

Womack, C. L. and M. F. Schuster (1259). Potato leafhopper (*Empoasca fabae*) (Homoptera: Cicadellidae) feeding on seedling cotton (*Gossypium hirsutum*). *Journal Of Economic Entomology* 79(5): 1259-1263.

Wullschleger, S. D. and D. M. Oosterhuis (1259). Photosynthetic carbon production and use by developing cotton leaves and bolls. *Crop Science* 30(6): 1259-1264.

Brown, S. M., T. Whitwell, et al. (1256). Conservation tillage systems for cotton (*Gossypium hirsutum*) production. *Soil Science Society Of America Journal* 49(5): 1256-1260.

Gulin, V. V., V. A. Kal'Chenko, et al. (1256). Gamma-induced null-mutations at the Est-1 locus in *Gossypium hirsutum* L. *Genetika* 25(7): 1256-1260.

Priestley, D. A. and A. C. Leopold (1252). Alleviation of imbibitional chilling injury by use of lanolin. *Crop Science* 26(6): 1252-1254.

Ramalho, F. D. S., W. L. Parrot, et al. (1236). Feeding sites of young stage tobacco budworm (*Heliothis virescens*) on cotton. *Ciencia E Cultura* 38(7): 1236-1241.

Fernandez, C. J., J. T. Cothren, et al. (1224). Partitioning of biomass in well-watered and water-stressed cotton plants treated with mepiquat chloride. *Crop Science* 31(5): 1224-1228.

De, S. J. G. and S. J. V. Da (1211). Partitioning of carbohydrates in annual and perennial cotton (*Gossypium hirsutum* L.). *Journal Of Experimental Botany* 38(192): 1211-1218.

Firoozabady, E. (1211). The effects of cell cycle parameters on the cell wall regeneration and cell division of cotton protoplasts (*Gossypium hirsutum*). *Journal Of Experimental Botany* 37(181): 1211-1217.

Kuznetsov, V. V., B. T. Kydyrov, et al. (1203). Heat shock-induced salt resistance in cotton plants: Participation of polyamines, ethylene and proline. *Fiziologiya Rastenii* 38(6): 1203-1210.

Rothrock, C. S. (1202). Influence of soil temperature, water, and texture on *Thielaviopsis basicola* and black root rot of cotton. *Phytopathology* 82(10): 1202-1206.

Gulin, V. V., L. P. Tsupina, et al. (1187). Influence of microwave radiation on pollen of cotton (*Gossypium hirsutum* L.). *Genetika* 27(7): 1187-1192.

Abdullaev, A., L. G. Gorenkova, et al. (1181). Effect of a mixture of fatty alcohols on some physiological and biochemical parameters in cotton plants. *Fiziologiya Rastenii* 38(6): 1181-1187.

Stadelbacher, E. A., J. E. Powell, et al. (1167). Parasitism of *Heliothis zea* and *Heliothis virescens* (Lepidoptera: Noctuidae) larvae in wild and cultivated host plants in the delta of Mississippi (USA). *Environmental Entomology* 13(5): 1167-1172.

Gwyn, J. J. and D. M. Stelly (1165). Method to evaluate pollen viability of upland cotton: Tests with chromosome translocations. *Crop Science* 29(5): 1165-1169.

Wilson, F. D. (1164). Pink bollworm (*Pectinophora gossypiella*) resistance, lint yield, and lint yield components of okra-leaf cotton (*Gossypium hirsutum*) in different genetic backgrounds. *Crop Science* 26(6): 1164-1167.

Oosterhuis, D. M. and S. D. Wullschleger (1154). Osmotic adjustment in cotton (*Gossypium hirsutum* L.) leaves and roots in response to water stress. *Plant Physiology* 84(4): 1154-1157.

- Villalobos, F. J. and E. Fereres (1153). Evaporation measurements beneath corn, cotton, and sunflower canopies. *Agronomy Journal* 82(6): 1153-1159.
- Beltrao, N. E. D., J. C. F. D. Santana, et al. (1147). Evaluation of cowpea cultivars under intercropping system with annual cotton. *Pesquisa Agropecuaria Brasileira* 21(11): 1147-1154.
- Nobrega, L. B. D., N. E. D. M. Beltrao, et al. (1145). Mixtures of herbicides on weed control in annual cotton. *Pesquisa Agropecuaria Brasileira* 22(11-12): 1145-1150.
- Wullschleger, S. D. and D. M. Oosterhuis (1141). Canopy development and photosynthesis of cotton as influenced by nitrogen nutrition. *Journal Of Plant Nutrition* 13(9): 1141-1154.
- Bowman, D. T. and J. E. Jones (1137). A diallel study of bract surface area/lint weight per boll ratio in cotton (*Gossypium hirsutum*). *Crop Science* 24(6): 1137-1141.
- Butler, G. D. J. and F. D. Wilson (1137). Activity of adult whiteflies (Homoptera: Aleyrodidae) within plantings of different cotton strains and cultivars as determined by sticky-trap catches. *Journal Of Economic Entomology* 77(5): 1137-1140.
- Kunce, C. M. and R. N. Trelease (1134). Heterogeneity of catalase in maturing and germinated cotton (*Gossypium hirsutum* cultivar Deltapine 62) seeds. *Plant Physiology* 81(4): 1134-1139.
- Li, R., D. M. Stelly, et al. (1128). Cytogenetic abnormalities in cotton (*Gossypium hirsutum* L.) cell cultures. *Genome* 32(6): 1128-1134.
- Musaev, D. A., A. T. Gafurov, et al. (1127). The signal characters and competitive ability of cotton inbred lines. *Genetika* 22(7): 1127-1137.
- Kennedy, C. W., W. C. J. Smith, et al. (1123). Root cation exchange capacity of cotton cultivars in relation to aluminum toxicity. *Journal Of Plant Nutrition* 9(8): 1123-1134.
- Acaster, M. A., W. S. Scott, et al. (1121). Carbamoyltransferase reactions in plants: A survey for enzymic diversity and the potential for herbicidal activity of transition state analogue inhibitors. *Journal Of Experimental Botany* 40(219): 1121-1126.
- Kohel, R. J. and J. A. Lee (1119). Genetic analysis of Egyptian glandless cotton (*Gossypium barbadense* cultivar Bahtim 110). *Crop Science* 24(6): 1119-1121.
- Wells, R. and W. R. J. Meredith (1119). Heterosis in upland cotton (*Gossypium hirsutum*): I. Growth and leaf area partitioning. *Crop Science* 26(6): 1119-1123.
- Womack, C. L. and M. F. Schuster (1118). Testing the reported positive growth response for cotton, *Gossypium hirsutum*, treated with aldicarb. *Journal Of Economic Entomology* 79(4): 1118-1120.
- Guinn, G., J. R. Dunlap, et al. (1117). Influence of water deficits on the abscisic acid and IAA contents of cotton flower buds and flowers. *Plant Physiology* 93(3): 1117-1120.
- Wallace, T. P. and Z. K. M. El (1114). Inheritance of resistance in three cotton cultivars to the HV1 isolate of bacterial blight. *Crop Science* 29(5): 1114-1119.
- Lascano, R. J., B. C. H. M. Van, et al. (1113). Energy and water balance of a sparse crop: Simulated and measured soil and crop evaporation. *Soil Science Society Of America Journal* 51(5): 1113-1121.
- Khaitova, Z. R., E. L. Milyaeva, et al. (1110). Hormonal regulation of growth and development in mutant lines of cotton. *Fiziologiya Rastanii* 36(6): 1110-1117.

Leigh, T. F., A. H. Hyer, et al. (1109). Observed population increase, nymphal weight gain, and oviposition nonpreference as indicators of *Lygus hesperus* (Hemiptera: Heteroptera: Miridae) resistance in glandless cotton (*Gossypium hirsutum*). *Journal Of Economic Entomology* 78(5): 1109-1113.

Nelson, E. B. (1108). Rapid germination of sporangia of *Pythium* spp. in response to volatiles from germinating seeds. *Phytopathology* 77(7): 1108-1112.

Hedin, P. A., W. L. Parrott, et al. (1107). Effects of cotton plant allelochemicals and nutrients on behavior and development of tobacco budworm. *Journal Of Chemical Ecology* 17(6): 1107-1122.

Kurth, E., G. R. Cramer, et al. (1102). Effects of sodium chloride and calcium chloride on cell enlargement and cell production in cotton roots. *Plant Physiology* 82(4): 1102-1106.

Mulrooney, J. E., P. A. Hedin, et al. (1100). Effects of PIX, a plant growth regulator, on allelochemical content of cotton (*Gossypium hirsutum* cultivar Stoneville 213) and growth of tobacco budworm (*Heliothis virescens*) larvae (Lepidoptera: Noctuidae). *Journal Of Economic Entomology* 78(5): 1100-1104.

Hills, M. J. and H. Beevers (1084). An antibody to the castor bean glyoxysomal lipase (62 kD) also binds to a 62-kD protein in extracts from many young oilseed plants. *Plant Physiology* 85(4): 1084-1088.

Cotty, P. J. (1082). Evaluation of cotton cultivar susceptibility to *Alternaria* leaf spot. *Plant Disease* 71(12): 1082-1084.

Elzen, G. W., H. J. Williams, et al. (1079). Quantification of volatile terpenes of glanded and glandless *Gossypium hirsutum* cultivars and lines by gas chromatography. *Journal Of Agricultural And Food Chemistry* 33(6): 1079-1082.

Armbrust, D. V. (1078). Effect of particulates (dust) on cotton (*Gossypium hirsutum* cultivar Dunn 120) growth, photosynthesis and respiration. *Agronomy Journal* 78(6): 1078-1081.

Eid, E. T., M. S. Ismail, et al. (1077). Effect of mepiquat chloride on McNair 220 cotton variety (*Gossypium hirsutum* L.) under Egyptian conditions. *Annals Of Agricultural Science* 31(2): 1077-1088.

Balasubramanian, G., S. Chelliah, et al. (1075). Effect of host plants on the biology of *Spodoptera litura*. *Indian Journal Of Agricultural Sciences* 54(12): 1075-1080.

Bariola, L. A., C. C. Chu, et al. (1074). Timing applications of plant growth regulators and last irrigation for pink bollworm (Lepidoptera: Gelechiidae) control. *Journal Of Economic Entomology* 83(3): 1074-1079.

Rasulov, B. K. (1066). Light-dependent control of the reaction of ribulose-1,5-bisphosphate carboxylation in intact cotton leaves. *Fiziologiya Rastenii* 38(6): 1066-1074.

Ivanishchev, V. V. (1065). Oxaloacetate decarboxylase activity in chloroplasts from C-3-plant leaves. *Fiziologiya Rastenii* 37(6): 1065-1071.

Burris, E., A. M. Pavloff, et al. (1064). Evaluation of two procedures for monitoring populations of early season insect pests (Thysanoptera: Thripidae and Homoptera: Aphididae) in cotton under selected management strategies. *Journal Of Economic Entomology* 83(3): 1064-1068.

Nobrega, L. B. D., D. J. Vieira, et al. (1061). Influence of node number and pruning height in herbaceous cotton (*Gossypium hirsutum*). *Pesquisa Agropecuaria Brasileira* 20(9): 1061-1068.

Ali, A., R. G. Luttrell, et al. (1060). Feeding sites and distribution of fall armyworm (Lepidoptera: Noctuidae) larvae on cotton. *Environmental Entomology* 19(4): 1060-1067.

Petersen, K. L., S. Moreshet, et al. (1059). Stomatal responses of field-grown cotton to radiation and soil moisture. *Agronomy Journal* 83(6): 1059-1065.

Radin, J. W., W. Hartung, et al. (1958). Correlation of stomatal conductance with photosynthetic capacity of cotton only in a carbon dioxide-enriched atmosphere: Mediation by abscisic acid? *Plant Physiology* 88(4): 1058-1062.

Hassidim, M., Y. Braun, et al. (1957). Studies on proton-translocating ATPase in plants of varying resistance to salinity: II. Potassium strongly promotes development of membrane potential in vesicles from cotton (*Gossypium hirsutum* cultivar Acala San Jose) roots. *Plant Physiology* 81(4): 1057-1061.

Hicks, S. K., C. W. Wendt, et al. (1957). Allelopathic effects of wheat straw on cotton germination, emergence and yield. *Crop Science* 29(4): 1057-1061.

Hedin, P. A., J. N. Jenkins, et al. (1955). Effects of bioregulators on flavonoids, insect resistance, and yield of seed cotton. *Journal Of Agricultural And Food Chemistry* 36(5): 1055-1061.

Kerbabaeva, Z. A., A. S. Konoplya, et al. (1955). Effects of marker genes on commercial characters in isolines of cotton *Gossypium hirsutum* L. and *Gossypium barbadense* L. *Genetika* 26(6): 1055-1063.

Hayashi, T., S. M. Read, et al. (1954). UDP-glucose: (1 fwdarw 3)-beta-glucan synthases from mung bean and cotton: Differential effects of calcium and magnesium on enzyme properties and on macromolecular structure of the glucan product. *Plant Physiology* 83(4): 1054-1062.

Barreiro, N. M., J. C. F. D. Santana, et al. (1949). Performance of herbaceous cotton lines in the state of Alagoas, Brazil. *Pesquisa Agropecuaria Brasileira* 21(10): 1049-1054.

Turley, R. B., S. M. Choe, et al. (1949). Characterization of a complementary DNA clone encoding the complete amino acid sequence of cotton isocitrate lyase. *Biochimica Et Biophysica Acta* 2: 223-226.

Abdel, A. M. H., M. S. Ismail, et al. (1947). Effect of growth retardant N,N,N-tributyl-3-(trifluoromethyl)benzenemethaniminium chloride (DOWCO 391X) on seed cotton yield and yield components. *Annals Of Agricultural Science* 31(2): 1047-1062.

Meredith, W. R. J. (1945). Yield and fiber-quality potential for second-generation cotton hybrids. *Crop Science* 30(5): 1045-1048.

Temple, P. J. (1945). Growth form and yield responses of four cotton cultivars to ozone. *Agronomy Journal* 82(6): 1045-1050.

Endrizzzi, J. E. and R. Nelson (1941). Linkage analysis and arm location of the open bud (ob-1) and yellow petal (Y-2) loci in chromosome 18 of cotton. *Genome* 32(6): 1041-1043.

Lutrick, M. C., H. A. Peacock, et al. (1941). Nitrate monitoring for cotton (*Gossypium hirsutum*) lint production on a typic paleudult. *Agronomy Journal* 78(6): 1041-1046.

Eck, H. V. (1935). Effects of water deficits on yield, yield components and water use efficiency of irrigated corn (*Zea mays*). *Agronomy Journal* 78(6): 1035-1040.

Mamood, A. N., G. D. Waller, et al. (1934). Dispersal of upland and Pima cotton pollen by honey bees (Hymenoptera: Apidae) visiting upland male-sterile flowers. *Environmental Entomology* 19(4): 1034-1036.

Ashworth, E. N., G. A. Davis, et al. (1933). Factors affecting ice nucleation in plant tissues. *Plant Physiology* 79(4): 1033-1037.

Parkhurst, D. F., S. C. Wong, et al. (1932). Gradients of intercellular carbon dioxide levels across the leaf mesophyll. *Plant Physiology* 86(4): 1032-1037.

Santana, J. C. F. D., E. O. D. Santos, et al. (1931). Evaluation of herbaceous cotton lines and cultivars in the northeast of Brazil: II. Agronomic and fiber characteristics. *Pesquisa Agropecuaria Brasileira* 21(10): 1031-1048.

Novick, R. G., J. E. Jones, et al. (1929). Genetic trait effects on nonlint trash of cotton. *Crop Science* 31(4): 1029-1034.

Mauney, J. R. and T. J. Henneberry (1927). Causes of square abscission in cotton (*Gossypium hirsutum*) in Arizona (USA). *Crop Science* 24(6): 1027-1030.

Pessarakli, M. and T. C. Tucker (1925). Ammonium (nitrogen-15) metabolism in cotton (*Gossypium hirsutum*) under salt stress. *Journal Of Plant Nutrition* 8(11): 1025-1046.

Lege, K. E., C. W. Smith, et al. (1924). Genotypic and cultural effects on condensed tannin concentration of cotton leaves. *Crop Science* 32(4): 1024-1028.

Waller, G. D., B. J. Estes, et al. (1922). Residual life and toxicity to honey bees (Hymenoptera: Apidae) of selected pyrethroid formulations applied to cotton in Arizona (USA). *Journal Of Economic Entomology* 81(4): 1022-1026.

Fry, K. E. (1920). Earliness factors in three pima cotton (*Gossypium barbadense*) genotypes. *Crop Science* 25(6): 1020-1023.

Turaev, A. M., A. N. Mun, et al. (1920). Isolation, fractionation and cultivation of cotton microspores. *Doklady Akademii Nauk Sssr* 311(4): 1020-1023.

Rosenthal, W. D., G. F. Arkin, et al. (1919). Water deficit effects on transpiration and leaf growth. *Agronomy Journal* 79(6): 1019-1026.

Turcotte, E. L. and R. G. Percy (1918). Inheritance of a second virescent mutant in American pima cotton. *Crop Science* 28(6): 1018-1019.

Correll, J. C., T. R. Gordon, et al. (1917). Vegetative compatibility and pathogenicity of *Verticillium albo atrum*. *Phytopathology* 78(8): 1017-1021.

Revenkova, E. V., A. S. Kraev, et al. (1917). Transformation of cotton (*Gossypium hirsutum* L.) by supervirulent *Agrobacterium tumefaciens* strain A281. *Molekulyarnaya Biologiya* 24(4): 1017-1023.

Kohel, R. J. (1916). Genetic analysis of a white mutant in cotton. *Crop Science* 28(6): 1016-1018.

Silva, M. D. A., J. R. P. Parra, et al. (1915). Compared biology of *Tetranychus urticae* in cotton (*Gossypium hirsutum*) cultivars: II. Life table of fertility. *Pesquisa Agropecuaria Brasileira* 20(9): 1015-1020.

Umbeck, P. F. and J. M. Stewart (1915). Substitution of cotton cytoplasm from wild diploid species for cotton germplasm improvement. *Crop Science* 25(6): 1015-1019.

Garas, N. A. and A. C. J. Waiss (1911). Differential accumulation and distribution of antifungal sesquiterpenoids in cotton stems inoculated with *Verticillium dahliae*. *Phytopathology* 76(10): 1011-1017.

Kittock, D. L., B. B. Taylor, et al. (1911). Partitioning yield reduction from early cotton planting. *Crop Science* 27(5): 1011-1015.

Chapman, K. D. and R. N. Trelease (1908). Acquisition of membrane lipids by differentiating glyoxysomes: Role of lipid bodies. *Journal Of Cell Biology* 115(4): 995-1008.

Yunuskhonov, S. H. (1906). The comparative study of cottonseed proteins of A and D genome groups in connection with the origin of *Gossypium hirsutum* and *Gossypium barbadense* allotetraploids. *Genetika* 20(6): 998-1006.

Garas, N. A., S. Wilhem, et al. (1905). Relationship of cultivar resistance to distribution of *Verticillium dahliae* in inoculated cotton plants and to growth of single conidia on excised stem segments. *Phytopathology* 76(10): 1005-1010.

Vaissiere, B. E., J. O. Moffett, et al. (1005). Honeybees (*Apis mellifera*) as pollinators for hybrid cottonseed (*Gossypium hirsutum*) production on the Texas (USA) high plains. *Agronomy Journal* 76(6): 1005-1010.

Geng, S., Q. Zhang, et al. (1004). Stability in yield and fiber quality of California cotton. *Crop Science* 27(5): 1004-1010.

McCormick, S. P., D. Bhatnagar, et al. (1002). An inhibitor of aflatoxin biosynthesis in developing cottonseed. *Canadian Journal Of Botany* 66(5): 998-1002.

Thomson, N. J. and D. J. Luckett (1002). Heterosis and combining ability effects in cotton: II. Heterosis. *Australian Journal Of Agricultural Research* 39(6): 991-1002.

Minton, E. B. and W. R. J. Meredith (1001). Root-knot nematode effect on nine cotton cultivars in Mississippi (USA). *Crop Science* 27(5): 1001-1004.

Rosolem, C. A. and D. S. Mikkelsen (1001). Potassium absorption and partitioning in cotton as affected by periods of potassium deficiency. *Journal Of Plant Nutrition* 14(9): 1001-1016.

Radin, J. W., J. R. Mauney, et al. (1000). Water uptake by cotton roots during fruit filling in relation to irrigation frequency. *Crop Science* 29(4): 1000-1005.

Aujla, M. S., C. J. Singh, et al. Comparison of various irrigation methods in cotton. *Journal of Research Punjab Agricultural University* 29(3): 374-377. {a} Regional Research Station, (PAU), Bathinda, India

Baloch, M. J., B. A. Soomro, et al. Stability analysis for comparing cotton varieties (*Gossypium hirsutum* L.). *Pakistan Journal of Scientific and Industrial Research* 37(12): 528-530. Cotton Res. Inst., Sakrand, Pakistan

Bhakare, H. A., A. S. Kulkarni, et al. Chemical characteristics, fatty acid and phospholipid composition of some cotton germplasm lines of *Gossypium hirsutum* and *Gossypium arboreum*: I. PKV Punjabrao Krishi Vidyapeeth Research Journal 16(2): 205-210. LIT, Nagpur Univ., Nagpur, India

Bhakare, H. A., A. S. Kulkarni, et al. Glycolipid composition of some germplasm lines of *Gossypium hirsutum* and *Gossypium arboreum*: II. PKV Punjabrao Krishi Vidyapeeth Research Journal 16(2): 211-214. Dep. Oil Technol., Laxminarayan Inst. Technol., Nagpur Univ., Nagpur, India

Buonauro, R. Preliminary investigations on lipoxygenase activity in cotton cotyledons inoculated by phytopathogenic bacteria. *Annali della Facolta di Agraria Universita degli Studi di Perugia* 48: 9-15. Istituto di Patologia Vegetale, Universita degli Studi di Perugia, Perugia, Italy

Chand, T. and R. K. Jain Chemical control of root-knot nematode (*Meloidogyne incognita*) through seed dressing in cotton (*Gossypium hirsutum*). *Indian Journal of Nematology* 20(2): 224-227. Dep. Nematol., Haryana Agric. Univ., Hisar 125004, India

Dadlani, M., V. Vashisht, et al. A comparison of field grow-out and electrophoresis methods for testing genetic purity of cotton hybrid seed. *Seed Research New Delhi* 22(2): 160-162. Div. Seed Sci. and Technol., Indian Agric. Res. Inst., New Delhi-110 012, India

Deshpande, L. A., R. M. Kokate, et al. Cytomorphological studies in induced tetraploid *Gossypium arboreum* and its interspecific hybrid with tetraploid *Gossypium hirsutum* L. *Indian Journal of Genetics and Plant Breeding* 51(2): 194-202. {a} Cotton Res. Station, Mahboob Baugh Farm, Marathwada Agric. Univ., Parbhani 431401, India

Gill, J. S., A. S. Sidhu, et al. Effect of temperature, humidity and food on *Earias vittella* (Fabricius) in cotton. *Journal of Research Punjab Agricultural University* 29(2): 216-218. Dep. Entomol., Punjab Agric. Univ., Ludhiana, India

Kale, P. S., N. R. Holey, et al. Fungicides for controlling seed borne infection in cotton seeds. Seed Research New Delhi 20(1): 60. Dep. Plant Pathol., Post Graduate Inst., Punjabrao Krishi Vidyapeeth, Akola 444 104, India

Katole, S. R., H. S. Thakare, et al. Occurrence of *Chiloloba acuta* Wied. (Cetoniinae-Coleoptera) on sorghum in Vidarbha, its damage and control. PKV Punjabrao Krishi Vidyapeeth Research Journal 16(2): 147-150. {a} Dep. Entomol., PKV, Akola, India

Keerio, M. D., M. S. Kalwar, et al. Biometrical analysis of some metrical traits of *Gossypium hirsutum* L. combining ability. Pakistan Journal of Scientific and Industrial Research 38(11-12): 454-456. Dep. Plant Breeding Genet., Agriculture University, Tandojam, Pakistan

Krishnadoss, D. and M. Kadambavanasundaram Genetic variability, heritability and genetic advance in tetraploid species of cotton. Indian Journal of Genetics and Plant Breeding 53(1): 55-59. Sch. Genetics, Tamil Nadu Agric. Univ., Coimbatore 641003, India

Luo, X. and J. M. Widholm Cryopreservation of photosynthetic plant cell suspension cultures. Plant Cell Tissue and Organ Culture 47(2): 183-187. {a} Dep. Crop Sci., Univ. Illinois, Edward R. Madigan Lab., 1201 W. Gregory, Urbana, IL 61801, USA

Mahmood, T., M. Arshad, et al. Response of some cotton varieties to cotton leaf curl virus. Pakistan Journal of Scientific and Industrial Research 38(1): 30-32. Central Cotton Res. Inst., Multan, Pakistan

Nadarajan, N. and S. R. S. Rangasamy Genetic analysis of certain fibre characters in *Gossypium hirsutum* L. Indian Journal of Genetics and Plant Breeding 52(3): 245-251. {a} Dep. Agric. Botany, Agric. College, Madurai 625104, India

Patil, S. M. and D. K. Morey Conservation of water through land surface modification and use of antitranspirants in cotton. PKV Punjabrao Krishi Vidyapeeth Research Journal 16(2): 180-184. {a} NARP, Sindewahi, India

Purohit, M. S. and A. D. Deshpande Effects of fertilizers on cotton bollworms in relation to plant protection. Journal of Maharashtra Agricultural Universities 19(2): 172-174. {a} Gujarat Agric. Univ., Navsari 396 450, India

Sandhu, B. S., M. S. Gill, et al. LD 491 variety of desi cotton (*Gossypium arboreum* L.). Journal of Research Punjab Agricultural University 32(2): 243. Dep. Plant Breeding, Punjab Agric. Univ., Ludhiana, India

Setatou, H. B. and A. D. Simonis Correction of PREVIEWS 98817852. Effect of time and rate of nitrogen application on cotton. Correction of publication year from 1996. Fertilizer Research 43(1-3): 49-53. {a} Soil Sciences Institute, 541 10 Thessaloniki, Greece

Siddiqui, M. A. and R. A. Patil Heterosis in crosses of *Hirsutum* cotton. Journal of Maharashtra Agricultural Universities 19(2): 241-244. Agril. Res. Stn., Badnapur, India

Singh, J., M. S. Mahal, et al. A plan for sampling cotton jassid and whitefly populations on *hirsutum* cotton. Journal of Research Punjab Agricultural University 32(1): 46-50. Dep. Entomol., Punjab Agric. Univ., Ludhiana, India

Soares, J. J. and F. M. Lara Feeding and oviposition preference of *Anthonomus grandis* Boheman, 1843, among cotton genotypes. Cientifica Jaboticabal 21(2): 333-338. {a} CNPA, EMBRAPA, 58100-970, Campina Grande, PB, Brazil

