


Nanded	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	10EC @7ml in 10 litres of water. Terminate the crop as early as possible. Avoid ratooning to break the PBW life cycle.
Parbhani	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Hingoli	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Buldhana	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Akola	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Washim	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Amravati	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Yavatmal	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Wardha	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Nagpur	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Chandrapur	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
A.P./										
TELANGANA										
Adilabad	0.0	0.0	0.0	0.0	3.0	0.0	0.0	0.0	0.0	At Guntur, the crop is 110 to 135 days and has entered boll bursting and picking stages. The weather was dry during the reporting period. Spraying against sucking pests (Thrips/ Jassid) and <i>Spodoptera</i> was taken up. In heavy soils, in farmers' fields, pickings are in Wherever irrigation facilities are available, farmers are giving light irrigation and they are suggested to be vigilant about pink bollworm incidence. Foliar application of nutrients (19:19:19, KNO ₃) given in irrigated crop. Spraying of 0.1% Carbendazim 50% WP was done to control Grey mildew. Advisory: If PBW observed more than 10 per cent, spray Fenvalerate 20 EC @ 5ml or Cypermethrin 10 EC @ 7ml in 10 litres of water only if plants have minimum 6-8 green bolls. Do not pick when cotton is wet following the rains. Store and pack dry cotton after drying for 2-3 days.
Warangal	0.0	0.0	0.0	21.0	4.0	0.0	0.0	0.0	0.0	
Khammam	0.0	0.0	0.0	27.0	3.0	0.0	0.0	0.0	0.0	
Karimnagar	0.0	0.0	0.0	11.0	3.0	0.0	0.0	0.0	0.0	
Mahabubnagar	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Guntur	0.0	0.0	0.0	3.0	0.0	0.0	0.0	0.0	0.0	
Prakasam	0.0	0.0	0.0	3.0	0.0	0.0	0.0	0.0	0.0	
KARNATAKA										
Dharwad	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	The crop is 175 to 180 days old, Second kapas picking was taken up. Kapas picking is almost over in crop sown during May and June first fortnight. Cleaning and grading of Kapas for marketing was also done during the reporting week. Suggested not to pick the kapas in early hours of the morning as the kapas is wet due to dew. Farmers were advised to sell the kapas, picking wise separately, to get higher price. Advisory: The cotton stalks may be used for compost making using suitable microbial cultures or to be rotoslashd in the field itself to incorporate in to the soil, which will enhance the soil fertility. Ratoon crop from the existing cotton crop is not desirable. The kapas should be dried under sunlight for a day. Kapas from each picking should be sent to market separately.
Haveri	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Mysore	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
TAMIL NADU										
Perambalur	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4.0	At Srivilliputhur, the crop is 75 to 115 DAS at boll formation to boll development stage. The weather during the reporting period was moderately hot without any rainfall. Weeding and top dressing of fertilizers in many parts of area taken up. Sucking pests (jassid, aphids and whitefly) and bollworms were observed in some areas. Root rot incidence was noticed in some areas. Picking started in some farmers' fields. Advisory:
Salem	0.0	0.0	0.0	0.0	0.0	0.0	0.0	9.0	0.0	
Trichy	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	3.0	
Virudhunagar	0.0	0.0	0.0	3.0	0.0	0.0	0.0	0.0	0.0	

										Application of Acetamiprid 20 % SP @100 g / ha suggested for the management of sucking pests. Farmers are advised to spray Chlorantraniliprole 18.5% SC @ 3ml/10 liter of water for the control of bollworms. To manage root rot disease, drenching of Carbendazim 50%WP @ 0.2% is recommended
--	--	--	--	--	--	--	--	--	--	--

Rainfall (mm) legend



0.0 mm rainfall (no rainfall) . Blank space express data Not available. Source : <http://imdagrmet.gov.in>