

ICAR-Central Institute for Cotton Research

Weekly Advisory for Cotton Cultivation from 21st to 26th July 2015 (36th Standard Week)

"The advisory is based on inputs received from the State Agricultural Universities of the respective states"

WEATHER ADVISORY

Date	July-15						Advisory
	21	22	23	24	25	26	
PUNJAB							Moderate to heavy Rains are expected this week and the subsequent week. Incidence of leafhopper incidence (1-8/3 leaves), whitefly (09-36/3leaves) and thrips (18-60/ 3 leaves) was observed. Whitefly incidence is likely to get reduced with the heavy rains. Leaf hopper and whitefly incidence can get aggravated with excessive urea application, especially if P and K are not available to plants. Leaf Curl Virus (CLCuD) has appeared at many locations and at few locations root rot and bacterial leaf blight problem was noticed after first irrigation. Square shedding has been observed due to cloudy weather. Do not spray against mealybugs since their parasitoids <i>Aenasius</i> are active now and mealybugs will be under natural control. Red hairy caterpillar is expected in some regions, particularly in the sandy areas. The damage is likely to be minimum and in patches. Recommended pest and disease control measures can be initiated only at economic threshold levels on any rain-free day. Before spray, weather forecast should be seen to avoid wash out of insecticide spray. Regular pest scouting is recommended.
Batinda	15	0	0	12	36	11	
Ferozepur	5	0	0	12	33	11	
Muktsar	2	0	0	6	44	6	
Mansa	3	0	0	7	52	8	
HARYANA							
Sirsa	5	0	0	6	33	9	
Hissar	3	0	0	7	56	0	
Fatehabad	4	0	0	6	35	12	
RAJASTHAN							Intermittent rains are expected. Incidence of leafhopper and whitefly is noticed. Recommended site specific remedial measure for root rot and parawilt may be initiated based on early symptoms. Strictly avoid spaying with pyrethroids or Fipronil. These insecticides will aggravate the whitefly problem. Neem oil sprays are preferred. Occurrences of cotton leaf curl virus disease and wilt (root rot) have been observed for which appropriate control measures are to be followed as per the recommended dose.
Hanumangarh	1	2	3	4	5	5	
Sri Ganganagar	4	5	6	7	10	6	
Banswara	3	5	6	8	6	6	
ORISSA							Good rains are expected during the week. Field operations may be difficult. However, wherever possible, weeding, earthing up and top dressing with N+P should be done. For green manuring, incorporate Sunhemp into the soil. Do not take up any insecticide sprays in view of the rains.
Koraput	9	17	26	18	22	27	
Kalahandi	3	9	15	20	14	14	
Balagiri	0	17	21	22	19	24	
GUJARAT							Good rains are predicted this week. Whitefly and leaf hopper infestation is expected to get reduced. Any insecticide sprays taken up this week will be completely wasted. Therefore do not spray. Wilt and root rot problems should be attended on priority with remedial measures recommended in the appendix. The crop is yet to enter boll formation stage, but pink bollworm may cause damage to flowers. though Bollgard-II bolls are resistant to pink bollworm, damage to flowers is not uncommon. Set up monitoring traps for pink bollworm. Check up for economic thresholds before contemplating any recommended management practices.
Amreli	18	11	30	10	4	3	
Bhavnagar	12	20	70	30	10	5	
Jamnagar	0	10	40	20	10	10	
Rajkot	8	24	50	20	10	10	
Baruch	34	30	60	10	5	10	
Sabarkantha	9	24	50	30	10	10	
Surendranagar	4	11	30	10	4	3	
Ahmedabad	16	24	60	0	5	5	
Vadodara	31	30	60	10	5	10	
Patan	0	5	30	0	0	3	
Mehsana	0	5	30	0	0	3	
MP							The crop is in vegetative stage. No incidence of insect- pests or

Khargaon	8	11	12	10	6	11	diseases has been recorded. However the crop is in water stress condition because there were no rains for the last twenty days. Intermittent rains are predicted this week. Top dressing with N+P is recommended during rain-free period. Wilt and root rot problems are expected after rains. These should be attended on priority with remedial measures recommended in the appendix of this advisory.
Dhar	17	8	28	23	0	16	
Khandwa	20	10	14	10	8	10	
MAHARASHTRA							
Nagpur	6	6	4	8	3	4	
Wardha	6	2	3	3	3	0	
Chandrapur	20	2	8	3	3	3	
Yavatmal	8	2	3	3	1	0	
Amravati	6	4	3	9	2	15	
Akola	6	2	3	2	5	4	
Buldhana	14	3	10	7	8	13	
Parbhani	12	3	0	5	5	4	
Nanded	5	3	0	5	7	5	
Beed	7	3	0	4	5	3	
Washim	10	2	2	2	4	5	
Dhule	12	31	3	5	0	22	
Jalgaon	10	10	19	10	14	13	
Jalna	9	0	0	0	6	0	
Aurangabad	3	16	7	5	5	10	
TELANGANA							
Adilabad	6	4	5	3	6	6	
Warangal	6	6	4	3	6	8	
Khammam	5	7	4	0	3	6	
Karimnagar	6	6	5	0	5	8	
Nalgonda	5	7	4	0	0	8	
AP							
Guntur	0	3	5	0	0	7	
Prakasam	0	10	7	2	2	2	
KARNATAKA							
Dharwad	3	4	7	7	5	7	
Haveri	2	3	9	8	9	5	
Mysore	13	3	3	7	5	6	
TAMILNADU							
Perambalur	2	0	0	2	0	0	
Salem	4	2	2	18	18	4	
Trichy	5	1	2	2	1	0	
Virudhunagar	1	1	2	2	1	0	

Weather is predicted to be cloudy with sporadic rains during the week. However revised predictions indicate heavy rains during the last week of July. Therefore weeding, inter-culture and top-dressing with N+P are strongly recommended to be completed during this week. There are no reports of insect pests or diseases. However, leaf hoppers (jassids) are reported from fields where only urea was applied. High levels of infestation were observed in hybrids susceptible to jassids especially under heavy urea application. Sporadic occurrence of safflower caterpillar *Perigea capensis* has been observed on cotton and soybean. The caterpillar is often confused as *Helicoverpa armigera*. There is no need for any control measure for this insect on cotton. Neem oil can be sprayed for sucking pest control.

Slight rains are predicted by the end of the week. Sowing should be taken up in irrigated regions. Wherever sowings were done, farmers are advised to take up inter-cultivation. Moisture conservation measures may be taken up. Crop sown after this week in rainfed farms is most likely to face moisture stress in the seedling stage. Though thrips and whiteflies may be observed, it is advisable to avoid any chemical sprays. Neem oil based sprays are recommended wherever leaf hoppers and whiteflies are seen on early sown crop. Field bunds of proposed cotton growing fields should be kept clean and weed free to avoid initial inoculum of Pest and Diseases. For summer sown Cotton, foliar application of 1 to 2% KNO₃ + 1% MgSO₄ is recommended twice at weekly intervals to mitigate yellowing, leaf reddening and stunted growth because of high temperature coupled with high wind velocity.

Weather is predicted to be cloudy with sporadic rains during this week and the subsequent few days. Hence, it is suggested to take up inter-culture operations, hoeing and weeding. Sowing of desi cotton varieties (Jayadhar, DDHC-11 and RAHS-14) may be continued till the end of July. If basal dose was not applied, fertilizer application is recommended on Desi cotton crop @ 40:25:25 N-P-K kg/ha sown during first fortnight of June. Sucking pests if observed should be managed based on recommendations in the advisory appendix.

Summer sown cotton -picking is in progress. Cloudy sky with sporadic and intermittent rainfall is predicted for this week and the subsequent period of 8-10 days.

Legend					
Rainfall in mm	< 5	5-20	20-50	50-80	> 80

MANAGEMENT STRATEGIES RECOMMENDED BY CICR

(Authored by K. R. Kranthi; No part of this advisory may be used in any form in any publication electronic or print or any other means without the permission of the author)

The strategies recommended in this brief note are based on results of experiments conducted by CICR and developed in consonance with various ecologically compatible guidelines issued by various National and Global agencies.

GENERAL CROP HEALTH MANAGEMENT PRACTICES

1. **Early maturing varieties or Bt-cotton hybrids** may be preferred in rain-fed regions.
2. **Early sowing** is preferred in rain-fed regions immediately after receiving the first showers of 80 mm rainfall.
3. **Sowing on ridges in rain-fed regions** especially in high density planting systems is most preferred.
4. **Bt-cotton hybrids** may be sown at 90 x 30 cm in rain-fed regions and at wider spacing under irrigation
5. **Non-Bt varieties** Suraj such as (CICR) NH 615 (VN-MAU, Parbhani), AKH 081 (Dr PDKV Akola), Phule Dhanwantari (MPKV Rahuri) and Anjali (LRK 516) are early maturing. If these varieties are sown before 15th June in high density planting at 60x10 cm (40x10cm for Phule Dhanwantari), the crop will escape drought stress and bollworms.
6. **Intercropping in high density non-Bt cotton varieties** can be taken up with soybean (seed treated with *Bradyrhizobium japonicum*), cowpea or blackgram in alternate rows at 45 cm row to row and 10 cm plant to plant.
7. **Intercropping in Bt hybrids** can be taken up with soybean (seed treated with *Bradyrhizobium japonicum*), cowpea or blackgram as one row between two Bt-hybrid rows
8. **Border rows (2-3 rows) of pigeonpea** around cotton fields will prevent infestation of mealy bugs and serve as refugia.
9. **Farm Yard Manure @ 5 to 10 t/ha** or compost should be applied just after the first rain.
10. **Azotobacter and PSB @ 25 g each / kg seed** should be used for nutrients fixation.
11. **Optimize nutrient management** for macro and micronutrients. Foliar spray of MgSO₄, 2% Urea followed by 2% DAP, to ensure proper Cry1Ac expression and also to reduce problems of leaf reddening. Sprays of 1% cobalt chloride and soil drenching with Bavistin 1 % in the initial stage of wilt was found to help in the recovery of plants.
12. **Prevention of Leaf Reddening:** Spray 2 % urea, 0.5% Zinc Sulphate and 0.2 % Boron, twice at 15 days interval on 90 days old crop.
13. **Retention of squares and flowers:** Spray Planofix 4.5 SL (NAA) hormone @ 21 ppm (7 ml per 15 litres of water).

INSECT PEST MANAGEMENT

General recommendations

DOs

1. **Select sucking pest resistant varieties/hybrids.** Sucking pest resistant Bt hybrids may require very few insecticide interventions.
2. **Inter-crop with cowpea or sorghum or soybean or blackgram** to encourage predators of sucking pests.
3. **Imidacloprid (8 g), Vitavax or Thiram (3 g)** per kg seed will protect varieties against sucking pests and diseases.
4. **Use nitrogenous fertilizers to the minimum** especially for sucking pest susceptible varieties.
5. **Maintain field sanitation** (weed free)
6. **Remove and destroy mealy bug infested plants.**
7. **Use Neem preparations and biological control options** for least disruptive pest management.
8. **Pheromone traps** are efficient for pest monitoring of Pink bollworm.
9. **Stem application or soil application** (near the root zone) of Imidacloprid, Dimethoate or Acephate at 30-40 DAS and 50-60 DAS for effective eco-friendly control of thrips, mirid bugs, mealy bugs and other sucking pests.

DON'Ts

10. **Avoid late sowing beyond 15th May in North India** to prevent aggravation of cotton leaf curl virus.
11. **As far as possible avoid chemical insecticides during the first two months of the crop** to conserve naturally occurring biological control. Ladybird grubs and beetles, *Chrysoperla* grubs and adults, Syrphid flies, *Geocoris* grubs and bugs, *Aenasius* spp., *Aphilinus* grubs and wasps, mirid bugs and Spiders are the most important naturally occurring predators and parasitoids that effectively control aphids, jassids, thrips, mirids, whiteflies and mealybugs.
12. **Do not spray against minor lepidopteran insects** such as the cotton leaf folder, *Sylepta derogata* and cotton semilooper, *Anomis flava*. The larvae cause negligible damage to cotton but serve as hosts for parasitoids such as *Trichogramma* spp., *Apanteles* spp and *Sysiropa formosa*, that attack *H. armigera* and other bollworms.
13. **Do not spray Bt-formulations on Bt cotton** to avoid further selection pressure.
14. **Avoid foliar application of neonicotinoid insecticides** such as Acetamiprid, Imidacloprid, Clothianidin and Thiomethoxam which are likely to aggravate insect resistance, since hybrid cotton seeds are treated with imidacloprid.
15. **Do not use WHO Class-I (Extremely Hazardous category) insecticides** such as Phosphamidon, Methyl parathion, Phorate, Monocrotophos, Dichlorvos, Carbofuran, Methomyl, Triazophos and Metasystox.
16. **Avoid Fipronil and Pyrethroids** to prevent whitefly outbreaks.
17. **Avoid insecticide mixtures.** Mixtures severely disrupt eco-systems thereby leading to pest outbreaks.

SUCKING PEST MANAGEMENT

Economic Threshold Level (ETL): If whitefly and/or leafhopper damage reaches economic threshold levels of grade-II damage of curling and crinkling of lower leaves and yellowing of margins in 25% plants or more, any one of the following pest control measures as suggested below can be used.

- a. Neem oil 1.0% + Neem Seed Kernel Extract 5.0% + 0.05-0.1% detergent
- b. *Verticillium lecanii* 10gms/lit of water, wherever good formulations are available from reliable manufacturers
- c. Diafenthiuron 50WP 800g /ha,
- d. Flonicamid 50 WG 200g a.i/ha or
- e. Buprofezin 25% SC 200 g a.i/ha.

Insecticides such as Dimethoate or Acephate or Ethion can also be used but may be considered as alternatives only, in view of factors that relate to ecological and environmental safety, efficacy and resistance.

If mirid bugs are observed to cause economic damage to squares, it is advised to spray Acephate 75 SP @ 1 g/lit or Dimethoate.

BOLLWORM MANAGEMENT

Bt cotton is effective in controlling bollworms.

The following strategies are being recommended for non-Bt cotton

At Economic Threshold Levels (ETLs) of 50% infested plants (plants having flared squares with entry hole) **for *Helicoverpa armigera*.**

1. **Use HaNPV** (*Helicoverpa armigera* Nuclear Polyhedrosis Virus) on Bt-cotton followed by the application of **5% Neem Seed Kernel Extract (NSKE)** a week later. **OR, use Phosalone** at ETL for the management of bollworms, *Spodoptera* or whitefly.
2. *Trichogramma*, if available, can be used on non-Bt genotypes at 70-80 DAS. Avoid *Trichogramma* egg parasitoid releases on Bt-cotton since maximum neonates get killed on Bt-cotton and with *Trichogramma* application becoming superfluous.
3. **Insecticides effective on Bollworms**, especially *Helicoverpa armigera*.
 - a. Chlorantraniliprole,
 - b. Flubendiamide,
 - c. Spinosad,
 - d. Emamectin benzoate and
 - e. Indoxacarb

These insecticides have a high selective toxicity towards the target pests while being less toxic to many beneficial insects in the cotton ecosystem. These insecticides are ideally suited in eco-sustainable insecticide resistance management programmes.

4. **Pink bollworm and Spotted bollworms:** ETL level of one live larva in 10 green bolls or 8 moths per night for three consecutive nights. Spray Quinalphos 25 EC Profenophos 50 EC @ 2 ml/lit of water / Spray of Thiodicarb 75 WP @ 20 g or any pyrethroid.

OTHER PESTS

1. *Spodoptera litura*: Collection of egg masses or application of *SINPV* (*Spodoptera litura* Nuclear Polyhedrosis Virus) @ 500 LE/ha or Spray 200 ml Novaluron 10 EC or 250g Thiodicarb 75WP in 250 litres of water per acre
2. To minimize **shoot weevil** damage, spray Profenofos @ 2 ml/lit
3. **Snail incidence in heavy rainfall areas:** baiting with 2% Metaldehyde (Snail kill) @ 12.5 kg/ha has to be taken up and it is to be applied at the hideouts of the snails, on the bunds and to the soil around the crop where the damage is seen

DISEASE MANAGEMENT

Parawilt or Sudden drying (New wilt) or Wilt / Root rot: Symptoms are noticed in some fields after drought followed by rains or irrigation. Spray cobalt chloride @10mg/litre (10ppm) on affected plants within few hours of onset of symptoms and/or Drench plants with a mixture of Copper-Oxy-Chloride 25g and 200g Urea in 10 ltr of water or Carbendazim 1g/L.

Boll Rot: Generally early formed lower bolls rot due to cloudy and drizzling conditions.

Spray Mancozeb 75 WP + Chlorothalonil 70 WP each @ 2 g/lit of water. For better results, mix 10g Selvet 99 or 50 ml Triton in 100 litres of fungicide solution.

Alternaria blight: spray Mancozeb@2.5 g per one litre of water.

Myrothecium leaf spot disease and/or Bacterial blight: Spray Streptomycin sulphate (15-20 g/ha) plus Copper oxychloride (1500-2000 g/ha) in 200-250 L of water.

WEED MANAGEMENT

Application of Pre-emergence weedicide Stomp 30EC or Basalin @45EC 2.5 lt/ha and harrow immediately to prevent degradation.

Herbicides are most effective on younger weeds.

Post emergence herbicides (application rate 50 to 75 g ai/ha)

Grasses: Spray Quizalofop-ethyl or Fenoxaprop ethyl or Fluazifop butyl,

Sedges and grasses: Spray Propaquizafop ethyl

Broadleaf weeds: Spray Pyriithiobac sodium

Post-emergence weedicides would provide effective and timely control especially when interculture operations or manual weeding becomes difficult in wet soils. Weedicides are effective against younger (less than 10-15 days old) weeds, especially grasses. For grassy weeds, Quizalofop ethyl, Fenoxaprop ethyl, fluazifop butyl, can be used. For sedges and grasses, Propaquizafop ethyl is effective. Pyriithiobac sodium is effective on broad leaf weeds. Farmers may consult the technical experts of the Agricultural Universities for further details.

WATER LOGGING MANAGEMENT

Cotton is very sensitive to excess water. In many parts of Central and South India, water logging can be problematic due to excess rains. Cotton grown on deep black soils and ill drained conditions is worst affected due to water logging. Provide adequate drainage channels or water ways (particularly in heavy soils) along the slope of the land for draining excess water under heavy rainfall situations. For better soil moisture conservation, preferably in areas where rainfall is 700-900mm, the land can be reshaped into ridges and furrows with the help of a ridge plough or a bund former. This technique and sowing cotton on ridges would conserve rainwater and the furrows acts drainage channels whenever heavy rains are received particularly in heavy clays.

Drainage channels must be opened up along the field borders so that excess water is removed from the fields. If sowing hasn't yet been completed, it is strongly recommended that to take up sowing immediately on ridges and furrows by planting on top of ridges. Heavy rains will not affect the crop because the furrows will drain away excess water. Apply fertilizers if the crop becomes pale due to water logging. If heavy rains are forecast, fertilizer application may be postponed so as to prevent losses due to surface run-off.

Foliar sprays with 0.5 to 1.0% DAP or 19:19:19 (soluble complex of Nitrogen) at weekly intervals will help the plants to recover from the effect of water logging.

Weekly weather Advisory Report Coordinating Team

Scientists	Address
Dr K R Kranthi	Director, CICR, Nagpur
Dr A H Prakash	PC and Head, CICR, Regional station, Coimbatore
Dr. D Monga	Head, CICR, Regional station, Sirsa
Dr. S. B. Singh	Head, Division of Crop Improvement, CICR, Nagpur
Dr Sandhya Kranthi	Head, Division of Crop Protection, CICR, Nagpur
Dr Blasé De souza	Head, Division of Crop Production, CICR, Nagpur
Dr. Isabell Agrawal	Sr. Scientist CICR, Coimbatore
Sh. M.Sabesh	Scientist, CICR, Coimbatore

Scientists in-charge for weather report (AICRP centres)

Scientists	Address	Phone Nos.	E-mail id
Dr. Pankaj Rathore	Punjab Agricultural University, Faridkot, Punjab	09464051995	pankaj@pau.edu
Dr (Ms) Suneet Pandher	Punjab Agricultural University, Faridkot, Punjab	09814513681	suneet@pau.edu
Dr.Sanjeev Kumar	Punjab Agricultural University, RRS,		k.sanjeev@pau.edu

Kataria	Bhatinda		
Dr. Jagdish Beniwal	CCS-Haryana Agricultural University, Hisar Haryana	09416325420	jbeniwal2016@gmail.com
Dr. Rishikumar.	CICR Regional Station, Sirsa, Haryana	09729106299	rishipareek70@yahoo.co.in
Dr. Roop Singh Meena	Swami Keshwanand Rajasthan Agricultural University, Sriganganagar, Rajasthan	09413024080	rsmeenars@gmail.com
Dr.B.S.Nayak	Orissa University of Agriculture & Technology, Bhubaneswar, Orissa	09437321675	bsnayak2007@rediffmail.com
Dr. Gofaldu	Navsari Agricultural University, Navsari, Gujarat	09662532645	girishfaldu@rediffmail.com
Dr A. N Paslawar	Panjabrao Deshmukh Krishi Vidyapeeth, Akola , Maharashtra	09822220272	adinathpaslawar@rediffmail.com
Arvond D. Pandagale	Marathwada Agricultural University, Nanded, Maharashtra	07588581713	arvindpandagale@yahoo.co.in
Dr. Satish Parsai	RVS Krishi Vishvva Vidhyalaya, Gwalior, Madhya Pradesh	09406677601	aiccipkhandwa@gmail.com
Dr.S.Bharathi	Acharya N. G. Ranga Agricultural University, LAM, Guntur, AP	0949072341	bharathi_says@yahoo.com
Dr.Aladakatti	University of Agricultural Sciences, Dharwad , Karnataka	09448861040	yaladakatti@rediffmail.com
Dr. M.Y.Ajaykumar	University of Agricultural Sciences Raichur, Karnataka	09880398690	dr.my.ajay@gmail.com
Dr. S. Somasundaram	Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu	09965948419	rainfed@yahoo.com
Dr.M.Gunasekaran	Tamil Nadu Agricultural University, Cotton Research Station, Srivilliputhur, Tamil Nadu	09443631359	gunasekaran.pbg@gmail.com