

## Central Institute for Cotton Research, Nagpur

### Fifteenth Weekly Advisory for Cotton Cultivation 1<sup>st</sup> to 7<sup>th</sup> September 2014

"The advisory is based on inputs received from the State Agricultural Universities of the respective states"

**Weed management:** Wherever weeds have emerged, weedicides would provide effective and timely control. Weedicides are effective against younger (less than 10-15 days old) weeds, especially grasses. For grassy weeds, Quizalofop ethyl, Fenoxaprop ethyl, fluazifop butyl, can be used. For sedges and grasses, Propaquizafop ethyl is effective and Pyriithiobac sodium is effective on broad leaf weeds. Farmers may consult the technical experts of the Agricultural Universities for further details.

**Water logging:** Cotton is very sensitive to excess water. In many parts of Central and South India, water logging can be problematic due to excess rains. Cotton grown on deep black soils and ill drained conditions is worst affected due to water logging. Provide adequate drainage channels or water ways (particularly in heavy soils) along the slope of the land for draining excess water under heavy rainfall situations. For better soil moisture conservation, preferably in areas where rainfall is 700-900mm, the land can be reshaped into ridges and furrows with the help of a ridge plough or a bund former. This technique and sowing cotton on ridges would conserve rainwater and the furrows acts drainage channels whenever heavy rains are received particularly in heavy clays.

Drainage channels must be opened up along the field borders so that excess water is removed from the fields. If sowing hasn't yet been completed, it is strongly recommended that to take up sowing immediately on ridges and furrows by planting on top of ridges. Heavy rains will not affect the crop because the furrows will drain away excess water. Apply fertilizers if the crop becomes pale due to water logging. If heavy rains are forecast, fertilizer application may be postponed so as to prevent losses due to surface run-off.

Foliar spray with 0.5 to 1.0% DAP at weekly intervals will help the plants to recover from the effect of water logging.

#### Cotton Sown Area (As on 22-08-2014)

S.No.	States	Normal Area (DES)*	Area in lakh ha		
			Bt	Non Bt	Total
1.	Andhra Pradesh + Telengana	19.830	18.559	1.669	20.228
	Andhra Pradesh (23.95%)	4.749	5.611	0.239	5.850
	Telangana (76.05%)	15.081	12.948	1.430	14.378
2.	Gujarat	26.490	26.880	2.930	29.810
3.	Haryana	5.640	6.310	0.080	6.390
4.	Karnataka	5.270	6.520	0.630	7.150
5.	Madhya Pradesh	6.390	5.445	0.285	5.730
6.	Maharashtra	39.160	37.398	1.822	39.220
7.	Odisha	0.970	0.000	1.240	1.240
8.	Punjab	5.170	4.300	0.200	4.500
9.	Rajasthan	4.000	3.950	0.208	4.158
10.	Tamil Nadu	1.250	0.056	0.014	0.070
11.	Uttar Pradesh	0.010	0.000	0.260	0.260
12.	Others	0.350		0.050	0.050
	<b>All INDIA</b>	<b>114.530</b>	<b>109.418</b>	<b>9.388</b>	<b>118.806</b>

\* Directorate of Economics & Statistics, DAC, Ministry of Agriculture, Krishi Bhavan, New Delhi

## Weather forecast for to '2014

Zones/ Weather parameter	Temperature ( Min, Max)						Rainfall					
	09/09	10/09	11/09	12/09	13/09	14/09	09/09	10/09	11/09	12/09	13/09	14/09
States												
Punjab	25,35	25,35	25,35	25,35	25,35	25,35	Partly cloudy sky					
Haryana	26,36	26,36	26,36	25,36	25,35	25,35	Partly cloudy sky					
Rajasthan	27,36	28,35	28,35	28,35	28,35	28,35	Partly cloudy sky with thundery development					
Gujarat	26,32	25,32	25,31	24,31	24,31	25,32	Light rain					
Maharashtra	23,27	23,28	24,30	24,30	23,29	23,29	Moderate rain	Light rain	Partly cloudy sky with thundery development		Light rain	
M.P.	22,28	22,28	22,28	22,29	23,29	23,29	Mainly or Generally cloudy sky with possibility of rain or thunderstorm					
Odisha	19,27	19,29	20,29	20,29	20,29	20,29	Generally cloudy sky with possibility of rain or thunderstorm					
A.P.	23,31	23,31	23,32	23,32	23,32	23,32	Mainly or Generally cloudy sky with possibility of rain or thunderstorm					
Karnataka	21,29	22,30	21,30	22,30	22,30	22,30	Light rain					
Tamil Nadu	23,32	23,32	23,32	23,32	23,32	23,32	Light rain					

Source: [www.imd.gov.in](http://www.imd.gov.in)

## STRATEGIES FOR MANAGEMENT OF PESTS, DISEASES &amp; WEEDS

## INSECT PEST MANAGEMENT

## General recommendations

## DOs

1. Select sucking pest resistant varieties/hybrids. Sucking pest resistant Bt hybrids may require very few insecticide interventions.
2. Inter-crop with cowpea or sorghum or soybean or blackgram to encourage predators of sucking pests.
3. Seed treatment with Imidacloprid @7gms/Kg of seed.
4. Use nitrogenous fertilizers to the minimum especially for sucking pest susceptible varieties.
5. Maintain field sanitation (weed free) and remove and destroy mealy bug infested plants &.
6. **Stem application or soil application** (near the root zone) of Imidacloprid, Dimethoate or Acephate at 30-40 DAS and 50-60 DAS for effective eco-friendly control of thrips, mirid bugs, mealy bugs and other sucking pests.

## DON'Ts

7. **If possible avoid chemical insecticides during the first two months of the crop** to conserve naturally occurring biological control. Ladybird grubs and beetles, *Chrysoperla* grubs and adults, Syrphid flies, *Geocoris* grubs and bugs, *Aenasius* spp., *Aphelinus* grubs and wasps, mirid bugs and Spiders are the most important naturally occurring predators and parasitoids that effectively control aphids, jassids, thrips, mirids, whiteflies and mealybugs.
8. **Do not spray against minor lepidopteran insects** such as the cotton leaf folder, *Sylepta derogata* and cotton semilooper, *Anomis flava*. The larvae cause negligible damage to cotton but serve as hosts for parasitoids such as *Trichogramma* spp., *Apanteles* spp and *Sysiropa formosa*, that attack *H. armigera* and other bollworms.

9. **Do not spray Bt-formulations on Bt cotton** to avoid further selection pressure.
10. **Avoid foliar application of neonicotinoid insecticides** such as Acetamiprid, Imidacloprid, Clothianidin and Thiomethoxam which are likely to aggravate insect resistance, since hybrid cotton seeds are treated with imidacloprid.
11. **Do not use WHO Class-I (Extremely Harzardous category) insecticides** such as Phosphamidon, Methyl parathion, Phorate, Monocrotophos, Dichlorvos, Carbofuran, Methomyl, Triazophos and Metasystox.

### SUCKING PEST MANAGEMENT

**Economic Threshold Level (ETL):** If whitefly and/or leafhopper damage reaches economic threshold levels of grade-II damage of curling and crinkling of lower leaves and yellowing of margins in 25% plants or more, any one of the following pest control measures as suggested below can be used.

- a. Neem oil 1.0% + Neem Seed Kernel Extract 5.0% + 0.05-0.1% detergent
- b. *Verticillium lecanii* 10gms/lit of water, wherever good formulations are available from reliable manufacturers
- c. Diafenthiuron (50WP 800g /ha),
- d. Flonicamid 50 WG 200g a.i/ha or
- e. Buprofezin 25% SC 200 g a.i/ha.

Insecticides such as Fipronil or Dimethoate or Acephate or Ethion can also be used but may be considered as alternatives only, in view of factors that relate to ecological and environmental safety, efficacy and resistance.

If mirid bugs are observed to cause economic damage to squares, it is advised to spray Acephate 75 SP @ 1 g/lit or Fipronil 5 SC @ 1.0 ml/lit of water

### BOLLWORM MANAGEMENT

Bt cotton is effective in controlling bollworms.

*The following strategies are being recommended for non-Bt cotton*

At Economic Threshold Levels (ETLs) of 50% infested plants (plants having flared squares with entry hole) for *Helicoverpa armigera*.

1. **Use HaNPV on Bt-cotton** followed by the application of **5% NSKE** a week later. **Or, use Phosalone** at 50% bollworm infested plants (plants having flared squares with entry hole) or for the management of *Spodoptera* or whitefly.
2. ***Trichogramma***, if available, can be used on non-Bt genotypes at 70-80 DAS. Avoid *Trichogramma* egg parasitoid releases on Bt-cotton since maximum neonates get killed on Bt-cotton and with *Trichogramma* application becoming superfluous.
3. **Insecticides effective on Bollworms**, especially *Helicoverpa armigera*.
  - a. Chlorantraniliprole (Coragen),
  - b. Flubendiamide (Fame),
  - c. Spinosad,
  - d. Emamectin benzoate and
  - e. Indoxacarb

These insecticides have a high selective toxicity towards the target pests while being less toxic to many beneficial insects in the cotton ecosystem. These insecticides are ideally suited in eco-sustainable insecticide resistance management programmes.

4. **Pink bollworm and Spotted bollworms:** ETL level of one live larva in 10 green bolls or 8 moths per night for three consecutive nights. Spray Quinalphos 25 EC Profenophos 50 EC @ 2 ml/lit of water / Spray of Thiodicarb 75 WP @ 20 g or any pyrethroid.
5. ***Spodoptera litura*:** Collection of egg masses or application of SNPV (*Spodoptera litura* Nuclear Polyhedrosis Virus) @ 500 LE/ha or Spray 200 ml Rimon 10 EC or 250g Larvin 75WP in 250 litres of water per acre
6. To minimize **shoot weevil** damage, spray Profenofos @ 2 ml/lit
7. In case of snail incidence in heavy rainfall areas, baiting with 2% Metaldehyde (Snail kill) @ 12.5 kg/ha has to be taken up and it is to be applied at the hideouts of the snails, on the bunds and to the soil around the crop where the damage is seen

### DISEASE MANAGEMENT

1. **Parawilt or Sudden drying (New wilt) or Wilt / Root rot:** Symptoms are noticed in some fields after drought followed by rains or irrigation. Spray cobalt chloride @10mg/litre (10ppm) on affected plants within few hours of onset of symptoms and/or Drench plants with a mixture of Copper-Oxy-Chloride 25g and 200g Urea in 10 ltr of water or Carbendazim 1g/L.
2. **Boll Rot:** Generally early formed lower bolls rot due to cloudy and drizzling conditions. Spray Mancozeb 75 WP + Chlorothalonil 70 WP each @ 2 g/lit of water. For better results, mix 10g Selvet 99 or 50 ml Triton in 100 litres of fungicide solution.
3. ***Alternaria* blight:** spray Mancozeb@2.5 g per one litre of water.
4. **Myrothecium leaf spot disease and/or Bacterial blight:** Spray Streptomycin sulphate (15-20 g/ha) plus Copper oxychloride (1500-2000 g/ha) in 200-250 L of water.

### WEED MANAGEMENT

Herbicides are most effective on younger weeds.

**Post emergence herbicides (application rate 50 to 75 g ai /ha)**

1. **Grasses:** Spray Quizalofop-ethyl or Fenoxaprop ethyl or Fluazifop butyl,
2. **Sedges and grasses:** Spray Propaquizafop ethyl
3. **Broadleaf weeds:** Spray Pyriithiobac sodium

### GENERAL CROP HEALTH MANAGEMENT

1. **Optimize nutrient management** for macro and micronutrients. Foliar spray of MgSO<sub>4</sub>, 2% Urea followed by 2% DAP, to ensure proper Cry1Ac expression and also to reduce problems of leaf reddening. Sprays of 1% cobalt chloride and soil drenching with Bavistin 1 % in the initial stage of wilt was found to help in the recovery of plants.
2. **Prevention of Leaf Reddening:** Spray 2 % urea, 0.5% Zinc Sulphate and 0.2 % Boron, twice at 15 days interval on 90 days old crop.
3. **Retention of squares and flowers:** Spray Planofix 4.5 SL (NAA) hormone @ 21 ppm (7 ml per 15 litres of water).

**DROUGHT MANAGEMENT****Odisha****Mid season drought (long dry spell )**

Condition			Suggested Contingency Measures	
Mid season drought (long dry spell)	Major Farming Situation	Normal Crop / Cropping System	Crop management	Soil nutrient & moisture conservation measures
At flowering/ fruiting stage	Red soil High rainfall Medium elevation	Cotton	Applying of Planofix hormone * spraying the crop with Imidacloprid for controlling of sucking pests	Apply 1250ml micronutrient/ha
	Red and Yellow soil High rainfall Medium elevation	Cotton		
	Black soil High rainfall Medium elevation	Cotton + Arhar		

**Terminal drought (Early withdrawal of monsoon)**

Condition			Suggested Contingency Measures	
Terminal drought (Early withdrawal of monsoon)	Major Farming Situation	Normal Crop / Cropping System	Crop management	Rabi Crop planning
	Red soil, High rainfall, Medium elevation	Cotton	Provide protective irrigation	Mulch with stovers Dibble rabi crop
		Cotton	Provide protective irrigation	
		Cotton + Arhar	Provide protective irrigation Harvest at physiological maturity stage	

In case of severe drought situation following measures may be taken apart from the above contingent plans.

- Opting for castor, sesamum and safflower (mid *Rabi* crop) (as per recommendations specific to the agro eco region) instead of cotton to manage severe drought situation in *Kharif season*.
- In the event of late planting of cotton due to delayed onset of monsoon, maintenance of higher plant population and optimum input management to the extent possible is suggested.
- Bacterial blight, fungal foliar spots and parawilt are the major diseases in this region. Long dry spells with intermittent rains may aggravate the incidence of the diseases. These should be managed with fungicidal sprays as and when required.
- Foliar spray of KCl or KNO<sub>3</sub> to partially alleviate moisture stress during drought.
- Application of anti-transpirants or hormones.

**Gujarat****1<sup>st</sup> -15<sup>th</sup> September (Terminal drought)**

- Weeding
- Top dressing of Urea if sufficient occurrence of rain
- Insecticide spray for control of sucking and bollworms

- Alternate furrow irrigation, if irrigation water is available
- Adopt topping to reduce evapotranspiration losses

### IMPORTANT NOTE: (PEST MANAGEMENT)

Farmers are advised not to spray pyrethroids early in the season singly or in combination against sucking pests such as the whiteflies not only for cotton but also on other *H. armigera* host plants such as soybean, as it may exacerbate bollworm problems in non Bt cotton, wherever cultivated.

### COTTON CROP SITUATION

(Based on inputs received from the State Agricultural Universities of the respective States)

#### NORTH INDIA

**Punjab:** The crop is in peak boll development stage. The infestation of whitefly is very high in cotton growing areas of Punjab which needs immediate spray of Spiromesifen 240SC (Oberon) @ 200ml or Difenthiuron 50WP (Polo) @ 200g/acre. The incidences of cotton leaf curl virus disease have also been noticed in districts of Bathinda, Mansa in addition to traditional infected areas of Abohar. The infestation of *Spodoptera litura*, American boll worm and spotted boll worm was recorded on *desi* cotton and non-Bt varieties of American cotton. Farmers are advised to visit and monitor their cotton field for insect pests and diseases regularly so that appropriate recommended management strategies may be followed to control them. After a heavy rainfall, Parawilt symptoms may notice in some areas which can be cured by spraying Cobalt chloride @ 10mg/litre (10ppm) on affected plant within few hours of onset of symptoms. At Faridkot, the crop is 120 days old at Reproductive stage (Crop bearing squares, flowers, bolls). Weather remained mildly hot and humid during the reporting period. Give sprays of Potassium nitrate @2kg/acre at weekly intervals. The farmers who have witnessed leaf reddening in their cotton fields are advised to spray MgSO<sub>4</sub> @1%(1 kg/100 lts of water) twice during flowering and boll development stage as a prophylactic remedial measure. Whitefly has become serious over the past few weeks. If necessary, spray Triazophos 40 EC @ 600ml/acre or Ethion 50 EC @ 800 ml/acre if whitefly population is higher than ETL level of 6/leaf after rainfall. Do not spray formulations or mixtures containing pyrethroids. Cotton leaf curl disease intensity is on the increase. Bacterial leaf blight has appeared owing to cloudy weather; Spray Blitox@500g+Streptocycline @3g /acre if the disease spreads. If plants show Parawilt symptoms after rainfall, spray Cobalt Chloride @ 10 mg/litre on affected plants within few hours to check it.

**Haryana:** The crop is normal at Vegetative/reproductive stage. Interculture if required and weeds must be removed around the fields, water channel and road side to check the whitefly population on alternate host of CLCuD. Irrigate the field as and when required. Mean population of whitefly adults was above ET in fields. Population of leafhopper was below ETL. Pink boll worm infestation was observed in *Desi* cotton/non Bt. varieties. Solenopsis mealybug (*Phenacoccus solenopsis*) incidence was in traces in few fields. Incidence of *Spodeptera* was observed in few farmers field. Low to moderate incidence of leaf-curl virus disease was observed. Myrothecium leaf spot disease was observed in traces. For the control of bollworms in *Desi* and non Bt cotton varieties, recommended insecticides may be sprayed. Population of whitefly may increase fast in case the dry spell prolongs. Since *Aenasius* parasitoid is quite active, the mealybug is likely to remain in low profile and there is no need of spraying any insecticide against this pest. For control of root rot disease in cotton, soil drenching with 0.2% carbendazim solution may done in root rot affected areas. If foliar diseases appears spray Streptomycin sulphate (6-8g) plus Copper oxychloride (600-800 g) in 200litres of water per acre 3-4 times at 15 days interval

## CENTRAL INDIA

**Maharashtra:** At Akola, in pre monsoon sown cotton area, the symptoms of parawilt was noticed in farmers field. Drench 25g COC and 100g Urea and SOP each in 10 litres of water and spray the same within a week. Sucking pests like jassid/ aphid to be controlled with recommended pesticides. Weeding is necessary up to 60 days after sowing. Spray of Glyphosate in cotton should be avoided during windy situation and not to use power spray for spraying-

**Odisha:** The crop is in square initiation and flowering stage (52 to 72 days). The weather is hot and humid. Incidence of aphids and jassids observed (below ETL). When sucking pest population exceeds ETL (> 20% infested plants) spray Buprofezin @1 ml/litre of water.

## SOUTH INDIA

**Andhra Pradesh:** Gap filling and thinning is being carried out in the late sown crop. Weed management by working Gorru & Guntaka as an inter cultivation operation was carried out. Post emergence application of weedicides is recommended at 4 to 6 leaf stage of weeds. For the control of grassy weeds, Quizalofopethyl @ 400 ml/acre and for the control of broad leaf weeds, Pyriithiobac sodium @ 250 ml/acre is recommended. Advised prophylactic spray for the control of leaf spots. Summer cotton sown in approximately in an acreage of 25 thousand hectares which is in squaring to flowering stage to boll development stages. Suitable plant protection against sucking pests (Jassids & Thrips) were recommended. In Coastal A.P., the cotton crop is in seedling to 75 days stage and with the availability of soil moisture depending on the crop stage, top dressing of 25 to 30 kg Urea and 15kg of MOP is recommended. In Telangana, the cotton sowings were completed and the crop is in 25 days (seedling stage) to 75 days (squaring and initiation of flowering). Top dressing of nitrogen and potassium fertilizers is under progress. Inter-cultivation by working in the Gorru & Guntaka is under progress for the conservation of moisture and control of weeds.

**Karnataka:** The crop duration varies from 60 days to 80 days at square formation stage. Weeding and spraying operations were undertaken. Suggested to drain out the excess water stagnated in the crop due to high intensity rainfall. Top dress the crop with Urea (25 kg/ac) for immediate recovery from water stagnation effect.- In 80-90 days old cotton crop, it is suggested to spray the crop with Mancozeb @ 2g/lit of water for controlling leaf spot disease. Avoid heavy irrigation in black soils where the crop is at peak square and boll formation stage. Irrigating in alternate furrows is suggested to save water, time, labour and to irrigate more area in available water

### Weekly Advisory Report Coordinating Team

Scientists	Address
Dr K R Kranthi	Director, CICR, Nagpur
Dr A H Prakash	PC and Head, CICR, Regional station, Coimbatore
Dr. D Monga	Head, CICR, Regional station, Sirsa
Dr. S. B. Singh	Head, Div of Crop Improvement, CICR, Nagpur
Dr Sandhya Kranthi	Head, Div of Crop Protection, CICR, Nagpur
Dr Blasé De souza	Head, Div of Crop Production, CICR, Nagpur
Dr. Isabella Agarwal	Sr. Scientist CICR, Coimbatore
Sh. M.Sabesh	Scientist, CICR, Coimbatore

Scientists In-charge for Weather Report (AICCIP Centres)			
Scientists	Address	Mobile No	E Mail ID
Dr. Paramajit Singh	Punjab Agricultural University, Bathinda, Punjab	9463628801	rsmeenars@gmail.com
Dr. Pankaj Rathore	Punjab Agricultural University, Faridkot, Punjab	9464051995	pankaj@pau.edu
Dr. Jagdish Beniwal	CCS-Haryana Agricultural University, Hisar 125 004, Haryana	9416325420	cotton@hau.ernet.in
Dr.S.L.Ahuja	CCS-Haryana Agricultural University, Sirsa, Haryana	9255947380	slahuja2002@yahoo.com
Dr.K.N.Bhatia	Swami Keshwanand Rajasthan Agricultural University, Sriganganagar, Rajasthan	9352700411	bsmeena1969@rediffmail.com
Dr.Harphool Meena	Maharana Pratap University of Agri. & Technology, Udaipur – 313 001, Rajasthan	9460246043	hpagron@rediffmail.com
Dr. Narendra Kumar	CSA University of Agri. & Technology, Kanpur – 208 002, Uttar Pradesh	9335699132	jaqdishk64@yahoo.com
Dr. Gofaldu	Navsari Agricultural University, Navsari – 396 450, Gujarat	9662532645	girishfaldy@rediffmail.com
Dr.M.D.Khanpara	Junagadh Agricultural University, Junagadh – 362 001, Gujarat	9426990070	cotton@jau.in
Dr.R.W.Bharud	Mahatma Phule Krishi Vidyapeeth, Rahuri – 413 722, Maharashtra	9850244087	cotton_mpkv@rediffmail.com
Dr. B . R. Patil	Panjabrao Deshmukh Krishi Vidyapeeth, Akola – 444 104, Maharashtra	9657725801	srs cottonpdkv1@yahoo.co.in
Dr.P.R.Zanwar	Marathwada Agricultural University, Parbhani – 431 402, Maharashtra	7588151244	crsned@indiatimes.com
Dr. Satish Parsai	RVS Krishi Vishwa Vidhyalaya, Gwalior – 474 002, Madhya Pradesh	9406677601	aiccipkhandwa@gmail.com
Dr. B.S.Nayak	Orissa University of Agriculture & Technology, Bhubaneshwar – 751 003, Orissa	9437321675	bsnayak2007@rediffmail.com
Dr.S.Bharathi	Acharya N. G. Ranga Agricultural University, LAM, Guntur, AP	949072341	bharathi_says@yahoo.com
Dr. Sharma	Acharya N. G. Ranga Agricultural University, Nandyal, AP	08514-242296	sharmarars@gmail.com
Dr.Aladakatti	University of Agricultural Sciences, Dharwad – 580 005, Karnataka	9448861040	yaladakatti@rediffmail.com
Dr. Bheemana	University of Agricultural Sciences Raichur – 584 102, Karnataka	9448633232	bheemuent@rediffmail.com
Dr. Amala Balu	Tamil Nadu Agricultural University, Srivilliputhur, Tamil Nadu		
Dr. M Gunasekaran	Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu	9443631359	gunasekaran.pbg@gmail.com

--- end of the report ---