

Central Institute for Cotton Research, Nagpur

Fourth Weekly Advisory for Cotton Cultivation 16th to 22nd June 2014

"The advisory is based on inputs received from the State Agricultural Universities of the respective states"

Weed management: Wherever weeds have emerged, weedicides would provide effective and timely control. Weedicides are effective against younger (less than 10-15 days old) weeds, especially grasses. For grassy weeds, Quizalofop ethyl, Fenoxaprop ethyl, fluazifop butyl, can be used. For sedges and grasses, Propanil is effective and Pyriproxyfen sodium is effective on broad leaf weeds. Farmers may consult the technical experts of the Agricultural Universities for further details.

Water logging: Cotton is very sensitive to excess water. In many parts of Central and South India, water logging can be problematic due to excess rains. Cotton grown on deep black soils and ill drained conditions is worst affected due to water logging. Provide adequate drainage channels or water ways (particularly in heavy soils) along the slope of the land for draining excess water under heavy rainfall situations. For better soil moisture conservation, preferably in areas where rainfall is 700-900mm, the land can be reshaped into ridges and furrows with the help of a ridge plough or a bund former. This technique and sowing cotton on ridges would conserve rainwater and the furrows acts drainage channels whenever heavy rains are received particularly in heavy clays.

Drainage channels must be opened up along the field borders so that excess water is removed from the fields. If sowing hasn't yet been completed, it is strongly recommended that to take up sowing immediately on ridges and furrows by planting on top of ridges. Heavy rains will not affect the crop because the furrows will drain away excess water. Apply fertilizers if the crop becomes pale due to water logging. If heavy rains are forecast, fertilizer application may be postponed so as to prevent losses due to surface run-off.

Foliar spray with 0.5 to 1.0% DAP or 19:19:19 (soluble complex of Nitrogen) at weekly intervals will help the plants to recover from the effect of water logging.

STRATEGIES FOR MANAGEMENT OF PESTS, DISEASES & WEEDS

INSECT PEST MANAGEMENT

General recommendations

DOs

1. Select sucking pest resistant varieties/hybrids. Sucking pest resistant Bt hybrids may require very few insecticide interventions.
2. Inter-crop with cowpea or sorghum or soybean or blackgram to encourage predators of sucking pests.
3. Seed treatment with Imidacloprid @7gms/Kg of seed.
4. Use nitrogenous fertilizers to the minimum especially for sucking pest susceptible varieties.
5. Maintain field sanitation (weed free) and remove and destroy mealy bug infested plants &.
6. **Stem application or soil application** (near the root zone) of Imidacloprid, Dimethoate or Acephate at 30-40 DAS and 50-60 DAS for effective eco-friendly control of thrips, mirid bugs, mealy bugs and other sucking pests.

DON'Ts

7. **If possible avoid chemical insecticides during the first two months of the crop** to conserve naturally occurring biological control. Ladybird grubs and beetles, *Chrysoperla* grubs and adults, Syrphid flies, *Geocoris* grubs and bugs, *Aenasius* spp., *Aphelinus* grubs and wasps, mirid bugs and Spiders are the most important naturally occurring predators and parasitoids that effectively control aphids, jassids, thrips, mirids, whiteflies and mealybugs.

8. **Do not spray against minor lepidopteran insects** such as the cotton leaf folder, *Sylepta derogata* and cotton semilooper, *Anomis flava*. The larvae cause negligible damage to cotton but serve as hosts for parasitoids such as *Trichogramma* spp., *Apanteles* spp and *Sysiropa formosa*, that attack *H. armigera* and other bollworms.
9. **Do not spray Bt-formulations on Bt cotton** to avoid further selection pressure.
10. **Avoid foliar application of neonicotinoid insecticides** such as Acetamiprid, Imidacloprid, Clothianidin and Thiomethoxam which are likely to aggravate insect resistance, since hybrid cotton seeds are treated with imidacloprid.
11. **Do not use WHO Class-I (Extremely Hazardous category) insecticides** such as Phosphamidon, Methyl parathion, Phorate, Monocrotophos, Dichlorvos, Carbofuran, Methomyl, Triazophos and Metasystox.

SUCKING PEST MANAGEMENT

Economic Threshold Level (ETL): If whitefly and/or leafhopper damage reaches economic threshold levels of grade-II damage of curling and crinkling of lower leaves and yellowing of margins in 25% plants or more, any one of the following pest control measures as suggested below can be used.

- a. Neem oil 1.0% + Neem Seed Kernel Extract 5.0% + 0.05-0.1% detergent
- b. *Verticillium lecanii* 10gms/lit of water, wherever good formulations are available from reliable manufacturers
- c. Diafenthiuron (50WP 800g /ha),
- d. Flonicamid 50 WG 200g a.i/ha or
- e. Buprofezin 25% SC 200 g a.i/ha.

Insecticides such as Fipronil or Dimethoate or Acephate or Ethion can also be used but may be considered as alternatives only, in view of factors that relate to ecological and environmental safety, efficacy and resistance.

If mirid bugs are observed to cause economic damage to squares, it is advised to spray Acephate 75 SP @ 1 g/lit or Fipronil 5 SC @ 1.0 ml/lit of water

BOLLWORM MANAGEMENT

Bt cotton is effective in controlling bollworms.

The following strategies are being recommended for non-Bt cotton

At Economic Threshold Levels (ETLs) of 50% infested plants (plants having flared squares with entry hole) for *Helicoverpa armigera*.

1. **Use HaNPV on Bt-cotton** followed by the application of **5% NSKE** a week later. **Or, use Phosalone** at 50% bollworm infested plants (plants having flared squares with entry hole) or for the management of *Spodoptera* or whitefly.
2. ***Trichogramma***, if available, can be used on non-Bt genotypes at 70-80 DAS. Avoid *Trichogramma* egg parasitoid releases on Bt-cotton since maximum neonates get killed on Bt-cotton and with *Trichogramma* application becoming superfluous.
3. **Insecticides effective on Bollworms**, especially *Helicoverpa armigera*.
 - a. Chlorantraniliprole (Coragen),
 - b. Flubendiamide (Fame),
 - c. Spinosad,
 - d. Emamectin benzoate and
 - e. Indoxacarb

These insecticides have a high selective toxicity towards the target pests while being less toxic to many beneficial insects in the cotton ecosystem. These insecticides are ideally suited in eco-sustainable insecticide resistance management programmes.

4. **Pink bollworm and Spotted bollworms:** ETL level of one live larva in 10 green bolls or 8 moths per night for three consecutive nights. Spray Quinalphos 25 EC Profenophos 50 EC @ 2 ml/lit of water / Spray of Thiodicarb 75 WP @ 20 g or any pyrethroid.
5. ***Spodoptera litura*:** Collection of egg masses or application of *SNPV* (*Spodoptera litura* Nuclear Polyhedrosis Virus) @ 500 LE/ha or Spray 200 ml Rimon 10 EC or 250g Larvin 75WP in 250 litres of water per acre
6. To minimize **shoot weevil** damage, spray Profenofos @ 2 ml/lit
7. In case of snail incidence in heavy rainfall areas, baiting with 2% Metaldehyde (Snail kill) @ 12.5 kg/ha has to be taken up and it is to be applied at the hideouts of the snails, on the bunds and to the soil around the crop where the damage is seen

DISEASE MANAGEMENT

Parawilt or Sudden drying (New wilt) or Wilt / Root rot: Symptoms are noticed in some fields after drought followed by rains or irrigation.

Spray cobalt chloride @10mg/litre (10ppm) on affected plants within few hours of onset of symptoms and/or Drench plants with a mixture of Copper-Oxy-Chloride 25g and 200g Urea in 10 ltr of water or Carbendazim 1g/L.

Boll Rot: Generally early formed lower bolls rot due to cloudy and drizzling conditions.

Spray Mancozeb 75 WP + Chlorothalonil 70 WP each @ 2 g/lit of water. For better results, mix 10g Selvet 99 or 50 ml Triton in 100 litres of fungicide solution.

***Alternaria* blight:** spray Mancozeb@2.5 g per one litre of water.

Myrothecium leaf spot disease and/or Bacterial blight: Spray Streptomycin sulphate (15-20 g/ha) plus Copper oxychloride (1500-2000 g/ha) in 200-250 L of water.

WEED MANAGEMENT

Herbicides are most effective on younger weeds.

Post emergence herbicides (application rate 50 to 75 g ai /ha)

Grasses: Spray Quizalofop-ethyl or Fenoxaprop ethyl or Fluazifop butyl,

Sedges and grasses: Spray Propaquizafop ethyl

Broadleaf weeds: Spray Pyriithiobac sodium

GENERAL CROP HEALTH MANAGEMENT

1. **Optimize nutrient management** for macro and micronutrients. Foliar spray of MgSO₄, 2% Urea followed by 2% DAP, to ensure proper Cry1Ac expression and also to reduce problems of leaf reddening. Sprays of 1% cobalt chloride and soil drenching with Bavistin 1 % in the initial stage of wilt was found to help in the recovery of plants.
2. **Prevention of Leaf Reddening:** Spray 2 % urea, 0.5% Zinc Sulphate and 0.2 % Boron, twice at 15 days interval on 90 days old crop.
3. **Retention of squares and flowers:** Spray Planofix 4.5 SL (NAA) hormone @ 21 ppm (7 ml per 15 litres of water).

COTTON CROP SITUATION

Based on inputs received from the State Agricultural Universities of the respective States

CENTRAL INDIA

Maharashtra: Pre monsoon showers started in some pockets of Vidarbha. Farmers should take up harrowing and cleaning the field. Purchase of seed and fertilizers should be done well in advance. Dry seeding of straight varieties could be taken up during 16th to 20th June 2014 but for Bt cotton, sowing should be done only after the receipt of 75 to 100 mm rainfall. Those who have sown during pre monsoon viz last week of May 2014, hoeing

or weeding is needed. If showers are there, weed growth will suppress the height of cotton. Looking into monsoon behavior, intercropping of mung, urd, bean and short duration soybean is recommended in rainfed situation to avoid the risk under Bt and non Bt cotton in 1:1 or 2:1 ratio. Sow the cotton on ridges and furrow or broad bed furrow for better conservation of moisture.

Odisha: Sowing will be started after the onset of monsoon. Farmers are advised to procure seeds from reliable sources and keep all the inputs like fertilizers, and herbicides ready well in advance. Fertile and well drained soils having depth not less than 3 feet should be selected for hybrid cotton. Soils with low fertility and depth should be taken for high density planting (HDP) with suitable varieties. Well decomposed FYM/compost should be applied @ 2.5-5.0 ton/ha before the final ploughing. Fertilisers should be applied as per soil test reports. For hybrids 120:60:60 and for varieties 90:45:45 kg N:P₂O₅: K₂O/ha is recommended. Entire P₂O₅, 50% K₂O and 25% N should be applied as basal dose at the time of sowing. Seeds should be treated with Azotobactor and PSB @ 25 g/kg seeds on the date of sowing. For normal planting, a spacing of 90 cm x 60 cm and for HDP 60 cm x 10 cm should be followed. For control of weeds, Pendimethalin should be applied as pre emergence spray @ 1.0 kg/ha at 1 day after sowing when sufficient moisture is there in the soil.

SOUTH INDIA

Andhra Pradesh: Dry weather prevailed during the period under report with a mean RH (Max.) 61.3% & RH (Min.) 42.3% with bright sunshine hours. Summer preparatory cultivation is under progress. Spodoptera litura trap catches were observed @ 2-3 moths/trap/week. Summer cotton sown in approximately in an acreage of 25 thousand hectares which is in squaring to flowering stage to boll development stages. Suitable Plant Protection against sucking pests (Jassids & Thrips) were recommended.

Karnataka: It is advised to maintain required plant population by filling the gaps with Bt cotton seeds only in sown crop of Bt cotton. Sowing of Bt cotton can be continued till July first week in all districts of cotton growing region of Karnataka. Intra-specific Bt hybrid cotton (H x H) to be sown at a spacing of 90 cm x 60 cm and inter-specific Bt hybrid cotton (H x B) to be sown at 120 cm x 60 cm. Refugia crop seeds given with Bt cotton seeds should be sown around the Bt cotton crop. Bhendi crop has to be sown in Bt cotton plot at 10:1 row proportion as "Trap Crop" to minimise the shoot weevil attack. One or two rows of Maize crop may also be grown around the Bt cotton plot as "Feast Crop" to increase the population of predators that control sucking pest on Bt cotton. Spray Pendimethalin 30 EC weedicide (4 ml/lit of water) on the soil immediately after cotton sowing within 48 hrs to control the weeds up to 30 days. Short duration crops like green gram (1:1), peas (1:1), beans (1:1) and coriander (1:2) may be taken up as an intercrop with Bt cotton. Fertiliser dose of 100:50:50 NPK kg/ha of which 25% N & K and entire P fertilizers has to be applied basally at the time of sowing. Apply the remaining N and K as top dressing in 3 equal splits i.e. 25% N & K each at 30, 45 and 60 DAS.

Weekly Advisory Report Coordinating Team

Scientists	Address	Mobile No	E Mail ID
Dr K R Kranthi	Director, CICR, Nagpur		
Dr A H Prakash	PC and Head, CICR, Regional station, Coimbatore		
Dr. D Monga	Head, CICR, Regional station, Sirsa		
Dr. S. B. Singh	Head, Div of Crop Improvement, CICR, Nagpur		
Dr Sandhya Kranthi	Head, Div of Crop Protection, CICR, Nagpur		
Dr Blasé De souza	Head, Div of Crop Production, CICR, Nagpur		
Dr. Isabell Agrawal	Sr. Scientist CICR, Coimbatore		
Sh. M.Sabesh	Scientist, CICR, Coimbatore		
Scientists In-charge for Weather Report (AICCIP Centres)			
Scientists	Address	Mobile No	E Mail ID
Dr. Paramajit Singh	Punjab Agricultural University, Bathinda, Punjab	9463628801	rsmeenars@gmail.com

Dr. Pankaj Rathore	Punjab Agricultural University, Faridkot, Punjab	9464051995	pankaj@pau.edu
Dr. Jagdish Beniwal	CCS-Haryana Agricultural University, Hisar 125 004, Haryana	9416325420	cotton@hau.ernet.in
Dr.S.L.Ahuja	CCS-Haryana Agricultural University, Sirsa, Haryana	9255947380	slahuja2002@yahoo.com
Dr.K.N.Bhatia	Swami Keshwanand Rajasthan Agricultural University, Sriganganagar, Rajasthan	9352700411	bsmeena1969@rediffmail.com
Dr.Harphool Meena	Maharana Pratap University of Agri. & Technology, Udaipur – 313 001, Rajasthan	9460246043	hpagron@rediffmail.com
Dr. Narendra Kumar	CSA University of Agri. & Technology, Kanpur – 208 002, Uttar Pradesh	9335699132	jagdishk64@yahoo.com
Dr. Gofaldu	Navsari Agricultural University, Navsari – 396 450, Gujarat	9662532645	girishfaldu@rediffmail.com
Dr.M.D.Khanpara	Junagadh Agricultural University, Junagadh – 362 001, Gujarat	9426990070	cotton@jau.in
Dr.R.W.Bharud	Mahatma Phule Krishi Vidyapeeth, Rahuri – 413 722, Maharashtra	9850244087	cotton_mpkv@rediffmail.com
Dr. B . R. Patil	Panjabrao Deshmukh Krishi Vidyapeeth, Akola – 444 104, Maharashtra	9657725801	srscottonpdkv1@yahoo.co.in
Dr.P.R.Zanwar	Marathwada Agricultural University, Parbhani – 431 402, Maharashtra	7588151244	crsned@indiatimes.com
Dr. Satish Parsai	RVS Krishi Vishwa Vidhyalaya, Gwalior – 474 002, Madhya Pradesh	9406677601	aiccpkhandwa@gmail.com
Dr. B.S.Nayak	Orissa University of Agriculture & Technology, Bhubaneshwar – 751 003, Orissa	9437321675	bsnayak2007@rediffmail.com
Dr.S.Bharathi	Acharya N. G. Ranga Agricultural University, LAM, Guntur, AP	949072341	bharathi_says@yahoo.com
Dr. Sharma	Acharya N. G. Ranga Agricultural University, Nandyal, AP	08514-242296	sharmarars@gmail.com
Dr.Aladakatti	University of Agricultural Sciences, Dharwad – 580 005, Karnataka	9448861040	yaladakatti@rediffmail.com
Dr. Bheemana	University of Agricultural Sciences Raichur – 584 102, Karnataka	9448633232	bheemuent@rediffmail.com
Dr. Amala Balu	Tamil Nadu Agricultural University, Srivilliputhur, Tamil Nadu		
Dr. M Gunasekaran	Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu	9443631359	gunasekaran.pbg@gmail.com

--- end of the report ---