

Central Institute for Cotton Research, Nagpur

Seventeenth Weekly Advisory for Cotton Cultivation 15th to 21st September 2014

"The advisory is based on inputs received from the State Agricultural Universities of the respective states"

Weed management: Wherever weeds have emerged, weedicides would provide effective and timely control. Weedicides are effective against younger (less than 10-15 days old) weeds, especially grasses. For grassy weeds, Quizalofop ethyl, Fenoxaprop ethyl, fluazifop butyl, can be used. For sedges and grasses, Propaquizafop ethyl is effective and Pyriithiobac sodium is effective on broad leaf weeds. Farmers may consult the technical experts of the Agricultural Universities for further details.

Water logging: Cotton is very sensitive to excess water. In many parts of Central and South India, water logging can be problematic due to excess rains. Cotton grown on deep black soils and ill drained conditions is worst affected due to water logging. Provide adequate drainage channels or water ways (particularly in heavy soils) along the slope of the land for draining excess water under heavy rainfall situations. For better soil moisture conservation, preferably in areas where rainfall is 700-900mm, the land can be reshaped into ridges and furrows with the help of a ridge plough or a bund former. This technique and sowing cotton on ridges would conserve rainwater and the furrows acts drainage channels whenever heavy rains are received particularly in heavy clays.

Drainage channels must be opened up along the field borders so that excess water is removed from the fields. If sowing hasn't yet been completed, it is strongly recommended that to take up sowing immediately on ridges and furrows by planting on top of ridges. Heavy rains will not affect the crop because the furrows will drain away excess water. Apply fertilizers if the crop becomes pale due to water logging. If heavy rains are forecast, fertilizer application may be postponed so as to prevent losses due to surface run-off.

Foliar spray with 0.5 to 1.0% DAP at weekly intervals will help the plants to recover from the effect of water logging.

Cotton Sown Area (As on 11-09-2014)

S.No.	States	Normal Area (DES)*	Area in lakh ha		
			Bt	Non Bt	Total
1.	Andhra Pradesh + Telengana	19.83	21.78	1.669	23.45
	Andhra Pradesh (23.95%)	4.749	6.861	0.239	7.1
	Telangana (76.05%)	15.081	14.92	1.430	16.350
2.	Gujarat	26.490	27.13	2.930	30.060
3.	Haryana	5.640	6.310	0.080	6.390
4.	Karnataka	5.270	6.97	0.630	7.600
5.	Madhya Pradesh	6.390	5.503	0.285	5.788
6.	Maharashtra	39.160	39.888	1.822	41.710
7.	Odisha	0.970	0.000	1.250	1.250
8.	Punjab	5.170	4.300	0.200	4.500
9.	Rajasthan	4.000	3.954	0.208	4.162
10.	Tamil Nadu	1.250	0.056	0.014	0.070
11.	Uttar Pradesh	0.010	0.000	0.260	0.260
12.	Others	0.350		0.050	0.050
	All INDIA	114.530	115.892	9.398	125.291

* Directorate of Economics & Statistics, DAC, Ministry of Agriculture, Krishi Bhavan, New Delhi

Source: Director, DOCD, Mumbai

Weather forecast for 21st to 26th 'Sep '2014

Zones/ Weather parameter	Temperature (Min, Max)						Rainfall					
	States	21/09	22/09	23/09	24/09	25/09	26/09	21/09	22/09	23/09	24/09	25/09
Punjab	25,34	25,35	26,35	26,35	26,35	26,35	Mainly clear sky					
Haryana	25,36	25,36	25,36	25,36	25,36	25,36	Mainly clear sky					
Rajasthan	24,38	24,37	24,37	24,37	24,37	24,37	Mainly clear sky					
Gujarat	26,33	26,33	26,33	26,34	26,34	26,34	Mainly clear sky					
Maharashtra	24,33	24,32	24,34	24,34	24,33	24,34	Partly cloudy sky with thunder development	Partly cloudy sky				
M.P.	21,31	22,31	22,32	21,32	21,32	21,31	Partly cloudy sky					
Odisha	18,26	18,26	18,27	19,28	19,28	19,28	Generally cloudy sky with possibility of rain or thunderstorm					
A.P.	23,32	24,33	23,33	23,33	23,33	22,33	Mainly or Generally cloudy sky with possibility of rain or thunderstorm					
Karnataka	21,31	22,32	22,32	22,33	20,33	21,33	Moderate rain					
Tamil Nadu	22,33	22,33	23,34	23,34	23,34	23,34	Partly cloudy sky		Light rain			

Source: www.imd.gov.in

STRATEGIES FOR MANAGEMENT OF PESTS, DISEASES & WEEDS

INSECT PEST MANAGEMENT

General recommendations

DOs

1. Select sucking pest resistant varieties/hybrids. Sucking pest resistant Bt hybrids may require very few insecticide interventions.
2. Inter-crop with cowpea or sorghum or soybean or blackgram to encourage predators of sucking pests.
3. Seed treatment with Imidacloprid @7gms/Kg of seed.
4. Use nitrogenous fertilizers to the minimum especially for sucking pest susceptible varieties.
5. Maintain field sanitation (weed free) and remove and destroy mealy bug infested plants &.
6. **Stem application or soil application** (near the root zone) of Imidacloprid, Dimethoate or Acephate at 30-40 DAS and 50-60 DAS for effective eco-friendly control of thrips, mirid bugs, mealy bugs and other sucking pests.

DON'Ts

7. **If possible avoid chemical insecticides during the first two months of the crop** to conserve naturally occurring biological control. Ladybird grubs and beetles, *Chrysoperla* grubs and adults, Syrphid flies, *Geocoris* grubs and bugs, *Aenasius* spp., *Aphelinus* grubs and wasps, mirid bugs and Spiders are the most important naturally occurring predators and parasitoids that effectively control aphids, jassids, thrips, mirids, whiteflies and mealybugs.
8. **Do not spray against minor lepidopteran insects** such as the cotton leaf folder, *Sylepta derogata* and cotton semilooper, *Anomis flava*. The larvae cause negligible damage to cotton but serve as hosts for parasitoids such as *Trichogramma* spp., *Apanteles* spp and *Sysiropa formosa*, that attack *H. armigera* and other bollworms.
9. **Do not spray Bt-formulations on Bt cotton** to avoid further selection pressure.

10. **Avoid foliar application of neonicotinoid insecticides** such as Acetamiprid, Imidacloprid, Clothianidin and Thiomethoxam which are likely to aggravate insect resistance, since hybrid cotton seeds are treated with imidacloprid.
11. **Do not use WHO Class-I (Extremely Hazardous category) insecticides** such as Phosphamidon, Methyl parathion, Phorate, Monocrotophos, Dichlorvos, Carbofuran, Methomyl, Triazophos and Metasystox.

SUCKING PEST MANAGEMENT

Economic Threshold Level (ETL): If whitefly and/or leafhopper damage reaches economic threshold levels of grade-II damage of curling and crinkling of lower leaves and yellowing of margins in 25% plants or more, any one of the following pest control measures as suggested below can be used.

- a. Neem oil 1.0% + Neem Seed Kernel Extract 5.0% + 0.05-0.1% detergent
- b. *Verticillium lecanii* 10gms/lit of water, wherever good formulations are available from reliable manufacturers
- c. Diafenthiuron (50WP 800g /ha),
- d. Flonicamid 50 WG 200g a.i/ha or
- e. Buprofezin 25% SC 200 g a.i/ha.

Insecticides such as Fipronil or Dimethoate or Acephate or Ethion can also be used but may be considered as alternatives only, in view of factors that relate to ecological and environmental safety, efficacy and resistance.

If mirid bugs are observed to cause economic damage to squares, it is advised to spray Acephate 75 SP @ 1 g/lit or Fipronil 5 SC @ 1.0 ml/lit of water

BOLLWORM MANAGEMENT

Bt cotton is effective in controlling bollworms.

The following strategies are being recommended for non-Bt cotton

At Economic Threshold Levels (ETLs) of 50% infested plants (plants having flared squares with entry hole) for *Helicoverpa armigera*.

1. **Use HaNPV on Bt-cotton** followed by the application of **5% NSKE** a week later. **Or, use Phosalone** at 50% bollworm infested plants (plants having flared squares with entry hole) or for the management of *Spodoptera* or whitefly.
2. *Trichogramma*, if available, can be used on non-Bt genotypes at 70-80 DAS. Avoid *Trichogramma* egg parasitoid releases on Bt-cotton since maximum neonates get killed on Bt-cotton and with *Trichogramma* application becoming superfluous.
3. **Insecticides effective on Bollworms**, especially *Helicoverpa armigera*.
 - a. Chlorantraniliprole (Coragen),
 - b. Flubendiamide (Fame),
 - c. Spinosad,
 - d. Emamectin benzoate and
 - e. Indoxacarb

These insecticides have a high selective toxicity towards the target pests while being less toxic to many beneficial insects in the cotton ecosystem. These insecticides are ideally suited in eco-sustainable insecticide resistance management programmes.

4. **Pink bollworm and Spotted bollworms:** ETL level of one live larva in 10 green bolls or 8 moths per night for three consecutive nights. Spray Quinalphos 25 EC Profenophos 50 EC @ 2 ml/lit of water / Spray of Thiodicarb 75 WP @ 20 g or any pyrethroid.
5. ***Spodoptera litura*:** Collection of egg masses or application of SNPV (*Spodoptera litura* Nuclear Polyhedrosis Virus) @ 500 LE/ha or Spray 200 ml Rimon 10 EC or 250g Larvin 75WP in 250 litres of water per acre
6. To minimize **shoot weevil** damage, spray Profenofos @ 2 ml/lit
7. In case of snail incidence in heavy rainfall areas, baiting with 2% Metaldehyde (Snail kill) @ 12.5 kg/ha has to be taken up and it is to be applied at the hideouts of the snails, on the bunds and to the soil around the crop where the damage is seen

DISEASE MANAGEMENT

1. **Parawilt or Sudden drying (New wilt) or Wilt / Root rot:** Symptoms are noticed in some fields after drought followed by rains or irrigation. Spray cobalt chloride @10mg/litre (10ppm) on affected plants within few hours of onset of symptoms and/or Drench plants with a mixture of Copper-Oxy-Chloride 25g and 200g Urea in 10 ltr of water or Carbendazim 1g/L.
2. **Boll Rot:** Generally early formed lower bolls rot due to cloudy and drizzling conditions. Spray Mancozeb 75 WP + Chlorothalonil 70 WP each @ 2 g/lit of water. For better results, mix 10g Selvet 99 or 50 ml Triton in 100 litres of fungicide solution.
3. ***Alternaria* blight:** spray Mancozeb@2.5 g per one litre of water.
4. **Myrothecium leaf spot disease and/or Bacterial blight:** Spray Streptomycin sulphate (15-20 g/ha) plus Copper oxychloride (1500-2000 g/ha) in 200-250 L of water.

WEED MANAGEMENT

Herbicides are most effective on younger weeds.

Post emergence herbicides (application rate 50 to 75 g ai /ha)

1. **Grasses:** Spray Quizalofop-ethyl or Fenoxaprop ethyl or Fluazifop butyl,
2. **Sedges and grasses:** Spray Propaquizafop ethyl
3. **Broadleaf weeds:** Spray Pyrithiobac sodium

GENERAL CROP HEALTH MANAGEMENT

1. **Optimize nutrient management** for macro and micronutrients. Foliar spray of MgSO₄, 2% Urea followed by 2% DAP, to ensure proper Cry1Ac expression and also to reduce problems of leaf reddening. Sprays of 1% cobalt chloride and soil drenching with Bavistin 1 % in the initial stage of wilt was found to help in the recovery of plants.
2. **Prevention of Leaf Reddening:** Spray 2 % urea, 0.5% Zinc Sulphate and 0.2 % Boron, twice at 15 days interval on 90 days old crop.
3. **Retention of squares and flowers:** Spray Planofix 4.5 SL (NAA) hormone @ 21 ppm (7 ml per 15 litres of water).

DROUGHT MANAGEMENT**Odisha****Mid season drought (long dry spell)**

Condition			Suggested Contingency Measures	
Mid season drought (long dry spell)	Major Farming Situation	Normal Crop / Cropping System	Crop management	Soil nutrient & moisture conservation measures
At flowering/ fruiting stage	Red soil High rainfall Medium elevation	Cotton	Applying of Planofix hormone * spraying the crop with Imidacloprid for controlling of sucking pests	Apply 1250ml micronutrient/ha
	Red and Yellow soil High rainfall Medium elevation	Cotton		
	Black soil High rainfall Medium elevation	Cotton + Arhar		

Terminal drought (Early withdrawal of monsoon)

Condition			Suggested Contingency Measures	
Terminal drought (Early withdrawal of monsoon)	Major Farming Situation	Normal Crop / Cropping System	Crop management	Rabi Crop planning
	Red soil, High rainfall, Medium elevation	Cotton	Provide protective irrigation	Mulch with stovers Dibble rabi crop
		Cotton	Provide protective irrigation	
		Cotton + Arhar	Provide protective irrigation Harvest at physiological maturity stage	

IMPORTANT NOTE: (PEST MANAGEMENT)

Farmers are advised not to spray pyrethroids early in the season singly or in combination against sucking pests such as the whiteflies not only for cotton but also on other *H. armigera* host plants such as soybean, as it may exacerbate bollworm problems in non Bt cotton, wherever cultivated.

COTTON CROP SITUATION

(Based on inputs received from the State Agricultural Universities of the respective States)

NORTH INDIA

Punjab: At Faridkot, the crop is nearly 130 days at reproductive stage (Crop bearing squares, flowers, bolls), Weather remained mildly hot and humid during the reporting period. Farmers are advised to drain out the stagnant water if there is medium to heavy rainfall. Give sprays of Potassium nitrate @2kg/acre at weekly intervals. The farmers who have witnessed leaf reddening in their cotton fields are advised to spray MgSO₄ @1%(1 kg/100 lts of water) twice during flowering and boll development stage as a prophylactic remedial measure. If hot and humid conditions persist, Whitefly incidence will increase. Spray Triazophos 40 EC @ 600ml/acre or Ethion 50 EC @ 800 ml/acre if its population is higher than ETL level of 6/leaf after rainfall. If numbers are too high, spray Polo (Diafenthiuron 50 WP @200g/acre). Cotton leaf curl disease intensity has increased to highest levels in the past week; Fungal foliar spots have appeared and might increase if cloudy weather persists. If plants show Parawilt symptoms after rainfall, spray Cobalt Chloride @ 10 mg/litre on affected plants within few hours to check it. At Bhatinda, cotton crop is in boll development stage. Boll opening has started in early maturing varieties/hybrids. The infestation of whitefly is quite high in cotton growing areas of the State due to favourable hot and humid conditions. Farmers are advised to monitor their crop for insect pests and diseases regularly to control or to keep the incidence and infestation below economic threshold level (ETL).

Haryana: The crop is normal at vegetative/reproductive stage, Irrigate the field as and when required. Mean population of whitefly adults was above ET in fields. Population of leafhopper was below ET. Incidence of *Spodoptera* was observed in farmers' field. Low to moderate incidence of leaf-curl virus disease was noticed. Myrothecium leaf spot disease was observed in traces. Bacterial leaf blight did not appear in cotton field. Farmers should be advised to monitor disease and insect-pest populations on weekly basis and to apply control measures only at ET. For the control of *Spodoptera*, spray 200 ml Rimon 10 EC in 200 L of water per acre. For the control of bollworms in Desi and non Bt cotton varieties, spray pesticides as recommended. Population of whitefly may increase fast in case the dry spell prolongs. For checking whitefly, Nimbecidine 300 ppm @ 1 litre/acre may be sprayed twice at five days interval followed by one spray of 300 ml Dimethoate 30 EC in 200 litres of water per acre. If foliar diseases appear, spray Streptomycine sulphate (6-8g) plus Copper oxychloride (600-800g) in 200 L of water per acre 3-4 times at 15 days interval.

Rajasthan: The crop is 98 to 103 days old at initial boll bursting stage, The weather was clear sky with high humidity, Weeding and spray for sucking pests has been taken up, At present, the crop is weed free. Jassids and white fly infestation continued to remain near ETL. Bollworm infestation was not recorded. *Anthraxnose* and *alternaria leaf blight* infestation was recorded.

CENTRAL INDIA

Gujarat: The crop is nearly 150 days old at reproductive stage (Crop bearing squares, flowers, bolls). Weather remained cloudy and rainy during the reporting period. Since heavy rainfall is there across the cotton belt, farmers are advised to drain out the stagnant water. Due to heavy rainfall, incidence of weeds is intense. The population of jassids was recorded above ETL and thrips, aphids, white fly and mealy bug were low throughout the reporting period. No incidence of bacterial blight and other cotton diseases.

Maharashtra: At Rahuri, the crop is 90 to 130 days old at boll development stage, Fertilizer application, weeding, Insecticide sprays were taken up. Weed infestation was up to 30 to 40 per cent, sucking pest attack around 25 to 30 per cent. Among diseases, Alternaria leaf blight incidence to the tune of 20 per cent and Tobacco Streak Virus up to 10 per cent were noticed. The control measures recommended may be followed to control the pest and disease attack. Incidence of semilooper and its leaf damage recorded in non BT cotton that should be ignored. At Nanded, heavy infestation of thrips and jassids is observed on irrigated as well as rainfed crop. Farmers are advised to spray Buprofenzin or Fipronil @ 20 ml / 10 lit for knapsack spray. Foliar application

of 13:0:45 @ 1% (100 g / 10 lit) is should be given for irrigated cotton which is at boll development stage. For surface irrigated cotton, irrigation lay outs may be prepared. Foliar application of Urea / DAP @ 2% (200 g / 10 lit) is recommended for rainfed at square formation and flowering stage. Spray of Carbendazim @ 10 g / 10 lit may be taken as precautionary measure for fungal diseases.

Odisha: The crop is 66 to 86 days old at flowering and boll formation stage. The weather was hot and humid. Incidence of aphids, jassids and mealy bug observed, sporadic incidence leaf folder, stem borer and Spodoptera were recorded. All three types of weeds i.e grasses, sedges and broad leaved weeds have infested the crop. Farmers are advised to drain out excess water from the field during heavy rainfall. For control of weeds, glyphosate should be applied as post emergence directed spray @ 1.0 kg/ha 50 days after sowing. To reduce the flower and fruit drop, spray Planofix @ 7.0 ml/15 litres of water. For reducing the aphids, jassids and mealy bug population, spray neem based pesticides @ 2.5 ml/litre of water. When sucking pest population exceeds ETL (> 20% infested plants) spray Buprofezin @1 ml/litre of water. To control the pests like Spodoptera, stem borer and leaf folder, spray Beauveria bassiana bio-pesticide @ 2 ml per litre of water.

SOUTH INDIA

Andhra Pradesh: Inter-cultivation was taken up by working with Gorru and Guntaka. Application of fertilizers in the form of N & K as top dressing was done depending upon the age of the crop. Post emergence application of weedicides is recommended at 4 – 6 leaf stage of weeds wherever inter-cultivation is not possible due to high moisture stress. For the control of grassy weeds, Quizalofopethyl @ 400 ml/acre and for the control of broad leaf weeds, Pyriithiobac sodium @ 250 ml/acre is recommended. Imidacloprid and water @ 60 DAS if jassid incidence crosses ETL i.e. 2/leaf, spraying of Acephate 75 SP 1.5 g/l or fipronil 5% SC 2ml/l is also recommended. Cloudy weather coupled with high humidity and rains is congenial for the development of diseases. Spraying of copper oxy chloride @ 30 g + Streptocycline @ 1 g in 10 lt. of water as prophylactic measure. Soil drenching with Copper oxy chloride @ 3 g/l (or) Carbendazim @ 1 g/l near the root zone is advocated against wilt diseases and root rot after draining the water. Spraying of Mancozeb @ 3 g/l or Propiconazole @ 1 ml/l is recommended to manage the leaf spots. In Coastal A.P the cotton crop is in 25 to 85 days stage. The rains received during the period under report resulted in water logging and high moisture stress. Draining out the water from the fields is recommended. Inter-cultivation and split application of fertilizers to be taken up at the earliest possible. Foliar application of 2% urea/ 2% KNO₃/ 1-2% DAP/ 2% 19-19-19 is recommended. In Telangana, the cotton crop is 35 days old at (vegetative) to 90 days (squaring, flowering & boll development) stage.. Topdressing of N and K fertilizers is under progress.

Tamil Nadu: The crop is 45 to 50 days old at square formation stage. The crop is normal. No significant incidence of pests and diseases or weeds noticed in the fields.

Weekly Advisory Report Coordinating Team

Scientists	Address
Dr K R Kranthi	Director, CICR, Nagpur
Dr A H Prakash	PC and Head, CICR, Regional station, Coimbatore
Dr. D Monga	Head, CICR, Regional station, Sirsa
Dr. S. B. Singh	Head, Div of Crop Improvement, CICR, Nagpur
Dr Sandhya Kranthi	Head, Div of Crop Protection, CICR, Nagpur
Dr Blasé De souza	Head, Div of Crop Production, CICR, Nagpur
Dr. Isabella Agarwal	Sr. Scientist CICR, Coimbatore
Sh. M.Sabesh	Scientist, CICR, Coimbatore

Scientists In-charge for Weather Report (AICCIP Centres)			
Scientists	Address	Mobile No	E Mail ID
Dr. Paramajit Singh	Punjab Agricultural University, Bathinda, Punjab	9463628801	rsmeenars@gmail.com
Dr. Pankaj Rathore	Punjab Agricultural University, Faridkot, Punjab	9464051995	pankaj@pau.edu
Dr. Jagdish Beniwal	CCS-Haryana Agricultural University, Hisar 125 004, Haryana	9416325420	cotton@hau.ernet.in
Dr.S.L.Ahuja	CCS-Haryana Agricultural University, Sirsa, Haryana	9255947380	slahuja2002@yahoo.com
Dr.K.N.Bhatia	Swami Keshwanand Rajasthan Agricultural University, Sriganganagar, Rajasthan	9352700411	bsmeena1969@rediffmail.com
Dr.Harphool Meena	Maharana Pratap University of Agri. & Technology, Udaipur – 313 001, Rajasthan	9460246043	hpagron@rediffmail.com
Dr. Narendra Kumar	CSA University of Agri. & Technology, Kanpur – 208 002, Uttar Pradesh	9335699132	jaqdishk64@yahoo.com
Dr. Gofaldu	Navsari Agricultural University, Navsari – 396 450, Gujarat	9662532645	girishfaldu@rediffmail.com
Dr.M.D.Khanpara	Junagadh Agricultural University, Junagadh – 362 001, Gujarat	9426990070	cotton@jau.in
Dr.R.W.Bharud	Mahatma Phule Krishi Vidyapeeth, Rahuri – 413 722, Maharashtra	9850244087	cotton_mpkv@rediffmail.com
Dr. B . R. Patil	Panjabrao Deshmukh Krishi Vidyapeeth, Akola – 444 104, Maharashtra	9657725801	srs cottonpdkv1@yahoo.co.in
Dr.P.R.Zanwar	Marathwada Agricultural University, Parbhani – 431 402, Maharashtra	7588151244	crsned@indiatimes.com
Dr. Satish Parsai	RVS Krishi Vishwa Vidhyalaya, Gwalior – 474 002, Madhya Pradesh	9406677601	aiccipkhandwa@gmail.com
Dr. B.S.Nayak	Orissa University of Agriculture & Technology, Bhubaneshwar – 751 003, Orissa	9437321675	bsnayak2007@rediffmail.com
Dr.S.Bharathi	Acharya N. G. Ranga Agricultural University, LAM, Guntur, AP	949072341	bharathi_says@yahoo.com
Dr. Sharma	Acharya N. G. Ranga Agricultural University, Nandyal, AP	08514-242296	sharmarars@gmail.com
Dr.Aladakatti	University of Agricultural Sciences, Dharwad – 580 005, Karnataka	9448861040	yaladakatti@rediffmail.com
Dr. Bheemana	University of Agricultural Sciences Raichur – 584 102, Karnataka	9448633232	bheemuent@rediffmail.com
Dr. Amala Balu	Tamil Nadu Agricultural University, Srivilliputhur, Tamil Nadu		
Dr. M Gunasekaran	Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu	9443631359	gunasekaran.pbg@gmail.com

--- end of the report ---