

## Central Institute for Cotton Research, Nagpur

### Twenty-Second Weekly Advisory for Cotton Cultivation 27<sup>th</sup> October to 2<sup>nd</sup> November 2013

"The advisory is based on inputs received from the State Agricultural Universities of the respective states"

**Weed management:** Wherever weeds have emerged, weedicides would provide effective and timely control. Weedicides are effective against younger (less than 10-15 days old) weeds, especially grasses. For grassy weeds, Quizalofop ethyl, Fenoxaprop ethyl, fluazifop butyl, can be used. For sedges and grasses, Propanil is effective and Pyriproxyfen sodium is effective on broad leaf weeds. Farmers may consult the technical experts of the Agricultural Universities for further details.

**Water logging:** Cotton is very sensitive to excess water. In many parts of Central and South India, water logging can be problematic due to excess rains. Cotton grown on deep black soils and ill drained conditions is worst affected due to water logging. Provide adequate drainage channels or water ways (particularly in heavy soils) along the slope of the land for draining excess water under heavy rainfall situations. For better soil moisture conservation, preferably in areas where rainfall is 700-900mm, the land can be reshaped into ridges and furrows with the help of a ridge plough or a bund former. This technique and sowing cotton on ridges would conserve rainwater and the furrows acts drainage channels whenever heavy rains are received particularly in heavy clays.

Drainage channels must be opened up along the field borders so that excess water is removed from the fields. If sowing hasn't yet been completed, it is strongly recommended that to take up sowing immediately on ridges and furrows by planting on top of ridges. Heavy rains will not affect the crop because the furrows will drain away excess water. Apply fertilizers if the crop becomes pale due to water logging. If heavy rains are forecast, fertilizer application may be postponed so as to prevent losses due to surface run-off.

Foliar spray with 0.5 to 1.0% DAP or 19:19:19 (soluble complex of Nitrogen) at weekly intervals will help the plants to recover from the effect of water logging.

Net Cotton Area sown as on 28-10-2013

State	Lakh hectares
Punjab	5.05
Haryana	5.57
Rajasthan	3.03
Uttar Pradesh	0.23
Gujarat	26.91
Madhya Pradesh	6.21
Maharashtra	38.72
Odisha	1.34
Andhra Pradesh	21.42
Karnataka	5.38
Tamil Nadu	1.16
Others	0.10
TOTAL	115.12

Source: Director, DOCD, Mumbai

Weather forecast for 29<sup>th</sup> Oct to 1<sup>st</sup> Nov 2013

Zones/ Weather parameter	Temperature ( Min, Max)				Rainfall			
	29/10	30/10	31/10	01/11	29/10	30/10	31/10	01/11
Punjab	15,30	15,29	15,29	15,29	Mainly clear sky			
Haryana	16,32	16,32	15,32	15,31	Mainly clear sky			
Rajasthan	16,35	16,35	15,35	15,35	Clear sky			
U.P.	18,31	18,31	17,30	17,30	Partly cloudy sky	Mainly clear sky		
Gujarat	21,35	21,35	20,35	20,35	Mainly clear sky			
Maharashtra	19,31	19,31	20,32	20,32	Mainly clear sky	Partly cloudy sky		
Madhya Pradesh	15,30	15,31	14,31	14,30	Clear sky			
Odisha	20,30	20,30	20,30	20,30	Partly cloudy sky			
Andhra Pradesh	22,29	22,30	21,30	21,30	Partly cloudy sky with possibility of rain or thunderstorm			
Karnataka	21,29	21,30	21,30	20,30	Light rain	Partly cloudy sky with possibility of rain or thunderstorm		Light rain
Tamil Nadu	21,31	23,31	24,31	23,31	Partly cloudy sky			

Source: www.imd.gov.in

## STRATEGIES FOR MANAGEMENT OF PESTS, DISEASES &amp; WEEDS

## INSECT PEST MANAGEMENT

*General recommendations**DOs*

1. Select sucking pest resistant varieties/hybrids. Sucking pest resistant Bt hybrids may require very few insecticide interventions.
2. Inter-crop with cowpea or sorghum or soybean or blackgram to encourage predators of sucking pests.
3. Seed treatment with Imidacloprid @7gms/Kg of seed.
4. Use nitrogenous fertilizers to the minimum especially for sucking pest susceptible varieties.
5. Maintain field sanitation (weed free) and remove and destroy mealy bug infested plants &.
6. **Stem application or soil application** (near the root zone) of Imidacloprid, Dimethoate or Acephate at 30-40 DAS and 50-60 DAS for effective eco-friendly control of thrips, mirid bugs, mealy bugs and other sucking pests.

*DON'Ts*

7. **If possible avoid chemical insecticides during the first two months of the crop** to conserve naturally occurring biological control. Ladybird grubs and beetles, *Chrysoperla* grubs and adults, Syrphid flies, *Geocoris* grubs and bugs, *Aenasius* spp., *Aphelinus* grubs and wasps, mirid bugs and Spiders are the most important naturally occurring predators and parasitoids that effectively control aphids, jassids, thrips, mirids, whiteflies and mealybugs.
8. **Do not spray against minor lepidopteran insects** such as the cotton leaf folder, *Sylepta derogata* and cotton semilooper, *Anomis flava*. The larvae cause negligible damage to cotton but

serve as hosts for parasitoids such as *Trichogramma* spp., *Apanteles* spp and *Sysiropa formosa*, that attack *H. armigera* and other bollworms.

9. **Do not spray Bt-formulations on Bt cotton** to avoid further selection pressure.
10. **Avoid foliar application of neonicotinoid insecticides** such as Acetamiprid, Imidacloprid, Clothianidin and Thiomethoxam which are likely to aggravate insect resistance, since hybrid cotton seeds are treated with imidacloprid.
11. **Do not use WHO Class-I (Extremely Hazardous category) insecticides** such as Phosphamidon, Methyl parathion, Phorate, Monocrotophos, Dichlorvos, Carbofuran, Methomyl, Triazophos and Metasystox.

### SUCKING PEST MANAGEMENT

**Economic Threshold Level (ETL):** If whitefly and/or leafhopper damage reaches economic threshold levels of grade-II damage of curling and crinkling of lower leaves and yellowing of margins in 25% plants or more, any one of the following pest control measures as suggested below can be used.

- a. Neem oil 1.0% + Neem Seed Kernel Extract 5.0% + 0.05-0.1% detergent
- b. *Verticillium lecanii* 10gms/lit of water, wherever good formulations are available from reliable manufacturers
- c. Diafenthuron (50WP 800g /ha),
- d. Flonicamid 50 WG 200g a.i/ha or
- e. Buprofezin 25% SC 200 g a.i/ha.

Insecticides such as Fipronil or Dimethoate or Acephate or Ethion can also be used but may be considered as alternatives only, in view of factors that relate to ecological and environmental safety, efficacy and resistance.

If mirid bugs are observed to cause economic damage to squares, it is advised to spray Acephate 75 SP @ 1 g/lit or Fipronil 5 SC @ 1.0 ml/lit of water

### BOLLWORM MANAGEMENT

Bt cotton is effective in controlling bollworms.

*The following strategies are being recommended for non-Bt cotton*

At Economic Threshold Levels (ETLs) of 50% infested plants (plants having flared squares with entry hole) for *Helicoverpa armigera*.

1. **Use HaNPV on Bt-cotton** followed by the application of **5% NSKE** a week later. **Or, use Phosalone** at 50% bollworm infested plants (plants having flared squares with entry hole) or for the management of *Spodoptera* or whitefly.
2. ***Trichogramma***, if available, can be used on non-Bt genotypes at 70-80 DAS. Avoid *Trichogramma* egg parasitoid releases on Bt-cotton since maximum neonates get killed on Bt-cotton and with *Trichogramma* application becoming superfluous.
3. **Insecticides effective on Bollworms**, especially *Helicoverpa armigera*.
  - a. Chlorantraniliprole (Coragen),
  - b. Flubendiamide (Fame),

- c. Spinosad,
- d. Emamectin benzoate and
- e. Indoxacarb

These insecticides have a high selective toxicity towards the target pests while being less toxic to many beneficial insects in the cotton ecosystem. These insecticides are ideally suited in eco-sustainable insecticide resistance management programmes.

4. **Pink bollworm and Spotted bollworms:** ETL level of one live larva in 10 green bolls or 8 moths per night for three consecutive nights. Spray Quinalphos 25 EC Profenophos 50 EC @ 2 ml/lit of water / Spray of Thiodicarb 75 WP @ 20 g or any pyrethroid.
5. ***Spodoptera litura*:** Collection of egg masses or application of *SNPV* (*Spodoptera litura* Nuclear Polyhedrosis Virus) @ 500 LE/ha or Spray 200 ml Rimon 10 EC or 250g Larvin 75WP in 250 litres of water per acre
6. To minimize **shoot weevil** damage, spray Profenofos @ 2 ml/lit
7. In case of snail incidence in heavy rainfall areas, baiting with 2% Metaldehyde (Snail kill) @ 12.5 kg/ha has to be taken up and it is to be applied at the hideouts of the snails, on the bunds and to the soil around the crop where the damage is seen

### DISEASE MANAGEMENT

**Parawilt or Sudden drying (New wilt) or Wilt / Root rot:** Symptoms are noticed in some fields after drought followed by rains or irrigation.

Spray cobalt chloride @10mg/litre (10ppm) on affected plants within few hours of onset of symptoms and/or Drench plants with a mixture of Copper-Oxy-Chloride 25g and 200g Urea in 10 ltr of water or Carbendazim 1g/L.

**Boll Rot:** Generally early formed lower bolls rot due to cloudy and drizzling conditions.

Spray Mancozeb 75 WP + Chlorothalonil 70 WP each @ 2 g/lit of water. For better results, mix 10g Selvet 99 or 50 ml Triton in 100 litres of fungicide solution.

***Alternaria* blight:** spray Mancozeb@2.5 g per one litre of water.

**Myrothecium leaf spot disease and/or Bacterial blight:** Spray Streptomycin sulphate (15-20 g/ha) plus Copper oxychloride (1500-2000 g/ha) in 200-250 L of water.

### WEED MANAGEMENT

Herbicides are most effective on younger weeds.

**Post emergence herbicides (application rate 50 to 75 g ai /ha)**

**Grasses:** Spray Quizalofop-ethyl or Fenoxaprop ethyl or Fluazifop butyl,

**Sedges and grasses:** Spray Propaquizafop ethyl

**Broadleaf weeds:** Spray Pyriithiobac sodium

### GENERAL CROP HEALTH MANAGEMENT

- **Optimize nutrient management** for macro and micronutrients. Foliar spray of MgSO<sub>4</sub>, 2% Urea followed by 2% DAP, to ensure proper Cry1Ac expression and also to reduce problems of leaf reddening. Sprays of 1% cobalt chloride and soil drenching with Bavistin 1 % in the initial stage of wilt was found to help in the recovery of plants.

- **Prevention of Leaf Reddening:** Spray 2 % urea, 0.5% Zinc Sulphate and 0.2 % Boron, twice at 15 days interval on 90 days old crop.
- **Retention of squares and flowers:** Spray Planofix 4.5 SL (NAA) hormone @ 21 ppm (7 ml per 15 litres of water).

## COTTON CROP SITUATION

*Based on inputs received from the State Agricultural Universities of the respective States*

### NORTH INDIA

The crop is in maturity phase and picking has been initiated. *Kapas* should be picked dry free from trash. It is advised to start picking in the morning after the dew dries from the opened bolls to avoid deterioration of seed cotton quality. Picking should be done after every 8 to 10 days to avoid losses. No agronomic or entomological interventions are needed henceforth. Suitable measures are to be taken for clean picking and storage to fetch better price of the harvested cotton in the market. Farmers are advised to harvest the crop when it has opened more than 50 per cent and there is no dew/ moisture. They are advised to cover their head with cloth. Picked cotton should be placed on cotton sheets or tarpaulins. Care should be taken to pick bolls without leaves, pedicel/ branches to have clean picked cotton.

**Haryana:** In Haryana during preceding week dry weather was observed. Cotton crop is in picking stage. In general, the crop is healthy. Do not irrigate the field after one third opening of bolls in the field. Avoid picking of rotten bolls. Dry the *kapas* before storage to avoid micro-organism damage. Farmers are advised to monitor their crop regularly.

**Uttar Pradesh:** Crop is in boll development and maturity stage. Farmers are advised to pick the bursting bolls and dry in the sun light and keep in jute bags.

### CENTRAL INDIA

**Maharashtra:** Cotton in Marathwada region is in boll development to boll bursting stage. Leaf reddening and Grey Mildew is recorded for which the recommended control measures are to be taken up. Square fall is noticed in some pockets. Spraying of NAA @ 1.2 ml per 10 lit water is advised.

**Odisha:** The cotton crop is at boll development to maturity stage (118-128 days). Postpone the picking of seed cotton during rainy days. Spray 2% DAP for better boll development. To check the incidence of sucking pests like aphids, jassids, thrips and whitefly spray recommended insecticides. Regular monitoring should be done for incidence of *Spodoptera* and leaf folders. In case of severe infestation of American boll worms ( at ETLs of 90-100% plants showing flared up squares) spray recommended control measures during 80 to 100 days after sowing.

### SOUTH INDIA

**Andhra Pradesh:** Cyclonic wide spread heavy rains received during past one week resulted in damage to the cotton crop in several districts. Wherever fields are inundated and water stagnation is observed, farmers are advised to drain-out the water from fields immediately. Application of booster dose of 30kg Urea and 10-15 kg MOP per acre is recommended. Foliar application of 2% Urea or 2% DAP or 1-2% KNO<sub>3</sub> along with 1% MgSO<sub>4</sub> is recommended to mitigate the stress conditions. For the control of sucking pests need based spraying of insecticides is recommended. Bollworms particularly *Spodoptera litura* is to be monitored through pheromone traps. Spraying of recommended doses is to be done if

Spodoptera incidence crosses ETL (one egg mass or one skeletonised leaf for 10 plants). Sudden death of the plants in patches or yellowing of leaves and wilting of plants should be managed by drenching the affected plants and soil with 0.3% (3 g/l) copper oxy chloride or 0.1% (1 g/l) Carbendazim or 0.1% (1 g/l) benomyl solution. Angular leaf spot and vein blight should be managed by spraying 0.3% (3 g/l) copper oxy chloride + 0.01% (100 mg/l) Streptocycline at 7 to 10 days interval. Fungal leaf spots should be managed by protective or curative spraying with appropriate fungicides at 7 to 10 days interval.

**Karnataka:** In 100 days old crop it is suggested to spray any recommended pyrethroid for controlling pink bollworm. Rotting of early formed lower bolls is expected due to cloudy and drizzling conditions and it is advised to spray the crop and developing bolls with Mancozeb 75 WP + Chlorothalonil 70 WP each @ 2 g/lit of water. In *herbaceum* and *arboreum* cotton crop (Desi varieties), it is suggested to spray the crop with Carbendazim 50 WP @ 1g/lit of water for effective control of cotton grey mildew disease. Seed cotton is to be picked on rain free days wherever the first formed bolls are opened and stored properly. Light irrigation is to be given after each kapas picking wherever irrigation facilities are available. Picking of seed cotton is nearing completion where the crop was sown during the month of May. It is suggested to use the cotton stalks for compost making instead of burning or using as fuel. Alternatively the cotton stalks can be rot slashed in the field itself by tractor operated rotovator.

**Tamil Nadu:** Sowing of cotton is also in progress by utilizing North East Monsoon rainfall. The weather prevailed during the reporting period was moderately cool. The onset of North East Monsoon brought moderate quantity of rainfall in many parts of the State. Sucking pests like aphids, thrips whiteflies and leaf hoppers incidence were noticed in some areas for which need based plant protection measures may be followed depending on the economic threshold level.

#### COTTON WEEKLY ADVISORY TEAM: 2013

Weekly Advisory Report Coordinating Team			
Scientists	Address		
Dr K R Kranthi	Director, CICR, Nagpur		
Dr A H Prakash	PC and Head, CICR, Regional station, Coimbatore		
Dr. D Monga	Head, CICR, Regional station, Sirsa		
Dr. P K Chakrabarty	Head, Div of Crop Improvement, CICR, Nagpur		
Dr Sandhya Kranthi	Head, Div of Crop Protection, CICR, Nagpur		
Dr Blasé De souza	Head, Div of Crop Production, CICR, Nagpur		
Dr. Isabell Agrawal	Sr. Scientist CICR, Coimbatore		
Sh. M.Sabesh	Scientist (SG), CICR, Coimbatore		
Scientists In-charge for Weather Report (AICCIP Centres)			
Scientists	Address	Mobile No	E Mail ID
Dr. Paramajit Singh	Punjab Agricultural University, Bathinda, Punjab	9463628801	rsmeenars@gmail.com
Dr. Pankaj Rathore	Punjab Agricultural University, Faridkot, Punjab	9464051995	pankaj@pau.edu
Dr. Jagdish Beniwal	CCS-Haryana Agricultural University, Hisar 125 004, Haryana	09416325420	cotton@hau.emet.in
Dr.S.L.Ahuja	CCS-Haryana Agricultural University, Sirsa, Haryana	09255947380	slahuja2002@yahoo.com
Dr.K.N.Bhatia	Swami Keshwanand Rajasthan Agricultural University, Sriganganagar, Rajasthan	09352700411	bsmeena1969@rediffmail.com
Dr.Harphool Meena	Maharana Pratap University of Agri. & Technology, Udaipur – 313 001, Rajasthan	09460246043	hpagron@rediffmail.com
Dr. Narendra Kumar	CSA University of Agri. & Technology, Kanpur – 208 002, Uttar Pradesh	09335699132	jagdishk64@yahoo.com
Dr. Gofaldu	Navsari Agricultural University, Navsari – 396 450, Gujarat	09662532645	girishfaldu@rediffmail.com
Dr.M.D.Khanpara	Junagadh Agricultural University, Junagadh – 362 001, Gujarat	09426990070	cotton@jau.in

Dr.R.W.Bharud	Mahatma Phule Krishi Vidyapeeth, Rahuri – 413 722, Maharashtra	09850244087	cotton_mpkv@rediffmail.com
Dr. B . R. Patil	Panjabrao Deshmukh Krishi Vidyapeeth, Akola – 444 104, Maharashtra	09657725801	srscottonpdkv1@yahoo.co.in
Dr.P.R.Zanwar	Marathwada Agricultural University, Parbhani – 431 402, Maharashtra	07588151244	crsned@indiatimes.com
Dr. Satish Parsai	RVS Krishi Vishwa Vidhyalaya, Gwalior – 474 002, Madhya Pradesh	09406677601	aiccipkhandwa@gmail.com
Dr. B.S.Nayak	Orissa University of Agriculture & Technology, Bhubaneshwar – 751 003, Orissa	9437321675	bsnayak2007@rediffmail.com
Dr.S.Bharathi	Acharya N. G. Ranga Agricultural University, LAM, Guntur, AP	0949072341	bharathi_says@yahoo.com
Dr. Sharma	Acharya N. G. Ranga Agricultural University, Nandyal, AP	08514-242296	sharmarars@gmail.com
Dr.Aladakatti	University of Agricultural Sciences, Dharwad – 580 005, Karnataka	09448861040	yralakatti@rediffmail.com
Dr. Bheemana	University of Agricultural Sciences Raichur – 584 102, Karnataka	09448633232	bheemuent@rediffmail.com
Dr. Amala Balu	Tamil Nadu Agricultural University, Srivilliputhur, Tamil Nadu		
Dr. M Gunasekaran	Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu	09443631359	gunasekaran.pbg@gmail.com

=== End of the Report ===