



भा. कृ. अनु. प.- केंद्रीय कपास अनुसंधान संस्थान, नागपुर
ICAR-central Institute for Cotton Research, Nagpur
An ISO 9001:2015 Certified Organisation

Fourteenth Weekly Advisory for Cotton Cultivation from 10th to 16th September, 2026

A. General advisory on soil health, nutrition, pest and disease management

Higher cotton yields rely on balanced soil nutrition, water management, and proactive, monitoring-based pest and disease management. Implementing a holistic and integrated crop management plan ensures proper growth and vigour, reduces input costs, and protects the crop from devastating pest and disease damage.

Soil health & nutrition management (INM)

- **Optimize macronutrients:** Improve soil health by prioritizing crop rotation, organic amendments, and cover crops, and by using rational chemical fertilization based on pre-sowing soil testing.
- **Apply micronutrients:** Address secondary and micronutrient deficiencies (such as Zinc, Boron, Manganese, and Magnesium) proactively via targeted foliar sprays as needed.

Disease management (IDM)

- **Deploy preventative practices:** Management of both soil-borne and foliar pathogens should be achieved through integrated cultural practices, including proper field sanitation and the application of biopesticides and biofertilizers as seed and soil treatments to enhance plant health and suppress disease incidence.
- **Control foliar outbreaks:** Regular field monitoring should be undertaken during periods of high humidity, and recommended fungicides should be applied judiciously only when prevailing environmental conditions are conducive to rapid disease development and spread.

Integrated pest management (IPM)

- **Conduct field scouting:** Perform weekly field scouting to monitor sucking pests (such as thrips, aphids, jassids and spider mites) alongside bollworms.
- **Utilize traps & biopesticides:** Deploy pheromone traps early in the season to track adult moth flights, predict egg-laying cycles and apply organic, natural or plant-based pesticides as a primary response.
- **Conserve natural enemies:** Safeguard beneficial insects like lady beetles, lacewings and big-eyed bugs by choosing highly selective insecticides over broad-spectrum formulas.
- **Restrict chemical interventions:** Use label-recommended chemical pesticides strictly as a last resort, deploying them only when pest populations reach established economic thresholds.

B. Location-specific advisory

PUNJAB		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Firozpur								0	0	0	0	0
	Faridkot	0	0	0	0	0	1	0	0	0.1	0	0	0
	Muktsar								0	0.1	0	0	0
	Bhatinda	0	0	0	0	0	0	0	0	0	0	0	0
	Sangrur								0	0	0	0	0
	Ludhiana	0	0	0	22	0	5.4	0	0	0	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm			64.5 to 115.5 mm			115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall			Heavy rainfall			Very heavy rainfall	

Crop condition

At Bathinda, the crop is 106 to 126 days old and is at boll development stage. Hoeing and weeding operations were taken up. Foliar sprays of 2% Potassium nitrate (13:0:45) were given in cotton fields wherever flowering has started in order to reduce flower drop and improve boll set. Weeds have infested the fields. Whitefly population ranged from 02-15 per three leaves and jassids 01-06 per three leaves. Incidence of green boll damage due to pink bollworm varies from 0 to 30 per cent. With reference to disease incidence, CLCuV was observed in a few fields.

At Faridkot, the crop is 114 to 124 days at boll development to boll maturity stage. Manual hand hoeing to remove leftover weeds from intra row spaces and insecticide sprays were done. Weeds have infested the fields but are controlled with application of recommended herbicides. The whitefly population was either near or above ETL at most of the locations. Jassid population was low to moderate in almost all of the locations with JIG 2-3. Pink bollworm incidence was below ETL at the locations surveyed.

Advisory:

At Bathinda, farmers are advised to apply four sprays of 2% Potassium nitrate (13:0:45) at weekly intervals in cotton fields in order to reduce flower drop and improve boll set. Desi cotton grown in medium to high fertility soils, generally attain unmanageable height and top portion of plants with excessive height usually remain unsprayed. Fruiting bodies of top portions contribute very little towards yield but favour bollworm build up. Detop the plants attaining height more than 1.5 m as and when required by using pruning secateur/sickle. If the whitefly adult population is above ETL, spray Afidopyropen 50 DC @ 400 ml per acre or Flonicamid 50 WG @ 80 g per acre or Clothianidin 50 WG @ 20 g per acre or Dinotefuran 20 SG @ 60 g per acre or Pyriproxyfen 20 EC @ 300 ml/acre or Fenpyroximate 5 EC @ 300 ml/acre. In case of thrips population above ETL, spray the crop with Profenophos 50 EC @ 500 ml/acre or Spinetoram 11.7 SC @ 170 ml per acre. If thrips and PBW population recorded above ETL then, spray Profenophos 50% EC @ 500 ml per acre. Flonicamid 50% WG @ 80 g per acre using 200 litres of water may be sprayed to control whitefly population below ETL. Install pheromone traps @ 2 per acre to monitor the population of PBW. Remove and destroy 'rosette flowers', if any. Spray Emamectin benzoate 5 SG @ 100 g/acre or Spinetoram 11.7 SC @ 170ml/acre using 200 litres of water, if green boll infestation is above 5% (ETL). If incidence of parawilt is observed in the fields, apply drenching with solution of copper oxychloride 50 % WP @ 250 g Or Carbendazim 50% WP @ 100 g plus Urea 2kg/100 litres of water at one-week intervals starting immediately after symptom development or spray Cobalt chloride solution 10 mg/ litre of water immediately after the appearance of symptoms on the affected plants and repeat twice at 5-7-day intervals. Manage the whiteflies with recommended doses of insecticides to check spread of CLCuV. Spray Azoxystrobin 18.2% + Difenoconazole 11.4% SC @ 200 ml in 200 litres of water to control fungal foliar leaf spots of cotton. In case sooty mould develops, apply 1-2 prophylactic/therapeutic sprays of Propiconazole 25%EC @10ml/10 litres (500ml/ha) or Copper oxychloride (COC) 50%WP @ 25g/10 litres (1250g/ha) at 10-15 days interval.

At Faridkot, farmers are advised to start Potassium nitrate (13:0:45) spray @ 2% to improve boll setting and reduce flower drop at 7-10 days interval. For this, dissolve 2 kg of Potassium nitrate in good quality 100 litre water and spray should be very fine so that it gets retained on leaves. If whitefly infestation exceeds ETL, spray Flonicamid 50 WG @ 80 g/acre or Dinotefuran 20 SG @ 60 g/acre. In case of higher infestation of whitefly adults and nymphal population, prefer Ethion 50 EC @ 800ml/acre or Afidopyropen @ 400 ml/acre alternately with Pyriproxyfen @ 500 ml/acre. Nitrogen fertilizer should be applied only after management of whitefly. In case of high thrips infestation apply light irrigation for its management. If the infestation increases, spray Profenofos 50 EC @ 500 ml/acre (in 200 litres of water for one acre). Monitor pink bollworm using funnel or delta traps baited with Gossypure, @ 2 traps per acre, and installed at about 15 cm above the crop canopy. Replace the lure after every 30 days or as per the label specifications. Spray Profenophos 50% EC @ 500 ml per acre or Emamectin benzoate 5SG @ 100 g/acre or Spinetoram 11.7 SC @ 170 ml/acre using 200 litres of water, if green boll infestation is above 5% (ETL). Keep the fields and fallow/barren areas around the farm/canals, etc., free from weeds, which act as collateral hosts of leaf curl virus. After irrigation, some plants might show wilting due to Para wilt which can be checked by spraying Cobalt chloride @ 10mg/ litre of water on the affected plants at the initial stage of wilt symptoms. For management of thrips-transmitted Tobacco Streak Virus (TSV), strictly follow field sanitation, particularly removal of Parthenium weed in and around the cotton fields and recommended management practices for thrips. Keep the fields and fallow/barren area around the farm/ canals etc. free from Kanghi buti (Sida sp.) and Peeli buti (Abutilon sp.) which act as collateral hosts of CLCuD. In case sooty mould develops, go for 1-2 prophylactic/therapeutic sprays of Propiconazole 25%EC @10 ml/10 litres (500ml/ha) Or Copper oxychloride (COC) 50%WP @25g/10 litres (1250g/ha) at 10-15 days interval.

HARYANA		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Hisar	0	0	0	0	0	0	0	0	0.2	0	0	0
	Jind								0	0.1	0	0	0
	Sirsa								0	0	0	0	0
	Rohtak	0.9	0	0	0	0	0	0	0	0.1	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4 mm			
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall			

Crop Condition:

At Hisar, the crop is 105 to 140 days from boll formation to boll opening stage. Manual hoeing, insecticide spray, foliar spray and irrigation (where there were no rains) were taken up during the reporting period. Majority of the fields are infested with weeds. Apply foliar application of 13:00:45 @ 2 kg per acre. Whitefly was found to cross economic threshold in several fields along with honey dew and sooty mould symptoms where control measures were not adopted well in time. Incidence of pink bollworm was noticed in green bolls in several fields near to economic threshold level. Less severity of cotton leaf curl virus disease was noticed in addition to boll rot in some fields. Root rot and wilt were also observed in patches of some fields. Farmers have started picking cotton that got opened due to stress.

At Sirsa, the crop is 114 to 134 days old at boll formation and boll development stages. Irrigation as well as split dose of Nitrogenous fertilizer application, hand hoeing wherever crop canopy is suitable were done. Insecticide spray to manage pests and diseases as well as micronutrient foliar spray were taken up. Weed infestation was noticed at a few locations in the cotton field with poor plant stand. The population of jassids ranged between 03-14/ three leaves, whitefly increasing above ETL between 19-59/ three leaves and thrips above ETL between 01-15 nymphs and adults / three leaves. Pink bollworm infestation was also reported above ETL with 05-15 % green boll damage.

Advisory:

At Hisar, farmers are advised to apply irrigation to cotton crops. Give foliar spray of 13:00:45 @ 2 kg per acre during the peak fruiting stage at 7 to 10 days interval. Need based foliar spray of Urea @ 2.0% and Zinc sulphate 21% @ 0.5% can also

be applied to obtain proper growth. Take up hand hoeing with kasola at appropriate stages. The population of sucking pests is crossing economic thresholds, and there is a need to monitor sucking pests (thrips, whitefly and jassids) on 20 plants/acre. To control jassids and whitefly, spray Flonicamid 50 WG @ 80g per acre and Afidopyropen 50 G/L DC @400ml. In case of moderate to high infestation of whitefly with nymphal population, spray Pyriproxyfen 10 EC @ 400 ml or Spiromesifen 240 SC @240ml per acre. Do not follow tank mixing of insecticides. Since the infestation of pink bollworm started appearing, install pheromone traps @ 2 / acre. Examine at least 100 flowers or 20 green bolls per acre for pink bollworm infestation. Collect and destroy infested and rosette flowers in an early flush of flowering stage. In case of pink bollworm infestation in flower/bolls crosses the economic threshold (>5 %), spray Emamectin benzoate 5SG @100g or Profenofos 50 EC @ 500 ml/acre or Quinalphos 20 AF @ 400 ml per acre. Follow-up spray of an alternate insecticide may be done after 10-12 days of the previous spray. In root rot and wilt affected patches, drench the affected roots with Carbendazim 50 WP@ 1.2 g/litre water. If parawilt is observed, apply 2-3 drenching with solution of Copper oxychloride 50 % WP @ 250g or Carbendazim 50% WP @120g plus Urea 2kg per 100 litres of water at one week interval starting immediately after symptom development. Take up picking of stress mediated good opened bolls after proper drying of moisture. For the management of boll rot, foliar spray of Carbendazim 12%+Mancozeb 63% WP @30g followed by Propineb 70% WP @25-30g or Propiconazole 25% EC @10ml or Azoxystrobin 18.2 % w/w+Difenoconazole 11.4% w/w SC @ 10ml or Fluxapyroxad 167 g/l + Pyraclostrobin 333 g/l SC @ 6 ml in 10 litres of water especially during flowering 60-120 days and as per weather conditions. In case sooty mould develops, apply 1-2 prophylactic/therapeutic sprays of Propiconazole 25%EC@10ml/10L (500ml/ha) Or Copper oxychloride (COC) 50%WP @ 25g/10L (1250g/ha) at 10-15 days interval.

At Sirsa, farmers are advised to irrigate the field wherever required and apply pending split dose of Nitrogenous fertilizer, if any. In case of sudden wilting, yellowing and reddening of leaves occur, apply light irrigation and broadcast 15-20 kg urea per acre. 2-3 foliar spray of NPK (13:00:45) @ 2 kg and MgSO₄ @ 1 kg in 200 litre water per acres and repeat 2-3 times must be applied at weekly intervals. In case of appearance of parawilt symptoms, apply 2-3 drenches with solution of Copper oxychloride 50 % WP @ 250 g Or Carbendazim 50% WP @ 120 g plus Urea 2 kg per 100 litres of water at one week interval starting immediately after symptom development or spray Cobalt chloride @ 10 mg (one gram in 100 litres water) or Sodium benzoate @ 50 mg per litre (5 g in 100 litres water) of water immediately after the appearance of symptoms (within 24 hours) on affected plants only. Keep monitoring whitefly population manually and/or using yellow sticky traps @ 8-10/acre and keep the field bunds and surroundings free from other alternate host plants of whitefly. Continue monitoring for other sucking pests like jassids and thrips by observing 20 plants per acre and from underneath three leaves from each plant. To manage whitefly adults, apply Dinotefuran 20 SG @ 60 g or Flonicamid 50 WG @ 80 g or Afidopyropen 50G/L @ 400 ml or Diafenthiuron 50WP (wherever appropriate humidity exists either due to irrigation or rainfall) @ 200 gm or Profenophos 50 EC @ 500 ml/acre in 200 litres of water. In case sticky, shiny and oily leaves observed with higher whitefly's nymphs, apply Pyriproxyfen 10 EC @500ml or Buprofezin 25 SC @400 ml or Spiromesifen 22.9 SC 240 ml per acre in 200 liters of water 3-5 days after spray to control adults. In the case of sooty mold, spray Propiconazole 25 EC @ 1 ml/litre or Copper oxychloride (COC) 50 WP @2.5 g/litre once or twice at 10-15 days interval. Manage jassid incidence by applying Tolfenpyrad 15 EC @ 300-400 ml or Fenpyroximate 5 EC @ 300 ml or Dinotefuran 20 SG @ 60 g/acre in 200 litres of water wherever the population crosses ETL (2 nymphs/leaf). Install yellow sticky traps @ 8-10 per acre to monitor whitefly population. Initial rowing population of thrips and PBW can be managed by spraying 5% neem seed kernel extract (NSKE) or Azadirachtin 1500 ppm (0.15% EC) @ 5ml/litre of water. Manage pink bollworm population by spraying Emamectin benzoate 5 SG @ 100 g or Spinetoram 11.7 SC @170 ml or Profenophos 50 EC @ 500 ml per acre and use 200 litre water per acre. The spotted bollworm at 5% fruiting body damage (flared square or in shed fruiting bodies) in Desi (arboreum) cotton can be managed by spraying of Spinosad @ 75 ml/acre in 200 litres of water. Under heavy rainfall and waterlogging conditions, maintain proper drainage in the field to remove excess water from the fields to prevent crop damage. Monitor root rot and wilt disease incidence regularly. Drench early symptomatic and surrounding plants with *Trichoderma* spp. 1% WP @ 50 g or Carbendazim 50% WP @ 12 g/ 10 litres of water to manage root-rot and wilt disease.

RAJASTHAN		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Ajmer	0	0	0	0	24	9	0	0	0.1	0.2	0.2	0.1
	Jodhpur	0	0	0	0	0	0	0	0.1	0	0.1	0.2	0
	Nagaur								0	0.1	0.2	0.1	0

	Pali	0	0	0	0	0	0	0	0	0.1	0.1	0.2	0.1
	Sri Ganganagar	0	0	0	0	18.1	0	0	0.1	0	0	0	0
Amount of rainfall & colour Code		0.1 to 2.4 mm	2.5 to 15.5 mm	15.6 to 64.4 mm	64.5 to 115.5 mm	115.6 to 204.4							
Rainfall category		Very light rainfall	Light rainfall	Moderate rainfall	Heavy rainfall	Very heavy rainfall							

Crop Condition:

In Sriganganagar and Hanumangarh districts, the crop is 110 to 126 days old at boll development to boll bursting stage. Manual hoeing/weeding and intercultural operations were taken up. Jassid ranged from 0.0 to 5.00/three leaves, whitefly incidence noticed 0.00 to 19.00/three leaves and thrips population observed ranging from 5.00 to 14.00/three leaves.

At Banswara, the crop is 64 to 70 days old at squaring, flowering and boll formation stages. Fields are clean. Timely spray for sucking pests was given. No infestation of weeds. Among sucking pests, the incidence of whitefly has started appearing in cotton fields but below ETL. Incidence of jassids were above ETL. No incidence of diseases.

Advisory:

In Sriganganagar and Hanumangarh districts, farmers are advised to monitor the crop for insect pests and disease regularly. The locations found infested with pink bollworm last year need to be monitored closely for PBW infestations this year. Install pheromone traps @ 5/ha to monitor pink bollworm activity. In case of recording rosette flowers of more than 5 out of 100 flowers or bolls infestation more than 5%, spray Profenophos 50%EC @2.5 ml/litre or Emamectin benzoate 5 SG@ 0.50 g/litre or Indoxacarb 14.5 SC @ 1.0ml/litre of water. Apply a spray of Flonicamid 50 WG @0.40gm/ litre or Afidopyropen 50 DC @ 1.50 ml/litre of water or Thiamethoxam 24 WG @ 0.5 g/litre of water to control jassids and whitefly while in case of nymph of whitefly, apply Pyriproxyfen 10% EC @2.5 ml/litre or Spiromesifen 22.9 SC @ 1.25 ml/litre of water. While in case of thrips infestation above ETL, spray Spinetoram 11.7 SC @0.84ml/litre or Profenophos 50EC @2.5 ml/ litre of water. In case sooty mould develops, give 1-2 prophylactic/therapeutic sprays of Propiconazole 25%EC @10ml/10L (500ml/ha) Or Copper oxychloride (COC) 50%WP @25g/10 litre of water (1250g/ha) at 10-15 days interval. In case of sudden wilting, yellowing and reddening of leaves occur, apply light irrigation and broadcast 15-20 kg urea per acre. 2-3 foliar spray of NPK (13:00:45) @ 2 kg and MgSO₄ @ 1 kg in 200 litre water per acre and repeat 2-3 times must be applied at weekly intervals. In case of appearance of parawilt symptoms, apply 2-3 drenches with solution of Copper oxychloride 50 % WP @ 250 g Or Carbendazim 50% WP @ 120 g plus Urea 2 kg per 100 litres of water at one week interval starting immediately after symptom development or spray Cobalt chloride @ 10 mg (one gram in 100 litres water) or Sodium benzoate @ 50 mg per litre (5 g in 100 litres water) of water immediately after the appearance of symptoms (within 24 hours) on affected plants only

In southern Rajasthan (Banswara, Bhilwara, Chittorgarh, Dungarpur, Pratapgarh, Rajsamand, Salumber and Udaipur), where manual weeding is not possible in moist soil condition, farmers are advised to go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre water if the field is infested with grassy weeds or Pyriproxyfen sodium 10 % EC @ 1.25 ml/litre water for broad leaved weeds or Pyriproxyfen sodium 6% EC + Quizalofop Ethyl 4% EC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Continue monitoring cotton fields for any signs of sucking pest infestation. Spray 5% neem seed kernel extract (NSKE) or Azadirachtin 1500 ppm (0.15% EC) @ 5 ml/ litre of water to control sucking pests if it goes beyond ETL. Install yellow sticky traps 8-10/acre to manage whitefly and jassids. In case of sucking pests exceeding ETL, spray any one of these insecticides: Buprofezin 25 EC @ 1.25 litre/ha or Diafenthiuron 50 WP @625 g/ha or Flonicamid 50 WG @ 200 g/ha. Install pheromone traps for pink bollworm monitoring (5/ha). Regularly monitor pink bollworm occurrence and destroy the affected flowers (rosette flowers) along with larvae.

ODISHA		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Koraput	0	0	0	0	0	0	0	2	2.3	18	19.7	9.4
	Kalahandi	4	0	0	0.8	0	0	0	1.4	3	24	17.9	7.4
	Balangir	8.2	0	0	0	0	0	0	1.8	4.6	19	17.9	9.8

Amount of rainfall & colour Code	0.1 to 2.4 mm	2.5 to 15.5 mm	15.6 to 64.4 mm	64.5 to 115.5 mm	115.6 to 204.4
Rainfall category	Very light rainfall	Light rainfall	Moderate rainfall	Heavy rainfall	Very heavy rainfall

Crop Condition:

In Odisha, the crop is 64 to 68 days old at flowering to boll formation stage. Drained excess rain water from fields to protect the crop from problem caused by water logging like root rot wilting and reddening of leaves, Application of third dose of fertilizer and spraying of herbicides to control weeds, neem-based pesticide (1500 ppm) for prevention of sucking pests like aphids and other foliage feeders like *Spodoptera* and semi-loopers were taken up during the reporting period. Sprayed 100g MgSO₄ with 100g Urea per 10 litre of water to manage leaf reddening of cotton in places where it occurred due to water logging conditions. All types of weeds, i.e., broad leaf, grasses and sedges, were noticed in most of the cotton fields. Infestation of aphids, jassids, *Spodoptera*, semi-loopers, leaf miners, leaf folders and stem weevil were reported from some areas but below ETL. Wilting and leaf reddening were reported from some places due to continuous rains and waterlogging conditions.

Advisory:

Farmers are advised to maintain proper drainage in cotton fields. Complete intercultural operations, weeding and earthing up operations as and when weather conditions improve. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre water if the field is infested with grassy weeds or Pyrethiobac sodium 10 % EC @ 1.25 ml/litre water for broad leaved weeds or Pyrethiobac sodium 6% EC + Quizalofop Ethyl 4% EC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Regularly monitor the crop to know about the incidence of any pests and diseases. To prevent sucking pests like aphids, jassids and thrips, spray neem-based pesticide @ one litre per acre in 200 litres of water. After 60 DAS of crop stage, spray Buprofezin @ 20 ml/10 litre of water when aphid population is above ETL (≥ 2 affected plants counted randomly showing symptoms cupping / crumpling of few leaves on the upper portion of plant (10% plants infested). To monitor pink bollworm and *Spodoptera* infestations, install pheromone traps baited with Gossypure and Spodpure, respectively @ 5 traps /ha. If parawilt or sudden wilt occurs due to heavy rain and waterlogging after a dry spell, spray Cobalt chloride @10 mg/litre (10ppm) or 2-3 drenching with solutions of Copper oxychloride 50 % WP @ 250 g Or Carbendazim 50% WP @ 100 g plus Urea 2 kg per 100 litres of water at one-week interval starting immediately after symptom development on affected plants. Spray the crop with 100g Magnesium sulphate mixed with 100g Urea in 10 litres of water in waterlogged areas to reduce leaf reddening conditions. Drench early symptomatic and surrounding plants with *Trichoderma* spp. 1% WP @50 g or Carbendazim50% WP @ 12 g/ 10 litres of water to manage root-rot and wilt diseases.

GUJARAT		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Amreli	0	0	0	0	0	0	0	0.1	0	0.3	0.1	0.8
	Bhavnagar	0	0	3	0	2.2	2	1	0.1	0.1	0.4	0.3	9.6
	Jamnagar	0	1	0	0	0	0	0	0.4	0.8	1.2	0.6	0.2
	Rajkot	0	0	0	0	0	0	0	0.2	0.6	0.8	0.5	1
	Junagadh	1.8	0	0	0	0	0	0	0.9	1.1	0.9	1.1	1.8
	Sabarkantha								0	0.1	0.3	0.1	2
	Surendranagar	0	0	0	0	0	0	0	0	0.2	0.4	0.4	1.4
	Ahmedabad	0	0	0	0	0	0	0	0.1	0.5	0.3	0.2	3.6

	Baroda	0	0	0	0	0	0	0	0.2	0.3	0.4	0.2	10.9
	Patan								0.1	0.2	0.6	0.2	0.1
	Mehesana								0	0.3	0.4	0.2	1.2
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4			
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall			

Crop Condition:

At Surat, the sown crop is 52 to 57 days old at vegetative growth stage. Due to deficit rains, most of the plants are in stress conditions. Weeding, fertilizer application and intercultural operations were taken up during the reporting period. Weeds like Chido (*Cyprus rotundus*), Satodi (*Trianthema monogyna*), Dhamdo (*Amaranthus viridis*) and others have infested the fields. Thrips incidence was noticed above ETL. Bacterial Leaf Blight disease has been recorded in few fields.

At Junagadh, the sown crop is 63 days old at square formation and flowering stage. Moisture stress was observed due to which parawilt was observed in isolated locations. Intercultural operations, weeding and insecticide spray were done during the reporting period. Kharif weeds (*Echinochloa colona*, *Cyperus rotundus*, and *Setaria glauca*) have infested the cotton fields. Amidst prevalence of dry spell, earthing up and irrigation are under progress. Incidence of thrips was noticed above ETL at most of locations.

Advisory:

At Surat, farmers are advised to carry out field sanitation. In case of heavy rain, drain out the excess rainwater from cotton fields. Cotton crop requires 240-40-00 NPK/ ha. Dose of Nitrogen is applied after sowing @ 30, 60, 75, 90 and 105 days in equal splits. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre water if the field is infested with grassy weeds or Pyriothobac sodium 10 % EC @ 1.25 ml/litre water for broad leaved weeds or Pyriothobac sodium 6% EC + Quizalofop Ethyl 4% EC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. While in case of thrips infestation above ETL, spray Spinetoram 11.7 SC @0.84 ml/litre or Profenophos 50 EC @ 2.5 ml/ litre of water. Check Fusarium wilt patches along with one-meter radius of healthy plants in desi cotton fields with suitable fungicide application. For the control of wilt and root rot disease, give spot application of Carbendazim 50% WP@ 1.2 g per litre of water or *Trichoderma harzianum* or *T. viride* 1% WP@10 g/ litre of water. If parawilt is observed, apply 2-3 drenchings with solution of Copper oxychloride 50 % WP @ 250 g Or Carbendazim 50% WP@ 100 g plus Urea 2 kg per 100 litres of water at one-week interval starting immediately after symptom development. On receiving sufficient rain, apply second instalment of split dose of nitrogenous fertilizer application. Complete the dose of Phosphorous, if not applied as a basal dose.

At Junagadh, farmers are advised to remove excess water from the cotton fields. Give top dressing of Ammonium sulphate 10 kg per acre or Urea @15 kg/acre and Potash @25kg/acre along with irrigation. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5 % EC @ 2 ml /litre water if the field is infested with grassy weeds or Pyriothobac sodium 10 % EC @ 1.25 ml/litre water for broad leaved weeds or Pyriothobac sodium 6% EC + Quizalofop Ethyl 4% EC @ 2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. While in case of thrips infestation above ETL, spray Spinetoram 11.7 SC @0.84 ml/litre or Profenophos 50 EC @ 2.5 ml/ litre of water. In case of jassid exceeding ETL, spray any one of these insecticides: Buperfezin 25 EC @ 1.25 litre/ha or Diafenthiuron 50WP @ 625 g/ha or Flonicamid 50 WG @200 g/ha. Manage parawilt damage by spraying water soluble fertilizer (13-0-45, NP-K) @ 100 g in 10 litres water and drench the affected plants 2-3 times at weekly interval with Copper oxychloride 50 WP@ 25 g or drench with solution of Carbendazim 50 WP@100 g plus Urea 2 kg/100 liter of water immediately after symptom appearance. For the management of wet weather blight and boll rot, foliar sprays of (Carbendazim 12%+Mancozeb 63% WP) @ 25 g or Propiconazole 25% EC @10 ml or (Azoxystrobin 18.2% w/w+Difenoconazole 11.4% w/w SC) @ 10 ml or (Pydiflumetofen 13.6% w/w + Propiconazole 11.4% w/w SE) @12 ml or (Fluxapyroxad 167 g/l + Pyraclostrobin 333 g/l SC) @ 6 ml in 10 litres of water is recommended.

MADHYA PRADESH	Actual Rainfall in last week(mm)	Predicted Rainfall in next week (mm)
	September	September

		01	02	03	04	05	06	07	09	10	11	12	13
	Kharggaon	0	0	0	0	0	0	0	4.1	0.4	0.2	10.8	13.6
	Dhar	0	0	0	15.9	4	0.2	0.5	1.8	0.4	0.4	5.2	7.4
	Khandwa	0	0	0	0	0	0	0	2.6	3.9	4.2	10.9	10.5
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm				64.5 to 115.5 mm		115.6 to 204.4	
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall				Heavy rainfall		Very heavy rainfall	

Crop Condition:

At Khandwa, the sown crop is 56 to 104 days old in its flowering growth stage. Spot weeding, fertigation, drip irrigation was given wherever required. Seasonal weeds like *Cynodon dactylon*, *Cyperus rotundus*, *Argemone maxicana* and *Phyllanthus niruri* have infested the fields. Incidence of jassids, aphids and whiteflies have been observed in some fields. No incidence of diseases reported so far.

Advisory

At Khandwa, farmers are advised to take up weeding on time. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4%MEC @2-2.5 ml /litre of water to control both grassy and broadleaved weeds. Install pheromone traps @2 / acre to monitor pink bollworm. Give a split dose of Nitrogen @ 25% applied by column method at the depth of 10 to 15 cm. Apply NSKE 5%+ Neem oil 5 ml/litre or neem oil-based formulations (300 or 1500 ppm) 5 ml/litre + 1.0 g laundry detergent emulsion to control sucking pests. If parawilt is observed, apply 2-3 drenching with solution of Copper oxychloride 50 % WP @ 250 g Or Carbendazim 50% WP @ 100 g plus Urea 2 kg per 100 litres of water at one-week interval starting immediately after symptom development. In case of a long dry spell, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water during the peak fruiting stage at 7 to 10 days interval. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton.

MAHARASHTRA		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Dhule								4	0.5	0.2	7	32.9
	Nandurbar								3.6	0.5	0.3	2.8	28.7
	Jalgaon	0	0	0	0	0	0	0	4.7	0.7	0.3	10.7	28.6
	Ahilyanagar	0	0	0	0.1	0	0	0	2.1	0.8	0.1	3.5	28
	Ch. Sambhaji Nagar	0.2	0.2	0	0	0	0	0	3.2	0.6	0.2	7.9	37.4
	Jalna	0	0	0	0	0	0	0	2.6	0.6	0.2	9.1	35.3
	Beed								1.2	0.4	0.2	8.1	29.6

	Nanded								0.8	0.8	1.9	18.3	20.7
	Parbhani	0	0.3	0	0	0	0	0	0.6	0.6	0.3	11.4	25.8
	Hingoli								1	0.7	0.5	12	22.1
	Buldhana	2	0	3	0	9	0	0	3.2	0.7	0.1	9	25.8
	Akola	0	0	0	0	0.8	0.7	0	2.2	0.5	0.3	14	20.7
	Washim	0	0	0	0	0	0	0	4.2	0.8	0.6	16.5	18.3
	Amravati	0	0	1	0	1	0.4	0	3.8	0.6	1.7	10.6	12.6
	Yavatmal	0	0	0	1	0	0	0	1.5	0.7	3.3	23.2	18.6
	Wardha	0	1	7	2.2	0	0	0	2.8	1	5.6	15.7	12.2
	Nagpur	0.2	0	7	2.4	4	0	0	3.8	1.2	6.4	14.7	15
Chandrapur	0	0	20	8	0	1	0	1.8	1.5	10	19.6	17.4	
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4			
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall			

Crop Condition:

At Akola, the sown crop is 71 to 76 days old at flowering and boll development stages. Hoeing and weeding operations were taken up. Insecticides/ fungicides/ micronutrient sprays were done. A long dry spell was experienced during the reporting week resulting in severe incidence of *Fusarium* wilt and *Macrophomina* dry root rot diseases in some areas. The crop growth is stunted due to moisture stress. Sucking pests like whitefly and thrips were noticed in some fields.

At Rahuri, the sown irrigated cotton crop is 64 to 99 days at flowering, boll formation and boll development stages based on the time sown. Irrigation, hoeing, weeding, spraying and fertilizer application were done. Weeds have infested the fields. Thrips and jassid incidence have crossed ETL in many fields with aphid incidence below ETL. Pink bollworm was recorded in rosette flowers. *Myloccerus* weevil has damaged 01% cotton plants. TSV has been recorded in many fields.

At Nanded, the sown crop is 60 to 85 days at square formation to boll formation stage. Intercultural operations, weeding, fertilizer application, plant protection and growth retardant sprays were done during the reporting period. Grassy, broad-leaved weeds and sedges have infested the fields. Jassids and aphids incidence were noticed but below ETL. No incidence of diseases.

Advisory:

At Akola, farmers are advised to open the furrows in the cotton field for moisture conservation purposes at 45-50 DAS. Carry out intercultural operations like hoeing and weeding in cotton. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + QuizalofopEthyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Apply first split of 40 Kg Nitrogen (90Kg Urea per ha) for irrigated hybrid cotton and 45 Kg N (97 Kg urea per ha) for rainfed hybrid/*hirsutum* cotton as top-dressing dose of chemical fertilizer. Apply NSKE 5% + Neem oil 5 ml/litre or neem oil-based formulations (300 or 1500 ppm) 5 ml/litre + 1.0 gm laundry detergent emulsion to control sucking pests like aphids, jassids and thrips, and also to repel pink bollworm females from ovipositing on cotton squares. To monitor pink bollworm, install 2 pheromone traps per acre for

surveillance purposes in the field. Observe 20 plants per acre as a representative of the cotton field for pest infestation. If parawilt is observed, apply 2-3 drenching with solution of Copper oxychloride 50 % WP @ 250 g or Carbendazim 50% WP @ 100 g plus Urea 2 kg per 100 litres of water at one-week interval starting immediately after symptom development. In *Fusarium* wilt and *Macrophomina* root rot affected areas, farmers are advised to uproot the infected plants and destroy them immediately to prevent the spread of disease. Drench the surrounding plants with Carbendazim 50% WP @ 1.2 g/L of water to prevent further spread. This should be followed by two drenches of *Trichoderma harzianum* or *T. viride* @ 5 g/litre of water at 15-day intervals. Farmers who adopted High-Density Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. If the crop is suffering from moisture stress due to prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton.

At Rahuri, farmers are advised to apply second split dose at 60 DAS. Keep the crop weed-free by hand hoeing or intercultural operations with a tractor or bullock operated implements. Remove *Parthenium* weed from the field and field bunds so as to avoid or to minimize TSV and thrips incidence. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriithobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriithobac sodium 6% + Quizalofop Ethyl 4%MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Monitor the fields to keep a check on the infestation of sucking pests. Install yellow sticky traps, spray homemade neem seed kernel extract (NSKE) 5% or Azadirachtin 10000 ppm 10 ml/10 litres of water. In cotton crops above 60 DAS, if sucking pests cross ETL, spray Flonicamid 50 WG 2g/10 lit water or Buprofezin 25 SC @ 20 ml /10 litres of water. Install pheromone traps @ 5 /ha to monitor pink bollworm and collect and destroy the rosette flowers along with PBW larvae at regular intervals to avoid further generation of pink bollworm. Farmers who adopted High-Density Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. If the crop is suffering from moisture stress due to prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton. Farmers having closer spacing (90 x 30 cm) cotton but not willing to spray PGR can go for removal of monopodia at 45 days after sowing and detopping at 75 to 80 days. If parawilt is observed, apply 2-3 drenching with solution of Copper oxychloride 50 %WP @ 250 g Or Carbendazim 50% WP @ 100 g plus Urea 2 kg per 100 litres of water at one week interval starting immediately after symptom development. Monitor the crop to avoid transmission of Tobacco Streak Virus (TSV) disease. Install blue sticky traps @20/ha to monitor and 100/ha for management of thrips. Uproot and destroy TSV infected plants to avoid spread of the disease and strictly follow field sanitation, particularly removal of *Parthenium* weed in and around the cotton fields and recommended management practices for thrips. Manage thrips (>ETL) incidence with recommended use of insecticides.

At Nanded, farmers are advised to take up intercultural operations and weeding to manage weeds. Give protective irrigation as per availability under moisture stress situation. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriithobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriithobac sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Give top dressing of Nitrogen @ 60 kg per ha to irrigated crops at 60 DAS whereas Nitrogen @ 48 kg/ha at 30 DAS to rainfed crop by drilling or ring method. Avoid broadcasting urea. Spray growth retardant, Mepiquat Chloride 5 % @ 500 ml/ha, at square formation stage in *Bt* cotton planted under HDPS and closer spacing cotton fields. Give foliar spray of NSKE 5% to manage sucking pests at 30 DAS. In crop of > 60 DAS, go for foliar application of Buprofezin 20 SC @ 1000 ml /ha or Dinotefuron 20 SG @ 150 g per ha to manage sucking pests once it crosses ETL. Field and field bunds should be kept weed free and *Parthenium* weed should be removed in and around the fields to minimize the thrips population build-up and TSV incidence. If parawilt is observed, apply 2-3 drenching with solution of Copper oxychloride 50 % WP @ 250 g or Carbendazim 50%WP @ 100 g plus Urea 2 kg per 100 litres of water at one-week interval starting immediately after symptom development. For the management of leaf spots and boll rot, foliar sprays of (Carbendazim 12%+Mancozeb 63% WP) @ 30 g or Propiconazole 25% EC @10 ml or (Azoxystrobin 18.2% w/w+Difenoconazole 11.4% w/w SC) @ 10 ml or (Fluxapyroxad 167g/l + Pyraclostrobin 333 g/l SC) @ 6 ml in 10 litres of water is recommended. Farmers who adopted High-Density Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. If the crop is suffering from moisture stress due to

prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4ml /10 lit of water to avoid natural shedding of squares and flowers of cotton.

TELANGANA		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Adilabad	0	0	2	0	0	0	0	2	2.2	6.8	28.7	16.5
	Warangal	0	4	3	7	0	0	0	1.4	0.4	9	23.3	8.2
	Khammam	0	0	2	0	0	0	0.6	2.4	0.3	5.5	11.9	5.8
	Karimnagar	0	0	1.7	9.6	0.3	0	0	0.5	1.3	6.5	24.4	8.5
	Mahabubnagar	0	0	0	0	0	0	0	0.2	0.8	3	5	1.3
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm			64.5 to 115.5 mm		115.6 to 204.4 mm		
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall			Heavy rainfall		Very heavy rainfall		

Crop Condition:

At Adilabad, the sown crop is at 67 to 72 days at flowering to boll formation stage. Inter cultural operations, weeding, insecticide and foliar sprays were given during the reporting period. Thrips and jassid incidence were observed in most of the fields above ETL. No incidence of diseases

At Warangal, the crop is 77 to 88 days old at flowering to boll formation stage. Inter-cultivation with Bullock drawn Danti or Guntaka, top dressing with Urea and Potash and insecticide sprays to control sucking pests were carried out during the reporting period. Weeds have infested the fields. Thrips, aphids and jassids were noticed above ETL. No incidence of disease.

Advisory:

At Adilabad, farmers are advised to take up periodical hoeing for moisture conservation and weed control. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2 ml/litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broadleaved weeds. Give prophylactic spray of neem based insecticide @ 5 ml per litre of water at 45 DAS to control sucking pests and deterring pink bollworm females from egg laying on newly formed squares. If the population of sucking pests exceeds ETL, give a spray of Flonicamid 50 WG @ 0.4 g per litre of water. Farmers who adopted High-Density Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0ml/litre of water. If the crop is suffering from moisture stress due to prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton. Monitor the crop to avoid transmission of Tobacco Streak Virus (TSV) disease. Install blue sticky traps @20/ha to monitor and 100/ha for management of thrips. Uproot and destroy TSV infected plants to avoid spread of the disease and strictly follow field sanitation, particularly removal of *Parthenium* weed in and around the cotton fields and recommended management practices for thrips. Manage thrips (>ETL) incidence with recommended use of insecticides.

At Warangal, farmers are advised to give top dressing of Nitrogen (25 kgs of Urea/acre) and Potassic (10 kgs MOP/acre) fertilizers at 40,60 and 80 DAS after receipt of rains in rainfed crops with sufficient moisture. With respect to weed control, in case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5%EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad

leaved weeds or Pyrethrin sodium 6% + Quinalphos Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Prophylactic spray of neem based insecticide @ 5 ml per litre of water at 45 DAS may be given for controlling sucking pests and deterring pink bollworm females from egg laying on newly formed squares. If population of sucking pests exceeds ETL, spray Flonicamid 50 WG @ 0.4 or Diafenthiuron 50 WP @ 1.25 g or Dinotefuron 20 SG @ 0.4g per litre of water from 60 DAS onwards. Install yellow- and blue-colored sticky traps (10 each per acre) as a preventive measure to monitor and control the sucking pest complex. Dry conditions are favorable for the infestation of pink bollworm. So, monitor the infestation of the pest by installing pheromone traps (2 No./Acre) in the cotton fields. Farmers who adopted High-Density Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. The crop is suffering from moisture stress due to prolonged dry spell. So, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. Go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of flowers of cotton. Monitor the crop to avoid transmission of Tobacco Streak Virus (TSV) disease. Install blue sticky traps @20/ha to monitor and 100/ha for management of thrips. Uproot and destroy TSV infected plants to avoid spread of the disease and strictly follow field sanitation, particularly removal of *Parthenium* weed in and around the cotton fields and recommended management practices for thrips. Manage thrips (>ETL) incidence with recommended use of insecticides.

ANDHRA PRADESH		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Guntur	0	7	0	0	0	0	14	4.2	0.3	5.3	7.6	4.2
	Prakasam	0	0	0.2	0.5	0	0	0	1.2	0.6	5.2	5.7	3
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm			64.5 to 115.5 mm		115.6 to 204.4 mm		
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall			Heavy rainfall		Very heavy rainfall		

Crop Condition:

At Guntur, sowings have been completed and also suggested not to take up further sowings. Intercultivation and weeding operations were completed in the cotton fields. In the present scenario, deficit rains have led to drying and withering of crop in majority of the fields. Incidence of sucking pests viz., thrips and leafhoppers were above ETL.

At Nandyal, the sown crop is 59 to 89 days old at squaring and flowering stages. Nearly 85 per cent of cotton area in Kurnool district was sown during June first fortnight under less moisture conditions wherein majority of the areas, crop is exhibiting wilting and moisture stress symptoms. Top dressing of fertilizer application and intercultural operations were taken up during the reporting period. Infestation of sucking pests were recorded around ETL in most of the locations. Pink bollworm infestation was also noticed in few fields

Advisory

At Guntur, farmers are suggested to not to take up further sowings. Give basal application of P_2O_5 @ 60 kg/acre and 50 kg N and 20 Kg K_2O per acre within 30 DAS Use Pendimethalin 38.7 CS @ 700 ml/acre in 200 litres of water as pre-emergence weedicide to control weeds in the early stage of the crop. Give foliar spray with 19:19:19 or 13:0:45 @ 1% per liter of water. Spray Thiamethoxam 25% WG @ 0.2g/lit or Fipronil 5% SC @ 2.0 ml/lit of water to control sucking pests. Monitor the crop to avoid transmission of Tobacco Streak Virus (TSV) disease. Install blue sticky traps @20/ha to monitor and 100/ha for management of thrips. Uproot and destroy TSV infected plants to avoid spread of the disease and strictly follow field sanitation, particularly removal of *Parthenium* weed in and around the cotton fields and recommended management practices for thrips. As a prophylactic measure against sucking pests and to check the initial pink bollworm infestation, spray neem-based insecticides at 1 litre per acre using 200 litres of water.

At Nandyal, farmers are advised to go for a spray of Urea or 13-0-45 @ 10 g/lit of water to tide over temporary dry spells. To manage sucking pests, spray neem oil @ 1 litre per acre (in 200 litres of water. Once it goes beyond ETL, give a spray of Imidacloprid @ 0.4 ml or Acetamiprid @ 0.2g per lit of water. Remove weeds on the bunds to avoid transmission of TSV virus. If pink bollworm crosses ETL, spray Profenophos @500 ml/acre using 200 litres of water. Farmers who adopted High-Density

Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. If the crop is suffering from moisture stress due to prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton. Monitor the crop to avoid transmission of Tobacco Streak Virus (TSV) disease. Install blue sticky traps @20/ha to monitor and 100/ha for management of thrips. Uproot and destroy TSV infected plants to avoid spread of the disease and strictly follow field sanitation, particularly removal of *Parthenium* weed in and around the cotton fields and recommended management practices for thrips. Manage thrips (>ETL) incidence with recommended use of insecticides.

KARNATAKA		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Dharwad	6	3	2	0	2	0	0	1.7	0.6	0.6	0.2	0.4
	Haveri	0	0.4	0	0	0	0	0	2.3	0.9	1	1.2	0.5
	Mysore	0	0	0	0	0	0	0	1	1	2.9	5.7	3.4
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm			64.5 to 115.5 mm		115.6 to 204.4 mm		
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall			Heavy rainfall		Very heavy rainfall		

Crop Condition:

At Dharwad, the crop is 71 to 76 days at flowering to boll formation stage. Hand weeding, inter cultivation for weed management, insecticides and foliar spray were taken up. Weeds infestation was noticed in certain fields. Shoot weevil, leaf hopper and aphids have infested the fields in low status but thrips in moderate status. *Alternaria* blight disease was reported from low to moderate status in certain fields.

At Raichur, the sown crop is 73 days old at squaring to flowering stage. Non receipt of rainfall is resulting in withering of cotton plants with various magnitudes, though the early sown crop is withstanding drought. Overall, 25-30 % of the cotton area is under irrigation. Inter cultivation operation has been taken up along with sprays for sucking pest control. Thrips incidence was reported above ETL and more influenced by lack of rainfall.

At Chamarajanagar, the crop is 100 to 115 days from boll development to boll opening stage. Spraying work is in progress. Incidence of leaf hopper – (04-06/3 leaves), aphids – (25-40/3 leaves) and PBW incidence below 5% were recorded. No incidence of diseases.

Advisory:

At Dharwad, farmers are advised to take up hand weeding and intercultural operations to manage weeds. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5% EC @ 2ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds. Do hand collection of shoot weevils from the okra as a trap crop. Apply Phosphorous and Potash by ring method. Apply Salicylic Acid @100 ppm spray for 75 days old crop. Spray neem based insecticide @ 5.0 ml per litre of water to manage sucking pests. After 60 DAS of crop age, spray Flonicamid 50 WG @0.4g per litre or Diafenthiuron 50 WP @ 1.2 g per litre of water to manage sucking pests if crosses ETL. Install pheromone traps @ 2 /acre to monitor pink bollworm moth activity. Destroy rosette flowers with PBW larvae. Farmers who adopted High-Density Planting System (90 x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. If the crop is suffering from moisture stress due to prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton.

At Raichur, farmers are advised to take up weeding and intercultivation on priority basis to avoid the land getting dried up. In case of moist soil condition where manual weeding is not possible, go for application of herbicides like Quizalofop ethyl 5%EC @ 2 ml /litre of water if the field is infested with grassy weeds or Pyriothobac sodium 10% EC @1.5 ml/litre of water for broad leaved weeds or Pyriothobac sodium 6% + Quizalofop Ethyl 4% MEC @2-2.5 ml /litre of water to control both grassy and broad-leaved weeds when the crop is at 25-30 days duration. Spray the recommended dose of Flonicamid 50% WG (4g/ 10 litres of water) against sucking pest populations that exceed ETL. Spray Profenophos 50 EC @ 2.5 ml or Imidacloprid 17.8 SL @ 0.3 ml per litre of water to control mealy bug infestation. Farmers who adopted High-Density Planting System (90x 15 cm) or closer spacing (90 x 30 cm) cotton have to go for spray of plant growth regulator (PGR) Mepiquat chloride 5% AS @1 ml/litres of water at 40–45-day crop or at square initiation and second spray after 15-20 days @1.0 ml/litre of water. If the crop is suffering from moisture stress due to prolonged dry spell, avoid spray of Mepiquat chloride. Alternatively, give foliar spray of 13:00:45 @ 2 kg per acre in 200 litres of water. If square drop is observed, go for foliar spray of NAA @ 4 ml /10 lit of water to avoid natural shedding of squares and flowers of cotton.

At Chamarajanagar, farmers are advised to install yellow sticky traps @ 8-12 per acre and pheromone traps, change of lures @ 15 days interval to monitor sucking pests. Spray Flonicamid 50% WG 0.4 ml /lt of water or Dinotefuran 20 SG @ 0.3 gm/lt of water and Imidacloprid 17.8 SL @ 0.5 ml /lt of water to control sucking pests above ETL. Spray Profenophos 50 EC @ 500 ml per acre or Emamectin benzoate 5SG @100 gm per acre or Indoxacarb 15 EC @ 200 ml per acre, if the incidence of pink bollworm is above 5 per cent (ETL) in flowers or green bolls. If the crop is above 120 days and incidence of pink bollworm is above ETL, spray Cypermethrin 10% EC @ 10-15 ml/10 lit of water or Deltamethrin 2.8 EC @ 10 ml/10 lit of water or Fenprothrin 10% EC @ 10 ml/10 lit of water.

TAMIL NADU		Actual Rainfall in last week(mm)							Predicted Rainfall in next week (mm)				
		September							September				
		01	02	03	04	05	06	07	09	10	11	12	13
	Perambalur	0	0	0	0	0	68.6	0	0.2	0.6	7.8	11.1	10
	Salem	0	0	5	0	0.4	0	0	0.2	1.2	1.9	8.6	4.8
	Trichy								0.2	0.3	2.7	7.7	12.6
	Virudhunagar	0	0	0	7	29.4	0	0	0.1	0.6	1.7	1.2	1.2
Amount of rainfall & colour Code		0.1 to 2.4 mm		2.5 to 15.5 mm		15.6 to 64.4 mm		64.5 to 115.5 mm		115.6 to 204.4 mm			
Rainfall category		Very light rainfall		Light rainfall		Moderate rainfall		Heavy rainfall		Very heavy rainfall			

Crop Condition:

At Coimbatore and surrounding areas, sowing has been initiated. The sown crop is at vegetative stage. Pre-emergence application of Pendimethalin (38.7% CS) @ 260 ml/ac or Pendimethalin (30%EC) @ 400 ml/ac was practiced within three DAS as recommended. During field preparation, soil application of *Trichoderma viride* and *Bacillus subtilis* @ 2.5 kg/ha along with 500 kg well decomposed FYM was carried out. During sowing, delinted seeds were treated with talc formulation of *Trichoderma viride* @ 4g/kg of seed or Thiram 75 WP@ 2.5 g/kg of seed to manage soil borne diseases in case of Non-Bt seeds. Leaf miner incidence has been noticed and initiated. prophylactic spray of neem oil 3% as recommended. Root rot incidence has been noticed.

At Srivilliputhur and surrounding areas, field preparation is in progress. Sowing is in progress. The sown crop is 10 to 20 days old at initial growth stage. Applied pre-emergence herbicide and still in progress in some places wherever sufficient moisture is available in the soil at the time of herbicide application. Weed species found were *Cyprus* spp and *Cynadon dactylon* and *Trianthema portulacastrum* etc. No incidence of pests and diseases.

Advisory

At Coimbatore and surrounding areas, farmers are advised to take up kharif sowing at the earliest possible. Sow the crop after receipt of at least 80-100 mm rainfall to avoid germination-related problems. Regulate irrigation based on soil type and receipt of rainfall. Give seed treatment with Carboxin 37.5% + Thiram 37.5% DS @ 3.5 g per kg of seeds (root rot and bacterial blight disease) or Fluxapyroxad (333 g/lit FS) @ 1.5 ml/kg seed or Tetraconazole 11.6% w/w (12.5% w/v) SL @ 1.5 ml/kg of seeds for seed-borne diseases (for root rot disease) as well as biofertilizers, viz., *Azotobacter*, PSB and KMB @ 10 ml per kg seed. Clean the bunds and irrigation channels and maintain them properly to ensure drainage during excess rains. Give pre-emergence application of Pendimethalin (38.7% CS) @ 700 ml/ac or Pendimethalin (30%EC) @ 1000 ml/ac within two DAS. Take up gap filling on the tenth DAS. Regularly monitor the crop for incidence of pests and diseases to adopt timely management. Select medium-duration hybrids.

At Srivilliputhur, isolated rainfall received in some parts of the district. Hence, farmers are advised to utilize the available moisture for herbicide application and sowing of crop. Remove cotton stalks after harvest to prevent the carry over life stages of stem weevil. Apply neem cake @100kg per acre during the last plough. Clean the farm bunds and other premises to destroy the alternate host of insect pests such as *Abutilon* and other grassy weeds. Do acid delinting of cotton to eliminate some externally seed borne pathogenic organisms and to kills eggs, larvae and pupae of seed harbouring insect pests and it also helps to remove immature, ill-filled, cut and damaged seeds. In stem weevil endemic areas, give seed treatment with Chlorpyrifos 20EC @ 10ml/kg of seed + drenching collar region with Chlorpyrifos 50 EC @ 1200 ml/ha on 15 DAS.

The detailed information regarding cotton production technology, e.g., selection of soil, varieties, fertilizer application, sowing methods, irrigation systems, management of weeds, insect pests and diseases, etc., can be availed from an Android-based **CICR Cotton App** developed by ICAR-CICR, Nagpur. The app can be downloaded free of cost from Google Play Store. Additionally, the crop growth stage-specific and weather-based weekly advisories are uploaded on the website of ICAR-CICR (<https://cicr.org.in/resource-weekly-advisory>) also to be consulted for the benefit of farmers.

Advisory issued by:

Dr. Vijay N. Waghmare

Director, ICAR-Central Institute for Cotton Research, Nagpur

Compilation by Dr. Isabella Agarwal

Editing Team:

Dr. G. T. Behere	Dr. A. S. Tayade	Dr. A. H. Prakash	Dr. S.K. Verma
Dr. S.K. Sain	Dr. Rishi Kumar	Dr. Babasaheb B. Fand	Dr. Ramkrushna GI
Dr. S.P Gawande	Dr. Shivaj Thube	Dr. R.R. Chapke	

Regional language translation:

Hindi	: Mr Ghanshyam Deogirikar & Mr Ashutosh Mishra
Punjabi	: Dr Amarpreet Singh
Tamil	: Dr. A. Manikandan & Dr. M Saravanan
Telugu	: Dr L. Rajesh Chaudhary
Marathi	: Dr Vrushali Deshmukh & Mrs Pooja Ghonge
Gujrati	: Dr H R. Desai & Dr Vivek Shah
Oriya	: Dr B. S. Nayak
Kannada	: Dr Neelkanth Hiremani